

REFERNCES

- Abbas, M.S.T. and Y.A.A. El-Deeb (1993):** On the natural enemies of the major pests infesting cotton in Egypt. Egypt, J. Agric. Res. 71 (1): 131-138.
- Abbott, W.S.(1925):** A method of computing the effectiveness of an insecticide. J. Econ. Entomol.. 18: 265 - 277.
- Abd El-Megeed, M.I. ; El-Refaei, S.A.; Zidan, A.A. and W.M. Watson (1984-1985):** The joint action of microbial and chemical insecticide combinations on the cotton leafworm, *Spodoptera littoralis* (Boisd.). Bull ent. Soc. Egypt, Econ. Ser., 14 : 119 - 126.
- Abd El-Megeed, M.I. ; Zidan, Z.H.; Watson, M.W.; Hegazy, G.M and M.S. Zemaity (1986):** The influence of IGR / insecticide mixture on the biotic potential of the cotton leafworm, *Spodoptera littoralis* (Boisd.). Med. Fac. Landbouww. Rijksuniv. Gent. 51 (1): 137 - 143.
- Abou – Zeid, N. A. (1972):** Studies of the rove beetle *Paedrus alfieri* Kok (Coleoptera : Staphylinidae). Ph.D. Thesis, Fac. of Agric. Cairo University.
- Abul-Nasr, S. and S.T. El-Sherif (1973 a):** Seasonal fluctuations of the egg-masses of the cotton leafworm, *Spodoptera littoralis* (Boisd.) in cotton fields. Bull. Soc. ent. Egypte, 57 : 353 - 360.

- Abul-Nasr, S. and S.T. El-Sherif (1973 b):** Seasonal fluctuations of moths of the cotton leafworm, *Spodoptera littoralis* (Boisd.) during the cotton season. Bull. Soc. ent. Egypte, 57: 413 - 418.
- Abul-Nasr, S.E.; Ammar, E.D. and S.M. Farrag (1979):** Rates of infestation by *Pectinophora gossypiella* (Saund.) and *Earias insulana* (Boisd.), (Lepidoptera), on flowering sites of the cotton plant. Deutsche Entomologische Zeitschrift. 26 (1/3): 165 - 172.
- Ahmad, S.; O'Neill, J.R.; Magur, D.L. and R.K. Nowalk (1978):** Toxicity of *Bacillus thuringiensis* to gypsy moth larvae parasitized by *Apanteles melanoscelus*. Environ. Entomol. 7 (1): 73 - 76.
- Ahmed, S. M. ; Shalaby, F. F. and M. K. El- Khishin (1978):** Antifeedant properties of some plant extracts for the larvae of *Spodoptera littoralis* (Boisd.). The 4th Conf. Pest Control, NRC, Cairo.
- Antonious, A.G. and G. Hegazy (1987):** Feeding deterrent activities of certain botanical extracts against the cotton leafworm, *Spodoptera littoralis* (Boisd.). Annals of Agric. Sci., Fac of Agric., Ain Shams Univ., Egypt, 32 (1): 719 - 729.
- Antonious, A.G. and M.A. Rizk (1994):** Response of *Spodoptera littoralis* (Boisd.) to neem

Azadirachta indica (A.juss) seeds oil. Egypt. J. Appl. Sci. 9 (10): 424 - 434.

- Antonious, A.G.; Hegazy, G.; El-Sharawy, M.F. and L.A.Youssef (1992):** Reaction of feeding the cotton leafworm, *Spodoptera littoralis* (Boisd.) on certain leaves. 3:Latent effect of larval feeding. Med. Fac. Landbouww. Univ. Gent, 57 (3a): 707 - 713.
- Attiah, H.H. (1977):** Ecological assessment of pesticide management on terrestrial ecosystem in Egypt. Proceeding of the VC / AID Univ. of Alexandria, A.R.E. Seminar /workshop in pesticide mangement. March 5 -10.
- Atwood, D.W.; Young, S.Y III and T.J. Kring (1997):** Impact of *B.t.* and thiodicarb alone and in combination on tobacco budworm, mortality and emergence of the parasitoid *Microplitis croceipes*. Proceedings Beltwide Cotton Conf., New Orleans, LA,U.S.A., Jan. 6 -10,2: 1305-1310.
- Auda, M. and D. Degheele (1985):** Joint action of pyrethroids with chitin synthesis inhibitors, organophosphorous and carbamate insecticides on a susceptible and resistant strain of *Spodoptera littoralis* (Boisd.). Med. Fac. Landbouww. Rijksuniv. Gent, 50 (26): 751 – 760..

- Avidov, Z. and I. Harpaz (1969):** Plant pests in Israel. Israel Univ. Press, Jerusalem, 549 p.
- Awadallah, K. T. (1968):** Studies on the aphidophagous syrphids commonly found in the U.A.R. (Diptera : Syrphidae). Ph.D. Thesis, Fac. of Agric., Cairo University.
- Ayyangar, G.S.G. and P.J. Rao (1989):** Neem (*Azadirachta indica*) (*A. juss*) extracts as larval repellents and ovipositional deterrents to *Spodoptera litura* (Fabr.) Indian J.Entomol., 51 (2): 121-124.
- Barakat, A.A.; Fahmy, H.S.M.; Kandil, M.A. and N.M.M. Ebrahim (1985):** Toxicity of the extracts of black pepper, cumin, fennel, chamomile and lupine against *Drosophila melanogaster*, *Ceratitis capitata* and *Spodoptera littoralis*. Indian J. Agric. Sci. 55 (2): 116-120.
- Berliner, E. (1911):** Ueber die Schlafsucht der Mehlmottenraupe. Z. Ges. Getreidew. 3: 63 – 70.
- Bishara, I. (1934):** The cotton worm *Prodenia littura* F. in Egypt. Bull. Soc. Roy. Ent. Egypt. 17 : 288 – 420.
- Blaney, W.M.; Simmonds, M.S.J.; Ley, S.V. and P.S.Jones (1988):** Insect antifeedants: a behavioural and electrophysiological investigation of natural and synthetically derived clerodane diterpenoids.

Entomologia- Experimentalis et Applicata. 46 (3):
267 - 274.

- Brewer, F.D. and E.G. King (1978):** Effects of parasitism by a tachinid *Lixophago diatrachae*, on growth and food consumption of a sugarcane borer larvae. Ann. Entomol., Soc. Amer. 71 (1): 19- 22.
- Brown, E.S. and C.F. Dewhurst (1975):** The genus *Spodoptera* (Lepidoptera: Noctuidae) in Africa and the Near East. Bull. Entomol. Res. 65: 221-262.
- Burges, H.D. and N.W. Hussey (1971):** Microbial control of insects and mites. Academic Press, London and New York 861 pp.
- Busoli, A.C. (1993):** Control of *Pectinophora gossypiella* (Saund.) and *Heliothis spp.* in cotton crop, Using pheromones and by the mating disruption method. Anais do Sociedade Entomologica do Brasil 22 (1): 139 - 148.
- Cai, S.H.; Xiong, Y.Q.; Ke, D.K. and B.J. He (1985):** Studies on the dynamics of pink bollworm population and the damage on cotton. Insect - Knowledge Kunhong Zhishi. 22 (2): 64 - 69.
- Cate, J.R.; Ridgwy, R.L. and P.D. Lingren (1972)** Effect of systemic insecticides applied to cotton on adults of an

- Ichneumonid parasite, *Campoletis perdinctus*. J. Econ. Entomol. 65 (2): 484 – 487.
- Chamberlain, D.J.; Crithley, B.R.; Campion, D.G.; Rafique, M. and M.I. Arif (1992):** Use of a multi – component pheromone formulation for control of cotton bollworms (Lepidoptera : Gelechiidae and Noctuidae) in Pakistan. Bull. Entomol. Res. 82 (4): 449 – 458.
- Chen, R.H. (1987):** Control of the first generation of *Pectinophora gossypiella*. Plant Protection. 13 (4): 2 – 4.
- Chu, C.C. and T.J. Henneberry (1990):** Pink bollworm seasonal distribution, yearly variation and male moth trap catch relationships to population increases in cotton. Southwestern Entomologist. 15 (3): 273 – 280.
- Corlett, G.R. (1961):** Investigation of the effect of *Bacillus thuringiensis* on mammals. Can. Wild life. Serv. Res. Proj. : 1 – 5 Univ. of Ottawa.
- Critchly, B.R. (1991):** Commercial use of pink bollworm pheromone formulations in Egyptian cotton pest management. Pesticide – Outlook. 2 (4): 9 – 13.
- Damo, M.M.C. and A.D. Solsoloy (1997):** Efficacy of *Bacillus thuringiensis* – chemical insecticide mixtures for bollworm, *Helicoverpa armigera* control. J. crop Sc. (Philippines), Philippine. 22 (1) p. 74.

- Darwish, Y.A. and A.I. Farghal (1990):** Evaluation of certain pesticides activity against the cotton white fly, *Bemisia tabaci* and associated natural enemies on cotton plants under field conditions in Assiut. Assiut, J. Agric. Sci. 21 (5): 331 – 339.
- Dejiu, G.: Wright D.I. and J. K. Waage (1992):** Influence of sublethal doses of insecticides on the population systems of *Plutella xylostella* and its parasitoids *Diadegma eucerocephala*. Proceedings 19 International Congress of Entomology. Beijing. China. June 28 – July 4, 276.
- Dhawan, A.K. and G.S. Simwat (1992):** Field evaluation of Neemrich 20 E.C. for management of insect pests on cotton during reproductive phase. In. Proc. 1st Nat. Symp., Allelopathy in Agroecosystems (Agriculture & Forestry), Feb. 12 – 14 held at CCS Haryana Agric. Univ., Hisar, India: 152 – 153. (c.f. Rev. Agric. Entomol., 80 (111): 1102, 1992).
- Dimetry, N.Z. and E.F. Abdalla (1988):** Feeding deterrent effects of various extracts of the bead tree, *Abrus precatorious*, seeds towards *Spodoptera littoralis* (Boisd.). Bull. ent. Soc. Egypt, Econ. Ser., 17 : 85 – 92.

- Dongre, T.K. and G.W. Rahalkar (1992):** Feeding and oviposition deterrents for spotted bollworm of cotton, *Earias vittella* (Fab.) in caulex of an alternate host plant, *Hibiscus sabdsriffa* L. J. Insect Sci. 5 (2): 130 – 133.
- Dongre, T.K. and G.W. Rahalkar (1993):** Feeding deterrent and toxic activity in the leaves of *Sida acuta* L. (Mal) against the larvae of *Earias vittella*. J. Insect Sci. 6 (1): 100 – 101.
- Dunbar, D.M.; Haya, H.K.; Doane, C.C.; Anderson, J.F. and R.H. Weseloh (1973):** Aerial application of *Bacillus thuringiensis* against larvae of elm spanworm and gypsy moth and effects on parasitoids of the gypsy moth. Conn. Agric. Exp. Stn. New Haven, Bull. 735.
- El-Hamaky, M.A.; Watson, W.M. and S.M. El-Fateh (1987):** Potency of certain insecticides and insect growth regulators and their mixtures, on *Spodoptera littoralis* (Boisd.). J. Agric. Res. Tanta Univ. 13 (4): 1198 – 1212.
- El-Henidy, A.H.; Abbas, M.S.T. and M.S.I. El-Dakroury (1978):** Seasonal abundance of certain predators in untreated Egyptian clover and cotton fields in Fayoum Governorate, Egypte. Bull. Soc. ent. Egypte, 62: 82 – 95.

- El-Henidy, A.H.; Abbas, M.S.T. and A.A. Khidr (1987):**
Comparative population densities of certain predators
in cotton fields treated with sex pheromones and
insecticides in Menofia Governorate, Egypt. Bull. ent.
Soc. Egypt, Econ. Ser., 16: 181 – 190.
- El-Maghraby, M.M.A.; Hegab, A. and S.I. Yousif Khalil
(1988):** Interactions between *Bacillus thuringiensis*
Berl., *Beauveria bassiana* (Bals.) and the host /
parasitoid system *Spodoptera littoralis* (Boisd.)/
Microplitis rufiventris Kok. J. Appl. Entomol. 106
(4): 417 – 421.
- El-Mandarawy, M.B.R. (1995):** Studies on the effects of
Bacillus thuringiensis Berliner and a chemical
insecticide on *Spodoptera littoralis* (Boisd.) and its
parasitoid *Microplitis rufiventris* Kok. Ph.D. Thesis,
Fac. of Sci., Cairo University.
- El-Minshawy, A.M. and E.M. Hegazi (1980):** Effect of
temperature and host age on the progeny production
and sex ratio of *Microplitis rufiventris* Kok.
(Hymenoptera : Braconidae). Z. age. Ent. 90: 52 – 57.
- El-Mosa, H. (1986):** Prospects of using sex pheromone for the
control of spiny bollworm in cotton growing in Syria.
Dirasat, 13 (5): 165 – 174.

- El-Sayed, E.I. and E.I. Sayed (1982):** Neem (*Azadirachta indica*, *A. juss*) seeds as antifeedants and ovipositional repellents for the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.). Bull. ent. Soc. Egypt, Econ. Ser., 13: 49 – 58.
- El-Sebae, A.H. (1981):** Mode of toxicity and hazards of insecticides used in Egypt. 1st Egyptian – Hungarian Conf. on Pl. Prot., Pl. Path., Entomol. And Pesticide Chem., Budapest, May 27 – 31.
- El-Shaarawy, M.F.; Antonious, A.G.; Hegazy, G. and L.A. Youssef (1992):** Reaction of feeding the cotton leafworm, *Spodoptera littoralis* (Boisd.) on certain plant leaves. 1: Antifeeding effect. Med. Fac. Landbouww. Univ. Gent. 57 (3a): 689 – 695.
- El-Sheikh, M.A.K.; Ibrahim, S. M. and S.A.S. El-Maasarawy (1993):** Food consumption and utilization in larvae of *Mythimma* (= *leucania*) *loreyi* (Dup.) parasitized by *Meteorus gyrator* Thun. Bull. Ent. Soc. Egypt, 71: 173 – 184.
- El-Zemaity, M.S. and S.A. El-Refai (1987):** Joint action of fenvalerate mixtures against the cotton leafworm, *Spodoptera littoralis* (Boisd.). Annals of Agric. Sci., Fac. of Agric., Ain Shams 32 (3): 1741 – 1749.

REFERENCES

- El-Zohairy, M.M.; Metwally, E.M.; Nasr, E.A.; El-Sebay, Y.; Radwan, S. M.E. and W.M.H. Desuky (1994):** Ecological studies on the immature stages of the cotton leafworm *Spodoptera littoralis* (Boisd.) in Sharkia Governorate. J. Agric. Res., Egypt, 72 (4): 1033 – 1050.
- Fayad, Y.H. and A. A. Ibrahim (1980):** Effect of some new insecticides of cotton leafworm on the number of predators in cotton fields. The 1st Conf. Pl. Prot. Inst. 2: 237 – 348.
- Fisher, R. and L. Rosner (1959):** Toxicology of the microbial insecticide, Thuricide. J. Agric. Food Chem. 7: 686 – 688.
- Gao, Z.R.; Zhao, H.Y. and Y.F. Jiang (1992):** Studies on economic threshold for pink bollworm on cotton in Henan Province. Acta Phytophylacica Sinica, 19 (2): 153 – 164.
- Gerling, D. (1969):** The parasites of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) eggs and larvae in Israel. Israel J. Ent. 4: 73 – 81.
- Gerling, D. (1971):** Occurrence, abundance and efficiency of some local parasitoids attacking *Spodoptera littoralis* (Lepidoptera: Noctuidae) in Selected cotton in Israel. Ann. Entomol., Soc. Amer. 64 (2): 492 – 499.

- Grosch, D.S. (1975):** Reproductive performance of *Bracon hebetor* after sublethal doses of carbaryl. J. Econ. Entomol. 68 (2): 659 – 662.
- Guirguis, M.W. and W.M. Watson (1981):** Laboratory and field studies on the efficiency of some insecticides against pink bollworm *Pectinophora gossypiella* (Saund.) Proc. 4th Arab Pest. Conf., Tanta Univ. 34: 105 – 111.
- Guirguis, M.W.; Watson, W.M. and F.A. Khalil (1976):** Efficiency of various insecticides in controlling the three cotton bollworms, *Pectinophora gossypiella* (Saund.), *Earias insulana* (Boisd.) and *Heliothis zea* (Boddie.). Bull. ent. Soc. Egypt, Econ. Ser., 10: 145.
- Guirguis, M.W.; Watson, W. M.; Gohar, K. and R. Salem (1991a):** Toxicity and latent effect of two wild plant extracts on the cotton leafworm, *Spodoptera littoralis*. Egypt, J. Agric. Res. 69 (1): 11 – 21.
- Guirguis, M.W.; Watson, W.M.; El-Feshawi, A.A. and M.A. Nassef (1991b):** Occurrence and seasonal abundance of the pink bollworm *Pectinophora gossypiella* Saund. infesting cotton. Egypt, J. Agric. Res. 69 (1): 63 – 71.
- Habib, A.; Rizk, G.N.; Farghaly, H.T.A. and Z.A. Ragab (1976):** Seasonal abundance of some predators in

- cotton and its relation to certain pests. Bull. Soc. ent. Egypte, 60: 191 – 196.
- Hafez, M. (1951):** Notes on the introduction and biology of *Microplitis demolitor* Wilk. (Hymenoptera : Braconidae). Bull. Soc. ent. Egypte, 35: 107 – 121.
- Hafez, M. (1953):** Studies on *Tachina larvarum* L. (Diptera : Tachinidae): 3: Biology and life history. Bull. Soc. ent. Egypte, 37: 305 – 335.
- Hafez, M. (1960):** The effect of some new insecticides on predators of the cotton leafworm in cotton fields. Agric. Res. Rev., Cairo, 38: 47 – 79.
- Hafez, M. and S.M. Hassan (1969):** On correct identity of Egyptian cotton leafworm (Lepidoptera : Noctuidae). Bull. Soc. ent. Egypte, 53: 215 – 219.
- Hafez, M.; Ali, A.M.; Mikhail, F. and F. Abd El-Kawi (1975):** Fluctuation of population of certain predators in cotton and clover fields in Assiut Governorate (Egypt). Agric. Res. Rev., Egypt, 53 (1): 47 – 51.
- Hafez, M.; Tawfik, M.F.S.; Azab, A.K. and A.A. Ibrahim (1976):** Survey and economic importance of parasites of the cotton leafworm, *Spodoptera littoralis* (Boisd.) in Egypt. Bull. Soc. ent. Egypte, 60: 179 – 189.
- Hafez, M.; Fayad, Y.H. and A.A. Sarhan (1977):** Preliminary indication of the effect of Nuvacron U.L.V on

abundance of predators in cotton fields in Egypt.
Agric. Res. Rev. Egypt, 55: 111 – 116.

Hamilton, J.T. and F.I. Attia (1977): Effects of mixtures of *Bacillus thuringiensis* and pesticides on *Plutella xylostella* and the parasite *Thyraeella collaris*. J. Econ. Entomol. 70: 146 – 148.

Hammad, S.M.; El-Minshawy, A.M. and A. Salama (1965): Studies on *Microplitis rufiventris* Kok. (Hymenoptera: Braconidae). Bull. Soc. ent. Egypte, 49: 215 – 219.

Hassan, K.A. (1998): Studied on some arthropodous natural enemies of the pink bollworm, *Pectinophora gossypiella* (Saund.) M.Sc. Thesis, Fac. of Agric., Moshtohor, Zagazig University.

Hassan, M.S. (1957): Studies on the damage and control of *Aphis maidis* Fitch in Egypt. Bull. Soc. ent. Egypte, 29: 213 – 230.

Hassan, A.G.; Hosny, M.M. and A.A. Soliman (1960): Analysis of the relative abundance of some beneficial insects on cotton plants treated by certain insecticides. Bull. Soc. ent. Egypte, 44: 179 – 184.

Hassanein, M.H. (1956): Studied on the activity, phenology and population density of the aphid lion, *Chrysopa vulgaris* Schneider. Ann. Agric. Sci. 12: 145 – 160.

- Hegazi, E.M. (1972):** Studies on certain natural enemies attacking the immature stage of certain lepidopterous pests [parasites of egg- larval and larval stage of the cotton leafworm, *Spodoptera littoralis* (Boisd.)] M.Sc. Thesis, Fac. of Agric., Alexandria University.
- Hegazi, E.M. (1976):** Further studies on certain natural enemies attacking the cotton leafworm, in Alexandria region. Ph.D. Thesis, Fac. of Agric., Alexandria University.
- Hegazi, E.M; Hammad, S.M. and A.M. El- Minshawy (1977a):** Suitability of *Spodoptera littoralis* (Boisd.) larvae for development of *Microplitis rufiventris* Kok. J. Agric. Sci., Camb. 89: – 662.
- Hegazi, E.M; Hammad, S.M. and A.M. El- Minshawy (1977b):** Field and laboratory observations on the parasitoids of the cotton leafworm in Alexandria. Z. ang. Ent. 84: 310 – 321.
- Hegazi, E. A.; Kolaib, M. O. and M. I. Abd El-Fattah (1980):** Development of *Microplitis rufiventris* kok at different temperatures. J. Agric. Sci., Camb. 94: 739-740.
- Hegazi, E. .M; Hammad, S.M.; Altahtawy, M. and S.K. El-Sawaf (1973):** parasof the larval stage of the cotton leafworm, *Spodoptera littoralis* (Boisd.) (Noctuidae,

- Lepidoptera) in Alexandria region. Z. ang. Ent. 74: 332 – 336.
- Hegazi, E.M.; Hammad, S.M.; Altahtawy, M. and S.K. El-Sawaf (1974):** Notes on biology of *Chelonus inanitus* (Hymenoptera: Braconidae). Z. ang. Ent. 75: 291 – 294.
- Hegazy, G.; Antonious, A.G.; El-Shaarawy, M.F. and L.A. Yousef (1992):** Reaction of feeding the cotton leafworm, *Spodoptera littoralis* (Biosd.) on certain plant leaves. 2: Effect of plant extracts. International symposium on crop protection, Med. Fac. landbouww Univ. Gent. 57 (3A) 697 – 705.
- Hongo, H. and A.K. Karel (1986):** Effect of plant extracts on insect pests of common beans. J. Appl. Ent. 102 (2): 164 – 169.
- Hosny, M.M.; Topper, C.P.; Moawad, G.M. and G.B. El-Saadany (1986):** Economic damage thresholds of *Spodoptera littoralis* (Biosd.) (Lepidoptera: Noctuidae) on cotton in Egypt. Crop Protection, 5 (2): 100 – 104.
- Hussein, Y.F. and Amira A. Ibrahim (1986):** Impact of successive insecticidal applications at different interval periods on the number of predators in cotton fields. Bull. ent. Soc. Egypt, Econ. Ser., 15: 47 – 58.

- Ibrahim, A.A. (1974):** Studies on biological control of *Spodoptera littoralis* (Boisd.) in A.R.E: Ph.D. Thesis Fac. Agric., University of Cairo.
- Ibrahim, A.A. and Y.H. Fayad (1980 a):** The effect of some new insecticides of bollworms on the number of predators in cotton fields. 1st Conf. Pl. Prot. Inst. 2: 349 – 375.
- Ibrahim, A.A. and Y.H. Fayad (1980 b):** Effect of granule insecticides on number of predators in cotton fields in Egypt. J. Agric. Res. Tanta Univ. 6 (1): 242 – 248.
- Ibrahim, A.A.; Fayad, Y.H.; Shalaby, F.F. and E.A. Kares (1986):** Morpho- logical studies on the immature stages of the parasite *Zele nigricornis* (Walk.) (Hymenoptera: Braconidae). Annals of Agric. Sci, Moshto- hor, 24 (2): 1147 – 1156.
- Idris, A.B. and E. Grafius (1993):** Pesticides affect immature stages on *Diadegma insulare* (Hymenoptera: Ichneumonidae) and its host the diamond back moth Lep. Plutellidae. J. Econ. Entomol. 86 (4): 1203 – 1212.
- Ismail, I.I. (1974):** Seasonal abundance of the rove- beetle *Paederus alfieri* Kock. [Coleoptera: Staphylinidae]. Bull. Soc. ent. Egypte, 58: 213 – 216.

- Kamal, M. (1951 a):** Biological control projects in Egypt, with a list of introduced parasites and predators. Bull. Soc. ent. Egypte, 35: 205 – 220.
- Kamal, M. (1951b):** The biological control of the cotton leafworm, (*Prodenia litura* F.) in Egypt. Bull. Soc. ent. Egypte, 35: 221 – 270.
- Kares, E.A. (1978):** Studies on certain economic pests. M.Sc. Thesis, Fac. of Agric., Alexandria University.
- Kares, E.A. (1985):** Studies on parasites of *Spodoptera littoralis* (Boisd.) in Egypt. Ph.D. Thesis, Fac. of Agric., Zagazig University.
- Kares, E.A.(1990 a):** Impact of diflubenzuron in *Spodoptera littoralis* (Boisd.) larvae parasitized by *Zelex nigricornis* (Walk.) Zagazig, J. Agric. Res. 17 (3B): 953 – 958.
- Kares, E.A.(1990 b):** Some biological aspects of *Zelex chlorophthalma* (Ness) (Hymenoptera: Braconidae), parasitoid on *Spodoptera littoralis* (Boisd.). Egypt, J. Appl. Sci. 5 (6): 43 – 50.
- Kares, E.A.(1991 a):** Effect of mixtures of *Bacillus thuringiensis* (Berliner) and chemical insecticides against larvae of the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). Egypt, J. Biol. P. Cont. 1 (2): 15 –23.

- Kares, E.A. (1991 b):** Toxicity of bioinsecticide, dipel on the potato tuber moth larvae *Phythorimaea operculella* (Zeller) and these parasitized by *Apanteles litae* Nixon var. *operculellae*. Annals of Agric. Sci., Moshtohor, 29 (4): 1781 – 1790.
- Kares, E.A.; Abdel-Rahman, A.M.; El-Moursy, A.A. and M.B.R. El – Mandarawy (1992):** Qualitative and quantitative studies on the haemolymph of *Artogeia rapae* L. larvae after a bioinsecticidal and gamma radiation treatment. Egypt, J. Biol. P. Cont. 2 (2): 131 – 141.
- Kares, E.A.; Fayad, Y.H. and A.A. Ibrahim (1988):** Effect of certain groups of insecticides on the abundance of predators in cotton fields. Annals of Agric. Sci. , Moshtohor, 26 (4): 2717 – 2733.
- Kares, E.A.; Hafez, A.A.; El-Khayat, E.F. and F.F. Shalaby (1993):** Sunflower and maize in cotton plantations to increase the natural role of entomophagous insects. Annals of Agric. Sci., Moshtohor, 31 (2): 1199 – 1211.
- Kares, E.A.; El-Moursy, A.A.; Zohdy, N.; Abd El-Rahman, A.M. and M.B.R. El-Mandarawy (1998):** Biological and toxicological studies on the parasitoid

- Microplitis rufiventris* Kok. Egypt, J. Agric. Res. 76 (4): 1499 – 1513.
- Kashyap, R.K. and A.N. Verma (1987):** Management of spotted bollworms (*Earias sp.*) in cotton. International J. Tropical Agric. 5 (1): 1-27.
- Kasopers, H. (1965):** Eroterungen Zur Prufung Von Fungiziden in Obsthau. Pflanzenschutz – Nachrichten. "Bayer" 18: 83 –92.
- Kaya, H.K. and D.M. Dunbar (1972):** Effect of *Bacillus thuringiensis* and carbaryl on elm spanworm egg parasite, *Telenomus alsophilae*. J. Econ. Entomol. 65 (4): 1132 – 1134.
- Kassem, S.H. and M.I. Zeid (1987):** Comparison of the insecticidal efficiency of certain insecticides and their mixtures with insect chitin inhibitors. Alex., J. Agric. Res. 32 (3): 335 – 345.
- Khalil, F.A. and W.M. Watson (1986):** Residual effectiveness of binary mixtures of Dimilin (IGR) with certain insecticides against *Spodoptera littoralis* (Boisd.) larvae. Agric. Res. Rev., Cairo, 61 (1): 99 – 108.
- Kokujev, N. (1914):** Hymenoptera parasitica nova fauna turanica-a. V.I. Plantkov collecta. Revusse d' Entomologie, St. Ptersburg 13: 513 – 514.

- Kolaib, M.A.O. (1978):** Studies on certain, natural enemies attacking the cotton leafworm *Spodoptera littoralis* (Boisd.) in Menoufia Governorate. M.Sc. Thesis, Fac. of Agric., Menoufia University.
- Kongkathip, B.; Kongkathip, N. and S. Wongthong (1993):** New insecticidal substance for cotton bollworm, (*Heliothis armigera*) from *Tinospora crispa*. Kasetsart Univ. Bangkok (Thailand). Fac. Sci. Ruambotkhayto ngan wichai 2536, 162 p.p. 56.
- Korat, D.M. and S. Lingappa (1996):** Monitoring of pink bollworm with sex pheromone traps and its relationship with larval population and field incidence. Karnataka, J. Agric. Sci. 9 (3): 432 – 437.
- Krug, E. and P. Proksch (1993):** Influence of alkaloids on survival and growth of *Spodoptera littoralis*. Biochem. Syntematics and Ecology 21 (8): 749.
- Lewis, W.J. (1970):** Life history and antomy of *Microplitis corceipes* (Hymenoptera: Braconidae), a parasite of *Heliothis spp.* (Lepidoptera: Noctuidae). Ann. Entomol, Soc. Amer., 63 (1): 67 – 71.
- Litchfield, J.T. and F. Willcoxon (1949):** A simplified method of evaluating dose – effect experiments. J. Pharmacol. And Exp. Therap. 96: 99 – 133.

REFERNCES

- Luo, S.B.; Yan, J.P.; Chai, C.J.; Liang, S.P.; Zhang, Y. and G.K.le (1986):** Control of pink bollworm, *Pectinophora gossypiella* with *Bacillus thuringiensis* in cotton fields. Chinese J. Biol. Con. 2 (4): 167 – 169.
- Madrid, F.J. and R.K. Stewart (1981):** Impact of Diflubenzuron spray on gypsy moth parasitoids in the field. J. Econ. Entomol. 74 (1): 1 –2.
- Makkar, A.W. and S.N. Kostandy (1995):** Relationship between spiny bollworm males attracted to pheromone traps and larval infestation in cotton bolls. Annals of Agric. Sci., Moshtohor, 33 (4): 1529 – 1538.
- Manaa, S.H. and A.A. Attia (1992):** Insecticidal properties of some lignin compounds, extracted from the root bark of *Artemisia argentea* L'her, on the cotton leafworm, *Spodoptera littoralis* (Boisd.) Assiut, J. Agric. Sci. 23 (2): 119 – 139.
- Mansour, E.S. (1997):** Using methods to substitute insecticides for controlling Egyptian cotton leafworm in Qualubia Governorate. M.Sc. Thesis, Fac. of Agric., Moshtohor, Zagazig University.

- Matter, M.M. (1993):** Effect of *Bacillus thuringiensis* diseased larvae of *Pieris rapae* on the parasitoid *Hyposter ebeninus*. Egypt, J. Biol. P. Cont. 3 (2): 213 – 227.
- Mattes, O. (1927):** Pakrankheiten der Meh- larven und versuche ueber ihre verwendbarkeit als biologisches bekaemp fungsmittel. Sitzungsber. Ges. Befoerder. Ges. Naturw. Marburg 62: 381 – 417.
- Mc Donald, R.C.; Kok, L.T. and A.A. Yousten (1990):** Response of fourth instar *Pieris rapae* parasitized by the braconid *Cotesia rubecula* to *Bacillus thuringiensis* subsp. *kuristaki* delta- endotoxin. J. Inventebr. Pathol. 56: 422 – 423.
- Mc Pherson, R.M.; Hensley, S.D.; Graves, J.B. and E.G. King (1976):** Dosage-mortality responses of the parasite, *Lixophaga diatracea* to selected insecticides in laboratory. J. Econ. Entomol. 69 (1): 105 – 108.
- Meisner, M.W.; Palevth, D. and N. Aharonson (1981):** Phagodeterreny induced by leaves and leaf extracts of *Catharanthus roseus* in the larvae of *Spodoptera littoralis*. J. Econ. Entomol. 74: 131 – 135.
- Metcalf, R.L.; Reynolds, H.T. and T.R. Fukuto (1967):** Carbamate insecticideswhere are they headed? Farm. Chem. 130: 45 – 48.

- Moawad, G.M.; Shalaby, F.F.; Metwally, A.G. and M.M. El-Gemeiy (1982-1983):** Laboratory pathogenecity tests with two commercial preparations of *Bacillus thuringiensis* (Berliner) on the first instar larvae of spiny bollworm. Bull. Soc. ent. Egypte, 64: 137 – 141.
- Moawad, G.M.; Amin, A.A.H. and A.M. Hossain (1994):** Spatial distribution patterns of two cotton bollworms, *Pectinophora gossypiella* and *Earias insulana* in Fayoum, Egypt. Annals of Agric. Sci., Cairo, 39 (2): 805 – 813.
- Moawad, G.M.; Sawires, Z.R.; El-Hamaky, M.A. and M.F.Gergis (1991):** The impact of sex pheromones and insecticides on the natural enemies in cotton fields in middle Egypt. Bull. ent. Soc. Egypt, Econ. Ser., 19: 237 – 242.
- Moawad, G.M.; Raslan, S.A.A.; Metwally, E.M.; Ibrahim, A.E. and W.M.H. Desuky (1996):** New approach for pink bollworm, *Pectinophora gossypiella* (Saunders) control in Egypt using fatal attraction. J. Agric. Res. 74 (4): 957 – 971.
- Mochida, O. and T. Okoda (1974):** A bibliography of *Spodoptera spp.* (Lepidoptera: Noctuidae). Misc. Bull. Kyushunot. Agric. Expt. Sta. 49: 1 – 110.

- Mohammed, A.A.; Ibrahim, I.L.; El-Khouly, A.S. and F.H. Tamer (1988):** Prey preference of the aphid predator *Cydonia vicina* Muls. (Coleoptera: Coccinellidae). Bull. Soc. ent. Egypte, 68: 149 – 158.
- Morallo-Rejesus, B.; Obra, J.B. and F.V. Salazar (1992):** Sensitivity of *Cotesia plutella* and *Diadegma semiclausum*, parasitoids of diamondback moth *Plutella xylostella* L., to several insecticides. Proceedings 19 International Congress of Entomology. Beijing. China. June 28- July 4, 308.
- Mourad, M.A.; Omar, M.E. and A.A. Mahran (1991):** Field potency of different insecticides against cotton bollworms. Egypt, J. Agric. Res. 69 (1): 57 – 62.
- Moussa, M.A.; Zaher, M.A. and F. Kotby (1960):** Abundance of the cotton leafworm, *Prodenia litura* F., in relative to host plant. 1: Host plants and their effect on biology. Bull. Soc. ent. Egypte, 44: 241 – 251.
- Nada, M.A. (1990):** Control of certain pests infesting cotton plants in Sharkia Governorate. Ph.D. Thesis, Fac. of Agric., Zagazig University.
- Naguib, M.A. (1980):** Effect of the predators population in clover fields and their population in adjacent cotton fields. Proc. 1st Conf. Pl. Prot. Res. Inst., pp. 309 – 321.

- Naik, M.I. Lingappa, S. and C.P. Mallapur (1996):**
Monitoring pink bollworm, *Pectinophora gossypiella*
(Saund.) using pheromone trap. Mysore, J. Agric. Sc.
30 (1): 43 – 47.
- Nandihalli, B.S.; Patil, B.V. and P. Hugar (1993):**
Optimisation of pheromone traps for cotton pink
bollworm, *Pectinophora gossypiella* Saund.
Karnataka J. Agric. Sc. 6 (3): 300 – 301.
- Nasr, F.N. (1979):** Studies on certain natural enemies of some
pests. M.Sc. Thesis, Fac. of Agric., Alexandria
University.
- Nassef, M.A.; El-Feshawi, A.A.; Abo-Sholooa, M.K.A. and
W.M. Watson (1996):** Seasonal abundance of certain
predators as influenced by cotton planting dates and
weather factors. Egypt, J. Agric. Res. 74 (3): 585.
- Nealis, V. and K. Van Frankenhuyzen (1990):** Interactions
between *Bacillus thuringiensis* Berliner and *Apanteles*
fumiferanae Vier. (Hym.: Braconidae) a parasitoid of
the sprue budworm, *Christoneura fumiferana* (Clem)
(Lep.: Tortricidae). Canadian Entomologist, 122 (7-
8): 585 – 594.
- Newsom, L.D. (1974):** Predator insecticide relationships.
Entomophaga 7: 13 – 23.

- Qureshi, Z.A.; and N. Ahmed (1989):** Efficacy of combined sex- pheromones for the control of three major bollworms of cotton. J. Appl. Entomol. 108 (4): 386 – 389.
- Qureshi, Z.A.; Ahmed, N. and T. Hussain (1993):** Pheromone trap catches as a means of predicting damage by pink bollworm larvae in cotton. Crop Protection, 12 (8): 597 – 600.
- Radwan, H.S.A.; Assal, O.M.; Samy, M.A. and M.R.R. Riskallah (1986):** Ovicidal action: Potentiation of synthetic pyrethroids by insect growth regulators against the cotton leafworm, *Spodoptera littoralis* (Boisd.). Bull. ent. Soc. Egypt, Econ. Ser., 14: 275 – 283.
- Rahman, M. (1970):** Effect of parasitism on food consumption of *Pieris rapae* larvae. J. Econ. Entomol. 63 (3): 820 – 822.
- Refaci, A.F.; Hegazy, M.A.; Hussein, N.M. and M.A. El-Hamaky (1990):** Efficiency of certain insecticides, insect growth inhibitors and their combinations against the cotton leafworm larvae in cotton plantations in Egypt. Med. Fac. Landbouww. Univ. Gent. 55 (2b): 577 – 582.

- Ridgway, R.L. and D.A. Lindquist (1966):** Systemic activity of shell SD - 9129. In cotton plants. J. Econ. Entomol. 59: 961 - 946.
- Ridgway, R.L.; Bariola, L.A.; Jones, S.L. and W.L. Lowrey (1986):** Stem treatments with systemic insecticides against *Heliothis zea* (Boddie) and *H. virescens* (F.). Bull. Entomol. Res. 57: 553 - 558.
- Rivany, E. (1961):** The phenology of *Prodenia litura* F. in Israel with reference to its occurrence in the Near East at large. Bull. Res. Coun. Israel. B. 10 (1-3): 100 - 106.
- Saad, A.S.A.; Marei, A.S.M.; Abu El- Amayem, M.M.; Madkour, A. and A. El-Bahrawi (1974):** Toxic effects of certain insecticides on cotton fiber quality. 2nd Egpt. P. Cont. Cong., Alex. (1): 255 - 267.
- Salama, H.S. and A. Sharaby (1988):** Feeding deterrence induced by some plants in *Spodoptera littoralis* and their potentiating effect on *Bacillus thuringiensis* Berliner. Insect Sci. and its application. 9 (5): 573 - 577.
- Salama, H.S. and F.N. Zaki (1984):** Impact of *Bacillus thuringiensis* Berl. on the predator complex of

- Spodoptera littoralis* (Boisd.) in cotton fields. Z. ang. Ent. 97 (5): 485 – 490.
- Saleh, M.M.E. (1986):** Ecological studies on certain corn borers in Egypt. M.Sc. Thesis, Fac. of Agric., Cairo. University.
- Salem, S. and H.S. Salama (1985):** Sex-pheromone for mass trapping of *Spodoptera littoralis* (Boisd.) in Egypt. Z.ang. Ent. 100 (3): 316 – 319.
- Salem, S.; Radwan, S.M.E. and M.A. Hamaky (1990):** Prospects of using sex – pheromone for the control of cotton bollworms, *Earias insulana* and *Pectinophora gossypiella* (Saund.) in cotton fields. Annals of Agric. Sci, Moshtohor, 28 (3): 1743 – 1752.
- Shalaby, F.F. (1968):** Morphology of immature stages and bionomics studies on *Microplitis rufiventris* Kok. (Braconidae, Hymenoptera) with reference to pathology of parasitism on the host. M.Sc. Thesis, Fac. Agric., Alexandria University.
- Shalaby, F.F.; Kares, E.A. and A.A. Ibrahim (1983a):** Effect of intercropping maize in cotton fields on the attractiveness of predaceous insects. Annals of Agric. Sci., Moshtohor, 20 (3): 109 – 123.

- Shalaby, F.F.; Kares, E.A. and A.A. Ibrahim (1983b):** Effect of a single insecticidal application on the populations of predaceous insects in cotton fields. *Annals of Agric. Sci., Moshtohor*, 20 (3): 125 – 134.
- Shalaby, F.F.; Ibrahim, A.A. and E.A. Kares (1986):** Effects of parasitism by *Microplitis rufiventris* Kok. On the susceptibility of *Spodoptera littoralis* (Boisd.) larvae to Bolstar 720 EC. *Bull. ent. Soc. Egypt, Econ. Ser.*, 15: 165 – 172.
- Shalaby, F.F.; Metwally, A.G.; El-Lakwah, F.A. and H.M. El-Gemeiy (1993a):** Effect of field application of a commercial product of *Bacillus thuringiensis* (Ber.) on *Pectinophora gossypiella* (Saund.) infesting cotton bolls. *Annals of Agric. Sci., Moshtohor*, 20: 135 – 144.
- Shalaby, F.F.; El-Khayat, E.F.; Hafez, A.A. and E.A. Kares (1993b):** Impact of chemical and bioinsecticides and their combinations on entomophagous insects in cotton fields and damage by the cotton leafwoand bollworms. *Egypt, J. Biol. P. Cont.* 3(1): 71 – 84.
- Shu, C.E. and C.Y. Cao. (1987):** Effects of damage of cotton squares and flowers on boll formation caused by

Pectinophora gossypiella. Insect knowledge 24
(1): 19 – 21.

Smith, G.J. and D.S. Grosch (1976): Fluoroacetate induced changes in the fecundity and fertility of *Bracon hebetor* females. J. Econ.Entomol. 69 (4): 521 – 522.

Smith, R.F.; Addikisson, P.L.; Gutierrez, A.P.; Harris, F.A.; Reynolds, H.T. and T.F. Watson (1975): Observation on pest management and pesticide management in the Arab Republic of Egypte. Report of United States VC/ AID pest management colloquium participants 1 –19.

Sneh, B. and S. Gross (1981): Toxicity of Avocado leaves (*Persea americana*) to young larvae of *Spodoptera littoralis* (Boisd.) (Lep.: Noctuidae). Z. ang. Ent. 92: 420 – 422.

Sobeiha, A.M.K.; Zidan, Z.H.; Mahmoud, F.A. and S. Tantawy (1980): Residual influence of Dimilin and Bactospine on the bio- effecacy of certain synthetic pyrethoids against the cotton leafworm, *Spodoptera littoralis* (Boisd.) Bull. ent. Soc. Egypte, Econo. Ser. 12: 165 – 170.

- Sun, Y.P. and E.R. Johnson (1960):** Analysis of joint action of insecticides against house flies. J. Econ. Entomol. 53: 887 – 892.
- Swan, L.A. (1964):** Beneficial insects. Harper and Row, New York, 429 pp.
- Taneja, S.L. and A.P. Jayaswal (1986):** Population dynamics of pink bollworm, *Pectinophora gossypiella* (Saunders) on upland cotton. Insect Science and its application 7 (4): 569 – 573.
- Tawfik, M.F.S. (1977):** Natural enemies of *Heliothis armigera* Hb., with special reference to the larval parasite, *Bracon brevicornis* Wesmael. M.Sc. Thesis, Fac. of Agric., Cairo University.
- Tawfik, M.F.S.; Hafez, M. and A. Ibrahim (1977a):** On the bionomics of *Microplitis rufiventris* Kok. Bull. Soc. ent. Egypte, 61: 123 – 135.
- Tawfik, M.F.S.; Hafez, M. and A. Ibrahim (1977b):** Biology of the immature forms of *Microplitis rufiventris* Kok (Hymenoptera: Braconidae) Bull. Soc. ent. Egypte, 61: 205 – 213.
- Tawfik, M.F.S.; Afifi, A.L; El-Dakroury, M.S.I. and F.M.H. Eid (1993):** The biology of *Chelonus* sp. ,A parasitoid newly reported for the spiny bollworm

- in Egypt (Braconidae: Hymenoptera). Egypt, J. Biol. P. Cont. 3 (1): 29 – 40.
- Teich, I.; Shani, A. and J.T. Klug (1985):** The role of mass trapping of the Egyptian cotton leafworm (*Spodoptera littoralis* Boisd.) in its integrated pest control. J. Envir. Sc. And Heal., A. Envir. Sci. and Eng., 20 (8): 943 – 955. [C.F. R.A.E. 81 (2), Abst. No. 1585].
- Temerak, S.A. (1976):** Studies on certain mortality factors affecting distribution and abundance of sugarcane borers in upper Egypt. Ph.D. Thesis, Fac. of Agric, Assiut University.
- Temerak, S.A. (1982):** Interaction between the bacterial pathogen *Bacillus thuringiensis* and the braconid parasitoid *Bracon brevicornis* on *Sesamia cretica* larvae in response to 5 treatments. 1st National Symposium and Workshops on Utilization of Microorganism in pest control.
- Thompson, W.R. (1946):** A catalogue of the parasites and predators of insect pests. 1: Parasites of Lepidoptera. Imperial Agricultural Burcaux, Institute of Entomology, parasite Service, Belleville, Ont, Canada 1 (7): 259 – 385.

- Thompson, W.R. (1953):** A catalogue of the parasite and predators of insect pests. 2: host parasite Catalogue part 2 host of the Hymenoptera (Agaonidae to Braconidae). Ottawa, ONT., Canada.
- Watanabe, C. (1938):** A braconid parasite of *Porthesia similis* (Fuessly) host record of Braconidae. Insecta mastum, 13 (1): 38.
- Watson, W.M.; Dahan, A.A., Khalil, F.A. and A. Shoeib (1981):** Effectiveness on sequential use of insecticides on pink bollworm, *Pectinophora gossypiella* infestation. 4th Arab Pest. Conf., Tanta Univ. 3A: 95 – 104.
- Watson, W.M.; Rashad, A.M.; Hussein, N.M. and M.W. Guirguis (1986):** Potencies of certain insecticides against the pink bollworm, *Pectinophora gossypiella* (Saund.) as influenced by chemical control program in Egypt. Bull. ent. Soc. Egypt, Econ. Ser., 15: 79 – 86.
- Willcocks, F.C. and S.Bahgat (1937):** The insect and related pests of Egypt. Vol. 1. Part 2. Insects and mites injurious to the cotton plant. Royal Agric. Soc., Entomological Section, Cairo, Egypt.

- Wollam, J.D. and W.G. Yendol (1976):** Evaluation of *Bacillus thuringiensis* and a parasitoid for suppression of the gypsy moth. J. Econ. Entomol. 69: 113 – 118.
- Youssef, E.A. (1997):** Damage and damage loss assessments of certain insect pests attacking cotton in Egypt. Ph.D. Thesis, Fac. of Agric., Ain Shams University.
- Youssef, L.A. (1991):** Physiological reactions of feeding the cotton leafworm on certain naturally occurring deterrents. M.Sc. Thesis, Fac. of Agric., Ain Shams University.
- Youssef, L.A. (1998):** Physiological disorders of the cotton leafworm, *S. littoralis* (Boisd.) fed on certain plants and synthetic media. Ph.D. Thesis, Fac. of Agric., Ain Shams University.
- Zaki; F.N.; Mariy, F.M.A.; Antounious, A.G.; Hegazy, G.M. and N.A. Farag (1987):** Effect of certain insect growth inhibitors on some biological aspects of the ectoparasite *Bracon brevicornis* (Wesm.), (Hymen.: Braconidae). Annals of Agric. Sci.; Ain Shams Univ. 32 (3): 1791 – 1801.
- Zidan, Z.H.; El-Saadany, G.B.; Sharaby, A., Omar, E.A. and S. Abdel – Aziz (1994):** Biological activity of essential oil of *Dodonea viscosa* plant against the

cotton leafworm, *Spodoptera littoralis* (Boisd.). 5th Conf. Agric. Dev. Res., Fac. Agric., Ain Shams Univ., Egypt, 2: 845 – 864.