

<b>CONTENTS</b>	<b>Page</b>
<b>I-INTRODUCTION</b>	<b>1</b>
<b>II-REVIEW OF LITERATURE</b>	<b>3</b>
II-1-Biological studies of some bollworm parasitoids	3
II-2-Biological studies and feeding capacity of some bollworm predators	10
II-3-Host plants of the Pink Bollworm (PBW) <i>Pectinophora gossypiella</i> (Saund.) and Spiny Bollworm (SBW) <i>Earias insulana</i> (Boisd. ) in the world	15
II-4-Survey of parasitoids and predators of PBW and SBW in the world	22
II-4-a-Bollworms' parasitoids	22
II-4-b-Bollworms' predators	27
II-5-Population dynamics, seasonal abundance and damage caused by <i>P. gossypiella</i> and <i>E. insulana</i> in the cotton fields	37
II-6-Population dynamics and seasonal abundance of parasitoids	42
II-7-Population dynamics and seasonal abundance of bollworm predators	48
<b>III-MATERIALS AND METHODS</b>	<b>57</b>
III-1-Laboratory experiments	57
III-1-1-Experiments on parasitoids	57
<b>CONTENTS</b>	<b>I</b>

\_\_\_\_\_

III-1-1-1-Biological aspects of <i>Bracon</i> and <i>Dibrachys</i>	57
III-1-1-2-The effect of kind of Food offered for adults of parasitoids on their egg production, pre-oviposition, post- oviposition and longevity	58
III-1-2-Experiments of predators	59
III-1-2-1-Eggs of PBW and SBW	60
III-1-2-2-Rearing of predators	60
III-1-2-2-1-Rearing of <i>Hippodamia</i> <i>tredecimpunctata</i>	60
III-1-2-2-2-Rearing of <i>Coccinella undecimpunctata</i>	60
III-1-2-3-Feeding capacity technique	61
III-1-2-4-The effect of kind of prey on some biological aspects of <i>H. 13-punctata</i> and <i>C. undecimpunctata</i>	61
III-1-3-Storage of two coccinellid predators, <i>H. 13-</i> <i>punctata</i> and <i>C. undecimpunctata</i> and its effect on mortality, adult longevity; fecundity and feeding capacity after storage:	62
III-1-3-1-The effect of storage on biological aspects of predators	
III-2-Field studies	65
III-2-1-Survey of PBW, SBW, parasitoids and predators	65
III-2-1-1-Survey of PBW and SBW host plants	65
III-2-1-2-Survey of parasitoids	65
III-2-1-3-Survey of predators	66
III-2-1-4- Percentages of parasitism	67
CONTENTS	II

---

III-2-2-Percentages of infestation and population dynamics of PBW and SBW	67
III-2-3-Times of collecting samples were as follows	68
III-2-3-1-Pin squares	68
III-2-3-2-Cotton blooms	68
III-2-3-3-Cotton bolls	68
III-2-4-Population dynamics of arthropod predators	68
III-2-4-1-The direct count method	69
III-2-4-2-Sweeping method	69
III-3-Statistical analysis	69
 IV-RESULTS AND DISCUSSION	 70
IV-1-Biological aspects of <i>Bracon hebetor</i> and <i>Dibrachys</i> sp.	70
IV-1-1- <i>Bracon hebetor</i> Say	70
IV-1-1-1-Duration of immature stages	70
IV-1-1-2-The relationship between kind of food offered for <i>Bracon hebetor</i> adults, and different constant temperatures, their longevity and fecundity	77
IV-1-1-3-Reproduction potential and life table parameters	83
IV-1-2- <i>Dibrachys</i> sp. (Fam: Pteromalidae)	89
IV-1-2-1-Duration of immature stages	89
IV-1-2-2-Duration of <i>Dibrachys</i> sp. on pupal stages of <i>E. insulana</i> and <i>P. gossypiella</i>	99

---

IV-1-2-3-The relationship between kind of food offered to <i>Dibrachys</i> sp. adults, at different constant temperatures, their longevity and fecundity	104
IV-1-2-4-Reproduction potential and life table parameters	109
IV-2-Bionomics and feeding capacity of some predaceous insect species	115
IV-2-1- <i>Hippodamia tredecimpunctata</i>	115
IV-2-1-1-Duration of immature stages	115
IV-2-1-2-Feeding capacity of <i>H. tredecimpunctata</i> larvae on <i>P. gossypiella</i> and <i>E. insulana</i> eggs	118
IV-2-2- <i>Coccinella undecimpunctata</i> L.	129
IV-2-2-1-Duration of immature stages	129
IV-2-2-2-Consumption of pink bollworm and spiny bollworm eggs by <i>C. undecimpunctata</i> larvae	132
IV-3-Storage of <i>H. tredecimpunctata</i> adults, at two low temperatures, under different regimes of nutrition	143
IV-3-1-Effect of storage for 3 and 5 months at 12 – 14°C on the fecundity and longevity of <i>Hippodamia tredecimpunctata</i> adults	145
IV-3-2-Storage of <i>C. undecimpunctata</i> adults, at two low temperatures under different regimes of nutrition	149

---



IV-3-3-Effect of storage for 3 and 5 months at 12 – 14 °C on the fecundity and longevity of <i>C. undecimpunctata</i> adults	151
IV-3-4-Feeding capacity of <i>H. 13-punctata</i> and <i>C.</i> <i>undecimpunctata</i> after storage	154
IV-4-Survey of some host plants which infested by (PBW) <i>Pectinophora gossypiella</i> and (SBW) <i>Earias insulana</i>	157
IV-5-Survey of parasitoids and predators on pink and spiny bollworms	162
IV-6-Survey and numbers of <i>Pectinophora</i> <i>gossypiella</i> parasitoids during diapause of its larvae	169
IV-7-Survey of Predators	177
IV-8-Percentage of infestation by pink bollworm and spiny bollworm to fruiting structures of cotton plants	182
IV-9-Population dynamics of <i>Pectinophora</i> <i>gossypiella</i> and <i>Earias insulana</i> in cotton fields	191
IV-10-Percentage of infested okra pods and karkadeih capsules by PBW&SBW in both locations	205
IV-11-Comparison between percentage of infestation in cotton, okra and karkadieh in two locations	210

---

IV-12-Population dynamics of PBW and SBW and their predators	214
IV-12-1-Seasonal fluctuations of the two bollworms	214
IV-12-2-Degree of cotton infestation by bollworms	214
IV-12-3-Seasonal abundance of total population of predators	228
IV-12-4-Abundance of total predator	228
IV-12-5-Abundance of different predators	231
IV-12-6-Counts of predators in cotton fields by using net sweeping technique, throughout 1997, 1998 and 1999 season, in untreated and treated fields	253
<b>V-SUMMARY</b>	<b>299</b>
<b>VI-REFERENCES</b>	<b>316</b>
<b>ARABIC SUMMARY (الملخص العربي)</b>	<b>346</b>

\_\_\_\_\_