

# References

- 1-Abraham VS, Swain JA, Forgash AJ, Williams BL and Musulin MM. 2000:**  
Ischaemic preconditioning protects against paraplegia after transient aortic occlusion in the rat.  
Ann Thorac Surg; 69: 475–479.
- 2-Acher CW, Wynn NM, Hoch JR and Popic P. 1994:**  
Combined use of cerebrospinal fluid drainage and naloxone reduces the risk of paraplegia in thoracoabdominal aneurysm repair.  
J Vasc Surg;19(2):236- 246.
- 3-Aebi M. 2005:**  
The adult scoliosis.  
Eur Spine J.;14(10):925-948.
- 4-Akbarnia BA, Marks DS, Boachie-Adjei O, Thompson AG and Asher MA. 2005:**  
Dual growing rod technique for the treatment of progressive early-onset scoliosis: a multicenter study.  
Spine .;30(17):46-S57.
- 5-Anand H, Kulkarni and Ambareesha M. 2007:**  
Scoliosis and Anaesthetic Considerations.  
Indian Journal of Anaesthesia; 51 (6) : 486-495.
- 6-Banoub M, Tetzlaff J and Schubert A. 2003:**  
Pharmacologic and physiologic influences affecting sensory evoked potentials: implications for perioperative monitoring.  
Anaesthesiology.; 99:716-739.
- 7-Barker L. 2002:**  
Positioning on the operating table.  
World Anaesthesia Online; 15(15):180-219
- 8-Bhardwaj A, Northington FJ, Koehler RC, Stiefel T, Hanley DF and Traystman RJ. 1995:**  
Adenosine modulates N-methyl-D-aspartate-stimulated hippocampal nitric oxide production in vivo.  
Stroke; 26: 1627–1633.

**9-Biglioli P, Spirito R and Roberto M. 2000:**

The anterior spinal artery: The main arterial supply of the human spinal cord. A preliminary anatomic study.

J Thorac Cardiovasc Surg; 119: 376–379.

**10-Boezaart AP, Eksteen JA and Spuy GV 1999:**

Intrathecal morphine: Double blind evaluation of optimal dosage for analgesia after major lumbar spinal surgery.

Spine; 24:1131-7.

**11-Borowitz D, Armstrong D and Cemy F. 2001:**

Relief of central airways obstruction following spinal release in a patient with idiopathic scoliosis.

Pediatr Pulmonol; 31:86-88.

**12-Borst HG, Schaudig SA and Rudolph W. 1964:**

Ateriovenous fistula of the aortic arch: repair during deep hypothermia and circulatory arrest.

J Thorac Cardiovasc Surg; 48: 443-447.

**13-Boyer J, Amin N and Taddonio R 1996:**

Evidence of airway obstruction in children with idiopathic scoliosis.

Chest; 109:1532-1535.

**14-Bronwell A K W. 1988:**

Malignant hyperthermia: relationship to other diseases.

British Journal of Anaesthesia; 60: 303-308.

**15-Burns LH, Jin Z and Bowersox SS. 1999:**

The neuroprotective effects of intrathecal administration of the selective N-type calcium channel blocker ziconotide in a rat model of spinal ischaemia.

J Vasc Surg; 30: 334–343.

**16-Cambria R, Davison JK and Zannetti S. 1997:**

Clinical experience with epidural cooling for spinal cord protection during thoracic and thoraco-abdominal aneurysm repair.

J Vasc Surg; 25:234-243.

**17-Cambria RP and Davison JK. 2000:**

Spinal cord ischaemic complications after thoracoabdominal aortic surgery. In: Gewertz BL, editor. Surgery of the aorta and its branches.

Philadelphia: W.B. Saunders ;: 212–228.

- 18-Cernaianu A, Olah A, Cilley Jr, Gaprindashvili T, Gallucci JG and DelRossi AJ. 1993:**  
Effect of sodium nitroprusside on paraplegia during cross-clamping of the thoracic aorta.  
Ann Thorac Surg; 56: 1035–1038.
- 19-Chang CH, Choi SH, Shim YH, Lee KY, Lee HM and Shin YS. 2006 :**  
Optimal endtidal concentration of sevoflurane to test an ankle clonus in children.  
Spine;31 (21): 813-816.
- 20-Christiansson L, Karacagil S, Thelin S and Bergqvist D. 1998:**  
Continuous monitoring of intrathecal pO<sub>2</sub>, pCO<sub>2</sub> and pH during surgical replacement of type 11 thoraco-abdominal aortic aneurysm.  
Eur J Vase Endovasc Surg; 15: 78-81.
- 21-Coillard C, Circo A and Rivard CH. 2008:**  
A new concept for the non-invasive treatment of Adolescent Idiopathic Scoliosis: the Corrective Movement principle integrated in the Spine cord system.  
Disabil Rehabil Assist Technol.;3(3):112-9.
- 22-Cole JW, Murray DJ, Snider RJ, Bassett GS, Bridwell KH and Lenke LG. 2003 :**  
Aprotinin reduces blood loss during spinal surgery in children.  
Spine.;28:2482-2485.
- 23-Cooper DM, Rojas JV and Mellins RB. 1984:**  
Respiratory mechanics in adolescents with idiopathic scoliosis.  
AmRev Respir; 130:16- 22.
- 24-Copley LA, Richards BS, Safavi FZ and Newton PO. 1999:**  
Hemodilution as a method to reduce transfusion requirements in adolescent spine fusion surgery.  
Spine.;24: 219-222.
- 25-Dawson EG, Sherman JE, Kanim LE and Nuwer MR. 1991:**  
Spinal cord monitoring Results of the Scoliosis Research Society and the European Spinal Deformity Society survey.  
Spine; 16: 361-4.
- 26-Day GA, Upadhyay SS and Ho EK. 1994 :**  
Pulmonary functions in congenital scoliosis.  
Spine; 19:1027-31.

**27-de Haan P and Kalkman CJ. 2001:**

Spinal cord monitoring somatosensory and motor-evoked potentials.  
Anesthesiol Clin North America; 19: 923-245.

**28-de Haan P, Vanicky I and Jacobs MJ. 2000:**

Effect of ischaemic pretreatment on heat shock protein 72, neurologic outcome, and histopathologic outcome in a rabbit model of spinal cord ischaemia.  
J Thorac Cardiovasc Surg; 120: 513-519.

**29-Deutsch H, Shrivastava R, Epstein F and Jallo GI. 2001:**

Pediatric intramedullary spinal cavernous malformations .  
Spine; 26: 427-31

**30-Drummond JC. 2000:**

Monitoring depth of anaesthesia: with emphasis on the application of the bispectral index and the middle latency auditory evoked response to the prevention of recall.  
Anaesthesiology.; 93(3): 876-882.

**31-Dunn F and Sharon G. 2008:**

Amazing brace.  
Macleans Magazine.; 3: 22-24

**32-Eckenhoff, J. E. ; Enderby, G. E. H. Larson, Alex; Ediudge, Anthony; and Judvine, D. E. 1963:**

Pulmonary Gas Exchange During Deliberate Hypotension.  
British J. Anaesth; 35: 750-758.

**33-Ehrlich M, Knolle E and Ciovica R. 1999:**

Menantine for prevention of spinal cord injury in a rabbit model.  
J Thorac Cardiovasc Surg; 117: 285-291.

**34-El-Azzazi.H. 2006:**

The practice conduct of anaesthesia.  
A review book, Flashlights on anaesthesia, 1st edition; 2: 3-12

**35-Elmore JR, Gloviczki P and Harper CM Jr. 1992:**

Spinal cord injury in Experimental thoracic aortic occlusion: Investigation of combined methods of protection.  
J Vasc Surg; 15: 789-799.

**36-Florentino-Pineda I, Blakemore LC, Thompson GH, Poe-Kochert C, Adler P and Tripi P. 2001:**

The effect of epsilon-aminocaproic acid on peri-operative blood loss in patients with idiopathic scoliosis undergoing posterior spinal fusion: a preliminary prospective study. *Spine.*; 26:1147-1151.

**37-Follis F, Dragan R and Blisard KS. 1999 :**

Retrograde perfusion of the spinal cord during aortic cross clamping: Initial observations in the swine model.

*J Thorac Cardiovasc Surg*; 118: 597–603.

**38-Freeman III, Canale ST and Beatty JH. 2007:**

Campbell 's. *Operative Orthopaedics*. 11th ed.

Philadelphia, Pa: Mosby Elsevier; *Scoliosis and Kyphosis.* ;23:400-421

**39-Gall O, Aubineau JV and Berniere J. 2001:**

Analgesic effect of low dose intrathecal morphine after spinal fusion in children.

*Anaesthesiology*; 94: 447-52.

**40-Gangemi JJ, Kern JA, Ross SD, Shockey KS, Kron IL and Tribble CG. 2000:**

Retrograde perfusion with a sodium channel antagonist provides spinal cord protection.

*Ann Thorac Surg*; 69: 1744–1748.

**41-Gibson PR. 2004:**

Anaesthesia for correction of scoliosis in children.

*Anaesth Intensive Care.*;32:548-559.

**42-Giglia JS, Zelenock GB and D'Alecy L. 1994:**

Prevention of paraplegia during thoracic aortic cross-clamping: Importance of patent internal mammary arteries.

*J Vasc Surg*; 19: 1044-1051.

**43-Ginsberg MD, Globus MYT and Dietrich R. 1993:**

Temperature modulation in ischaemic brain injury: an aynthesis of recent advances.

*Prog Brain Res*;96:13-22.

**44-Gorgulu A, Kins T and Cobanoglu S. 2000:**

Reduction of edema and infarction by memantine and MK-801 after focal ischaemia and reperfusion in rat.

*Acta Neurochir (Wien)*; 67: 1943–1946.

**45-Greene L . 2001:**

Musculoskeletal Care.  
AAOS,;13: 697 .

**46-Grottke O, Dietrich PJ, Wiegels S and Wappler F. 2004:**

Intraoperative wake up test and postoperative emergence in patients undergoing spinal surgery: a comparison of intravenous and inhaled anaesthetic techniques using short-acting anaesthetics.  
Anaesth Analg.;99: 1521-1527.

**47-Grundy BL. 1983:**

Intraoperative monitoring of sensory evoked potentials.  
Anaesthesiology; 58:72-87.

**48-Guille JT. 2007:**

Fusionless treatment of scoliosis.  
Orthop Clin North Am.;38(4) :541-545.

**49-Hawes M. 2006:**

Impact of spine surgery on signs and symptoms of spinal deformity.  
Pediatr Rehabil, 9(4):318-339.

**50-Hedequist DJ. 2007 :**

Surgical treatment of congenital scoliosis.  
Orthop Clin North Am.;38(4):497-509.

**51-Hell AK, Campbell RM and Hefti F. 2005:**

The vertical expandable prosthetic titanium rib implant for the treatment of thoracic insufficiency syndrome associated with congenital and neuromuscular scoliosis in young children.  
J Pediatr Orthop B.;14:287-293.

**52-Herdmann J, Lumenta CB and Huse KOW. 1993:**

Magnetic stimulation for monitoring of motor pathways in spinal procedures.  
Spine; 18: 551-9.

**53- Herold JA, Kron IL, Langenbur g SE, Blackbourne LH and Tribble CG. 1994:**

Complete prevention of postischaemic spinal cord injury by means of regional infusion with hypothermic saline and adenosine.  
J Thorac Cardiovasc Surg; 107: 536-541.

**54-Hirose K, Okajima K and Taoka Y. 2000:**

Activated protein C reduces the ischaemia reperfusion induced spinal cord injury in rats by inhibiting neutrophil activation.  
Ann Surg; 232: 272–280.

**55-Hoppenfield S, Gross A, Andrews C and Lonner B. 1997 :**

The ankle clonus test for assessment of the integrity of the spinal cord during operations for scoliosis.  
J Bone Joint Surg Am.; 79(2):208-212.

**56-Ishizaki M, Sugiyama S, Uchida H, Nawa S and Shimizu N. 1999:**

Identification and selective perfusion of the spinal cord-feeding arteries by intratechal pO<sub>2</sub> monitoring for spinal cord protection.  
Eur J Vasc Endovasc Surg; 18: 17– 24.

**57-Jacob S. 2002:**

Atlas of human anatomy. Vertebral Column And The Spinal Cord  
Churchill livingstone.; 5: 143-150

**58-Juvonen T, Biancari F, Rimpiläinen J, Satta J, Rainio P and Kiviluoma K. 2002:**

Strategies for Spinal Cord Protection during Descending Thoracic and Thoracoabdominal Aortic Surgery- Up-to-date Experimental and Clinical Results.  
Scand Cardiovasc J; 36:136–160

**59-Kafer ER. 1980 :**

Respiratory and cardiovascular functions in scoliosis and the principles of anesthetic management.  
Anaesthesiology; 52: 339- 51.

**60-Kafer ER. 1977:**

Respiratory and cardiovascular functions in scoliosis. Bull Eur Physio-pathol Respir; 13:299-321.

**61-Kalkman CJ, Drummond JC and Ribberink AA. 1991:**

Low concentrations of isoflurane abolish motor evoked responses to transcranial electrical stimulation during nitrous oxide/opioid anaesthesia in humans.  
Anaesthesia & Analgesia; 73: 410-5.

**62-Kalkman CJ, Drummond JC and Ribbernick AA. 1992:**

Effects of propofol, etomidate, midazolam, and fentanyl on motor evoked responses to transcranial electrical or magnetic stimulation in humans.  
Anaesthesiology; 76: 502-9.

- 63-Kanellopoulos GK, Kato H and Wu Y. 1997:**  
Neuronal cell death in the ischaemic spinal cord: The effect of methylprednisolone *Ann Thorac Surg*; 64: 1279–1285.
- 64-Kannan S, Meert KL, Mooney JF, Hillman-Wiseman C and Warriar I. 2002:**  
Bleeding and coagulation changes during spinal fusion surgery: a comparison of neuromuscular and idiopathic scoliosis patients.  
*Pediatr Crit Care Med.*;3:364-369.
- 65-Kaplan SS, Park TS, Gonzales ER and Gidday JM. 2000:**  
Hydroxyethyl starch reduces leukocyte adherence and vascular injury in the newborn pig cerebral circulation after asphyxia.  
*Stroke*; 31: 2218–2223.
- 66-Karapurkar A, Kudalkar A and Naik L. 2002:**  
Aprotinin, to reduce perioperative blood loss in scoliosis surgery.  
*Indian J Anaesth.*; 46:378-380.
- 67-Katircioglu SF, Ulus AT, Gokce P and Surucu S. 2000:**  
Iloprost protects the spinal cord during aortic cross-clamping in a canine model.  
*J Cardiovasc Surg*; 41: 89–93.
- 68-Kawaguchi M, Sakamoto T and Inoue S. 2000:**  
Low dose propofol as a supplement to ketamine-based anaesthesia during intraoperative monitoring of motor-evoked potentials.  
*Spine*; 25: 974-9.
- 69-Kitagawa K, Matsumoto M and Mabuchi T. 1997:**  
Ischaemic tolerance in hippocampal CA1 neurons studied using contralateral controls .  
*Neuro-science*; 81: 989-998.
- 70-Koshino T, Murakami G, Morishita K, Mawatari T and Abe T. 1999:**  
Does the Adamkiewicz artery originate from the large segmental arteries?  
*J Thorac Cardiovasc Surg*; 117: 898–905.
- 71-Kouchoukos NT, Masetti P, Rokkas CK, Murphy SF and Blackstone EH. 2001:**  
Safety and efficacy of hypothermic cardiopulmonary bypass and circulatory arrest for operations on the descending thoracic and thoracoabdominal aorta.  
*Ann Thorac Surg*; 72: 699–708.

**72-Kouchoukos NT and Rokkas CK. 1999:**

Hypothermic cardiopulmonary bypass for spinal cord protection: rationale and clinical results.

Ann Thorac Surg, Supp ;67 :1940-1942.

**73-Koumbourlis AC. 2006:**

Review: Scoliosis and the respiratory system.

Paed Resp Rev; 7:152-160.

**74-Langeron O, Vivien B and Paqueron X. 1999:**

Effects of propofol, propofol-nitrous oxide and midazolam on cortical somatosensory evoked potentials during sufentanil anaesthesia for major spinal surgery.

Br J Anaesth; 82: 340-5.

**75-Langley SM, Chai PJ, Jagers JJ and Ungerleider RM. 2000 :**

Preoperative high dose methylprednisolone attenuates the cerebral response to deep hypothermic circulatory arrest.

Eur J Cardiothorac Surg; 17: 279–286.

**76-Lapchak PA, Chapman DF, Nunez SY and Zivin JA. 2000:**

Dehydro-epiandrosterone sulfate is neuroprotective in a reversible spinal cord ischaemia model. Possible involvement of GABA receptors.

Stroke; 31: 1953–1957.

**77-Lazorthes G, Gouaze A, Zadeh JO, Santini JJ, Lazorthes Y and Burdin P.**

**1971:** Arterial vascularization of the spinal cord. Recent studies of the anatomic substitution pathways.

J Neurosurg; 35: 253–262.

**78-Lee TC, Yang LC and Chen HJ. 1998 :**

Effect of patient position and hypo-tensive anaesthesia on inferior vena caval pressure.

Spine; 23: 941-7.

**79-Lips J, de Haan P and Bodewits P. 2000:**

Neuroprotective effects of riluzole and ketamine during transient spinal cord ischaemia in the rabbit.

Anaesthesiology; 93: 1303–1311.

**80-Lips L. and Kalkman CJ. 2002:**

Intraoperative spinal cord monitoring: evoked potentials and cerebrospinal fluid oxygenation.

Neurosurgery; 2:842-835.

**81-Lonner, B. S. 2007:**

Emerging minimally invasive technologies for the management of scoliosis.  
Orthop Clin North Am.; 38(3): 431-440.

**82-Lonstein J. 1994:**

The Galveston technique using Luque or Cotrel- Dubousset rods.  
Orthop Clin North Am, 25:311-20.

**83-Lowry KJ, Tobias J and Kittle D. 2001:**

Post operative pain control using epidural catheters after anterior spinal fusion for adolescent scoliosis.  
Spine; 26:1290-3.

**84- Margaret E and Bjorkegren. 1917:**

Handbook of anatomy for students of massage. ( Joints Of The Vertebral Column).  
Bailliere, Tindall & Cox .; 4:89

**85- Maria A. Patestas and Leslie P. Gartner. 2006:**

Textbook of Neuroanatomy  
Blackwell Science Ltd.; 5: 54-60

**86-Matsuyama K, Chiba Y, Ihaya A, Kimura T, Tanigawa N and Muraoka R. 1997:**

Effect of spinal cord preconditioning on paraplegia during cross-clamping of the thoracic aorta.  
Ann Thorac Surg; 63: 1315–1320.

**87-Meert KL, Kannan S And Mooney JF. 2002:**

Predictors of red cell transfusion in children and adolescents undergoing spinal fusion surgery.  
Spine.;27:2137-2142.

**88-Mehta MH. 2005:**

Growth as a corrective force in the early treatment of progressive infantile scoliosis  
J Bone Joint Surg Br; 87 (9): 1237–47.

**89-Melissa A. 2007:**

Anaesthetic implications for surgical correction of scoliosis  
AANA Journal. August; 75(4): 277.

**90-Mendosa-Lattis S and Boezaart AP. 2006 :**

The adult spine. In: Boezaart AP, ed. Anaesthesia and Orthopaedic Surgery. New York, NY: McGraw- Hill.:155-180.

**91-Meylaerts SA, de Haan P, Kalkman CJ, Jaspers J, Vanicky I and Jacobs MJHM. 2000:**

Prevention of paraplegia in pigs by selective segmental artery perfusion during aortic cross-clamping.

J Vasc Surg; 32: 160–170.

**92-Minahan RE, Sepkuty JP and Lesser RP. 2001:**

Anterior spinal cord injury with preserved neurogenic 'motor' evoked potentials.

Clin Neurophysiol; 112: 1442-50.

**93-Motoyama EK, Deeney VF and Fine GF. 2006:**

Effects on lung function of multiple expansion thoracoplasty in children with thoracic insufficiency syndrome: a longitudinal study.

Spine.; 31(3):284-290.

**94-Munyo N, Kaste M and Lindsberg PJ. 1998:**

Tolerization against loss of neuronal function after ischemia-reperfusion injury.

Neuroreport; 9: 321–325.

**95-Murray DJ, Forbes RB, Titone MB and Weinstein SL. 1997:**

Transfusion management in pediatric and adolescent scoliosis surgery: efficacy of autologous blood.

Spine. ;22:2735-2740.

**96-Nakai T, Milusheva E, Baranyi M, Uchihashi Y, Satoh T and Vizi ES. 1999:**

Excessive release of [3H]-noradrenaline and glutamate in response to stimulation of ischaemic conditions in rat spinal cord slice preparation: Effect of NMDA and AMPA receptor antagonists.

Eur J Pharmacol; 366: 143–150.

**97-Nakao Y, Otani H, Yamamura T, Hattori R, Osako M and Imamura H. 2001:**

Insulin-like growth factor 1 prevents neuronal cell death and paraplegia in the rabbit model of spinal cord ischaemia.

J Thorac Cardiovasc Surg; 122: 136–143.

**98-Nakashima K, Todd MM and Warner DS. 1995:**

The relationship between cerebral metabolic rate and ischaemic depolarization. A comparison of the effects of hypo-thermia, pentobarbital and isoflurane.

Anaesth-esiology;82:1199-1208.

**99-Nicole O, Ali C and Docagne F. 2001:**

Neuroprotection mediated by glial cell linederived neurotrophic factor: Involvement of a reduction of NMDA-induced calcium influx by the mitogen-activated protein kinase pathway.  
J Neurosci; 21: 3024–3033.

**100-O'Rahilly, Muller, Carpenter and Swenson. 1982:**

Basic human anatomy, The vertebral column  
W B Saunders.; 39: (411- 432)

**101-Owen JH. 1999 :**

The application of intraoperative monitoring during surgery for spinal deformity.  
Spine; 24: 2649-62.

**102-Papastefanou SL, Henderson LM and Smith NJ. 2000 :**

Surface electrode somatosensory-evoked potentials in spinal surgery.  
Spine; 25: 2467-72.

**103-Park CK. 2000:**

The effect of patient positioning on intraabdominal pressure and blood loss in spinal surgery.  
Anaesth Analg; 91: 552-7.

**104-Parke WW, Whalen JL, Bunger PC and Settles HE. 1995:**

Intimal musculature of the lower anterior spinal artery.  
Spine; 20: 2073–2079.

**105-Parrino PE, Kron IL and Ross SD. 2000:**

Retrograde venous perfusion with hypo-thermic saline and adenosine for protection of the ischaemic spinal cord.  
J Vasc Surg; 32: 171–178.

**106-Parrino PE, Kron IL and Ross SD. 1999:**

Spinal cord protection during aortic cross clamping using retrograde venous perfusion.  
Ann Thorac Surg; 67: 1589–1594.

**107-Patil CG, Santarelli J and Lad SP. 2008:**

Inpatient complications, mortality, and discharge disposition after surgical correction of idiopathic scoliosis: a national perspective.  
Spine J.; 45:498-534

- 108-Pehrsson K, Larsson S, Oden A and Nachemson A. 1992:**  
Long-term follow-up of patients with untreated scoliosis. A study of mortality, causes of death, and symptoms.  
Spine; 17: 1091-6.
- 109-Pelosi L, Lamb J, Grevitt M, Mehdian SM, Webb JK and Blumhardt LD. 2002:**  
Combined monitoring of motor and somatosensory evoked potentials in orthopaedic spinal surgery.  
Clin Neurophysiol.;113:1082-1091.
- 110-Pelosi L, Stevenson M and Hobbs GJ. 2001:**  
Intraoperative motor evoked potentials to transcranial electrical stimulation during two anaesthetic regimens.  
Clin Neurophysiol; 112: 1076-87.
- 111-Petaja J, Pesonen E and Fernandez JA. 2001:**  
Activated protein C and inflammation in human myocardium after heart surgery.  
Am J Hematol; 67: 210-212.
- 112--Peter de Haan, Cor J. Kalkman, and Michael J.H. M and Jacobs. 2001:**  
Pharmacologic Neuroprotection in Experimental Spinal Cord Ischaemia.  
A systematic review. Journal of Neurosurgical Anaesthesiology;13(1): 3–12
- 113-Qayumi AK, Janusz MT and Dorovini-Zis K. 1994:**  
Additive effect of allopurinol and deferoxamine in the prevention of spinal cord injury caused by aortic cross-clamping.  
J Thorac Cardiovasc Surg; 107: 1203–1209.
- 114-Raw DA, Beattie JK and Hunter JM. 2003:**  
Review article: Anaesthesia for spinal surgery in adults.  
British Journal of Anaesthesia; 91:886-904.
- 115-Reamy BV and Slakey JB. 2001:**  
Adolescent idiopathic scoliosis: Review and current concepts.  
Am Fam Med,; 64:1-10
- 116-Relton JE and Hall JE. 1967:**  
An operation frame for spinal fusion: a new apparatus designed to reduce haemorrhage during operation.  
J Bone Joint Surg Br. ;49:327-332.

**117-Renshaw TS. 1988:**

Role of Harrington instrumentation and posterior spine fusion in the management of AIS.

Orthop Clin N Am, 19:257-269.

**118-Reuben SS and Connelly NR. 2000:**

Postoperative analgesic effects of celecoxib or rofecoxib after spinal fusion surgery.

Anaesth Analg; 91: 1221-5.

**119-Reuter DG, Tacker WA Jr, Babbs CF, Badylak SF, VoorheesWD 3<sup>rd</sup> and Konrad PE. 1995:**

Preliminary results of deferoxamine and L1 treatment of spinal cord ischaemia.

J Thorac Cardiovasc Surg; 109: 1017–1019

**120--Richardson ML. 2006:**

Approaches to differential diagnosis in musculoskeletal imaging.

Univ. of Washington School of Medicine 8:56-89

**121--Robinson, C. S. B. Galasko, C. Delaney, J. B. Williamson, and J. L. Barrie1995:**

Scoliosis and lung function in spinal muscular atrophy.

Eur Spine J; 4:268-273.

**122-Robinson, J. S. 1967:**

Hypotension Without Hypoxia.

Internat. Anaesth. Clin. 5: 467-480.

**123-Rodola F, D'Avolio S and Chierichini A. 2000:**

Wake-up test during major spinal surgery under remifentanyl balanced anaesthesia.

Eur Rev Med Pharmacol Sci; 4: 67-70.

**124-Rokkas CK, Helfrich LR and Lobner DC. 1995 :**

Dextrophan inhibits the release of excitatory amino acids during spinal cord ischaemia.

J Thorac Cardiovasc Surg;110: 27-35.

**125-Rokkas CK and Kouchoukos NT. 1998:**

Profound hypothermia for spinal cord protection in operations on the descending thoracic and thoracoabdominal aorta.

Semin Thorac Cardiovasc Surg;10(1):57-60.

**126-Rollason, W. N., and Hough, J. M. 1969:**

A Re-examination of some electrocardio graphic studies during hypotensive anaesthesia. The effect of rate of fall of blood pressure.  
British J. Anaesth; 41: 985-986.

**127-Rose PS and Lenke LG. 2007:**

Classification of Operative Adolescent Idiopathic Scoliosis: Treatment Guidelines.  
Orthop Clin N Am.;38:521-529.

**128-Rowbotham DJ, Alexander J and Arden C. 1998:**

Guidelines for the use of non-steroidal anti-inflammatory drugs in the perioperative period.  
The Royal College of Anaesthetists, UK;45:69-70

**129-Russ W, Kling D and Loesevitz A. 1984:**

Effect of hypothermia on visual evoked potentials in humans.  
Anaesthesiology; 61:207.

**130-Sakurai M, Abe K and Hayashi T. 2001:**

Adenovirus-mediate d glial cell linederived neurotrophic factor gene delivery reduces motor neuron injury after transient spinal cord ischaemia in rabbits. J Thorac Cardiovasc Surg; 120: 1148–1157.

**131-Salem, M. R.; Ivankovic, A. D.; and Shaker, M. H. 1971:**

Safety factors in deliberately induced hypotension.  
Middle East J. Anaesth; 3: 107-119

**132-Salzano R, Ellison LH and Altonji DF. 1994:**

Regional deep hypothermia of the spinal cord protects against ischaemic injury during thoracic aortic cross-clamping.  
Ann Thorac Surg;57:65-70.

**133-Samra SK, Dy EA and Welch KB. 2001:**

Remifentanil and fentanyl based anaesthesia for intraoperative monitoring of somatosensory evoked potentials.  
Anaesth Analg; 92: 1510-5.

**134-Sarah M. and Charles E. Seay, Martha and Pat Beard. 2007**

scoliosis genetic research  
American Journal of Human Genetics.;22:236-278

**135-Sarwark J and Sarwahi V. 2007:**

New strategies and decision making in the management of neuromuscular scoliosis.

Orthop Clin North Am.;38(4): 485-496.

**136-Sessler DI. 1997:**

Mild Perioperative hypothermia.

N Engl J Med. 336:1730-1737.

**137-Sharrock NE, Beckman JD, Connolly E and Savarese JJ. 2005:**

Anaesthesia for Orthopedic Surgery, Text book of Anaesthesia by Miller RD, 6th edition.

Elsevier Churchill Livingstone; 61: 466-509

**138-Shibata K, Takamoto S and Kotsuka Y. 2001:**

Doppler ultrasonographic identification of the critical segmental artery for spinal cord protection.

Eur J Cardiothorac Surg; 20: 527–532.

**139--Simpson J, Eide T and Newman S. 1996:**

Trimethaphan versus sodium Nitroprusside for the control of proximal hypertension during thoracic aortic cross-clamping: The effects on spinal cord ischaemia.

Anaesth Analg; 82: 68–74.

**140-Sinclair J R and Mason R A. 1984:**

Ankylosing Spondylitis; the case for awake intubation.

Anaesthesia; 39: 3-11

**141--Skagg S. 1996:**

Adolescent idiopathic scoliosis , The family physician's role in adolescent idiopathic scoliosis

Am Fam Physician; 53(7):2327.

**142--Sliwa JA and MacLean IC. 1992:**

Ischaemic myelopathy: A review of spinal vasculature and related clinical syndromes.

Arch Phys Med Rehabil; 73: 365–372.

**143-Spielholz NI, Benjamin MV and Engler GL. 1979:**

Somatosensory evoked potentials during decompression and stabilization of the spine: methods and findings.

Spine; 4:500.

- 144- Suzer T, Coskun E, Islekel H and Tahta K. 1999:**  
Neuroprotective effect of magnesium on lipid peroxidation and axonal function after experimental spinal cord injury.  
Spinal Cord; 37: 480–484.
- 145-Svensson LG, Hess KH, D'Agostino RS and Entrup MH. 1998:**  
Reduction of neurologic injury after high-risk thoracoabdominal aortic operation.  
Ann Thorac Surg ;66(1):132-138.
- 146-Svensson LG, Von Ritter CM and Groeneveld HT. 1986 :**  
Cross-clamping of the thoracic aorta. Influence of aortic shunts, laminectomy, papaverine, calcium channel blocker, allopurinol , and superoxide dismutase on spinal cord blood flow and paraplegia in baboons .  
Ann Surg; 204: 38–47
- 147-Szabo MD and Crosby G. 1988:**  
Effect of profound hypermagnesemia on spinal cord glucose utilization in rats.  
Stroke; 19: 747–749.
- 148-Szilagyi DE, Hageman JH and Smith RF. 1978:**  
Spinal cord damage in surgery of the abdominal aorta.  
Surgery;83:38-56.
- 149-Tatu Juvonen, Fausto Biancari, Jussi Rimpiläinen, Jari Satta, Pekka Rainio and Kai Kiviluoma.2002:**  
Strategies for spinal cord protection during descending thoracic and thoracoabdominal Aortic surgery : Up-to-date Experimental and Clinical Results .  
Scand Cardiovasc J; 36: 136–160.
- 150-Taylor JM and Gropper MA. 2006:**  
Critical care challenges in orthopedic surgery patients.  
Crit Care Med; 34: 191-199.
- 151-Tetik O , Islamoglu F and Yagdi T. 2001:**  
An intraaortic solution trial to prevent spinal cord injury in a rabbit model.  
Eur J Vasc Endovasc Surg; 22: 175– 179.
- 152-Tetzlaff JE and Yoon HJ. 1995:**  
Clinical Orthopedic Anaesthesia, ed.  
Butterworth Heinemann; 22: 233-300

**153-Thomas S, McNeill, Ronald L. Dewald, Kenn. Kuo, Edward J. Bennett and M. R. Salem. 1974:**

Controlled Hypotensive Anaesthesia in Scoliosis Surgery. *J Bone Joint Surg Am.*;56:1167-1172.

**154-Tobias JD, Hoernschemeyer DG and Anderson JT. 2009:**

Ankle clonus and wakeup tests during posterior spinal fusion: Correlation with bispectral index.  
*Am J Orthop.*; 38(4):75-77.

**155-Toyota S and Amaki Y. 1998:**

Hemodynamic evaluation of the prone position by transoeso-phageal echo.  
*J Clin Anaesth*; 9:32-35.

**156-Ueno T, Furukawa K, Katayama Y and Itoh T. 1994 :**

Protection against ischaemic spinal cord injury: One-shot perfusion cooling and percutaneous topical cooling.  
*J Vasc Surg*; 19: 882–887.

**157-Valadka AB, Gopinath SP, Contant CF, Uzura M and Robertson CS. 1998 :**

Relationship of brain tissue P02 to outcome after severe head injury. *Critical Care Med*; 26: 1576-81.

**158-Valentino K, Newcomb R and Gadbois T. 1993:**

A selective N-type calcium channel antagonist protects against neuronal loss after global cerebral ischaemia.  
*Proc Natl Acad Sci USA*; 90: 7894–7897.

**159-Vedantam R, Lenke LG and Bridwell KH. 2000:**

A prospective evaluation of pulmonary function in patients with adolescent idiopathic scoliosis relative to the surgical approach used for spinal arthrodesis.  
*Spine*; 25:82-90.

**160-Vedantam R, Lenke LG and Bridwell KH. 2000:**

A prospective evaluation of pulmonary function in patients with adolescent idiopathic scoliosis relative to the surgical approach used for spinal arthrodesis.  
*Spine*; 25: 82-90.

**161-Wada T, Yao H, Miyamoto T, Mukai S and Yamamura M. 2001:**

Prevention and detection of spinal cord injury during thoracic and thoracoabdominal aortic repairs.  
*Ann Thorac Surg*; 72: 80–84.

- 162-Waldhausen JH, Redding GJ and Song KM. 2007:**  
 Vertical expandable prosthetic titanium rib for thoracic insufficiency syndrome: a new method to treat an old problem.  
 J Pediatr Surg.;42(1):76-80.
- 163-Wazeka AN,DiMaio MF and Boachie-Adjei O. 2004:**  
 Outcome of pediatric patients with severe restrictive lung disease following reconstructive spine surgery.  
 Spine; 29:528-34.
- 164-Weibl H. 1973:**  
 Zur Topographie der Medulla spinalis der Albinoratte (Rattus Norvegicus). Adv Anat Embryol Cell Biol.; 47: 6.
- 165-Weinstein S L, Zavala D C and Ponsetti I V. 1981:**  
 Idiopathic scoliosis: long term follow up and prognosis in untreated patients.  
 Journal of Bone and Joint Surgery (Am); 63A: 702-712.
- 166-Weiss HR and Goodall D. 2008:**  
 Rate of complications in scoliosis surgery.  
 a systematic review of the Pub Med literature. Scoliosis; 3: 9
- 167-Wiess HR, Goodall D, Bess S, Wong MS, Patal V and Burger E. 2008:**  
 Adolescent idio-pathic scoliosis to operate or not.  
 A debate article of BioMed Central. Patient safety in surgery, 2:25.
- 168-Winnerkvist A, Bartoli S, Iliopoulo s DC, Hess KR, Miller CC 3<sup>rd</sup> and Sa. HJ. 2002:**  
 Spinal cord protection during aortic cross clamping: Retrograde venous spinal cord perfusion, distal aortic perfusion, cerebrospinal fluid drainage.  
 Scand Cardiovasc J; 36: 6–10.
- 169-Wisselink W, Patetsios P and Panetta TF. 1998:**  
 Medium molecular weight penta-starch reduces reperfusion injury by decreasing capillary leak in a model of spinal cord ischaemia.  
 J Vasc Surg; 27: 109–116.
- 170-Yagita Y, Kitagawa K, Ohtsuki T, Tanaka S, Hori M and Matsumoto M. 2001 :**  
 Induction of the HSP110/105 family in the rat hippocampus in cerebral ischaemia and ischaemic tolerance.  
 J Cereb Blood Flow Metab; 21: 811–819.

**171-York DH, Chabot RJ and Gaines RW. 1987:**

Response variability of somatosensory evoked potentials during scoliosis surgery.

Spine; 12: 864-76.

**172-Yuan N, Fraire JA, Margetis MM, Skaggs DL, Tolo VT and Keens TG. 2005:**

The effect of scoliosis surgery on lung function in the immediate postoperative period.

Spine.;30(19):2182-2185.

**173-Zikria BA, Subbarao C and Oz MC. 1989:**

Macromolecules reduce abnormal micro-vascular permeability in rat limb ischaemia reperfusion injury.

Crit Care Med; 17: 1306–1309.

**174-Zvara DA, Colonna DM, Deal DD, Vernon JC, Gowda M and Lundel JC. 1999:**

Ischaemic preconditioning reduces neurologic injury in a rat model of spinal cord ischaemia.

Ann Thorac Surg; 68: 874–880.