

References

- Abou-Hussein HH, El-Khawaga AM, Wahab AA(2005)* :C-reactive protein as a marker for early-onset neonatal sepsis: The good, the better and the best. The Egyptian Journal of Neonatology ; 6 (3): 151-159.
- Ahrens P, Kattner E, Kohler B, et al.(2004)*: Mutations of genes involved in the innate immune system as predictors of sepsis in very low birth weight infants. *Pediatr Res* 2004; 55: 652-56.
- Anwr, S, Mustafa, S. and Ashraf, S.(2000)*: Rapid identification of neonatal sepsis. *J. Pak. Med. Assoc.* 45: 94.
- Apgar B.S., Greenberg G., and Yen G. (2005)*: Prevention of group B streptococcal Disease in the Newborn. *Am fam physician*; 71 (5): 903 – 10. *Arch Dis Child*, in press.
- Atli, D., and Petur,L. : Neonatal sepsis(2005)*: diagnosis and initial antibiotic therapy. *Pediatr . clin. North Am.* 34: 219.
- Auriti, C., Maccallini. A. and Diliso, D.(2003)*: Risk factors for nosocomial Infection in a neonatal intensive care unit. *J. Hosp. Infect.* 53: 30.
- Bacher, C(2005)*: Group- B streptocoecal infection. *Crit. Care med.* 27: 120.

- Baltimore R.S. (2003):** Neonatal sepsis: Epidemiology and management. *Pediatr Drugs*; 5(11): 723 – 40.
- Baungart, S., Halls, E. and Campos, J.(2004):** Sepsis with coagulase negative staph in critically ill newborn. *Am. J. Dis. Child.* 137: 661.
- Berger, M.(2000):** Complement deficiency and neutrophil dysfunction as a risk factors for bacterial infections in newborn. *Rev. inf. Dis.* 12: 405.
- Berner R, Niemeyer CM, Leititis JU ,et al.(1998):** Plasma levels and gene expression of granulocyte colony stimulating factor, tumor necrosis factor alpha, IL-1 beta, IL-6, IL-8 and soluble intercellular adhesion molecule-1 in neonatal early onset sepsis. *Pediatr Res* ; 44: 469–77.
- Betty C and Inderpreet S.(2005):** Early onset neonatal sepsis. *The Indian Journal of Pediatrics* 2005; 72 (1): 23-26.
- Bonac B, Dergance M, Wraber B and Hojker S(2000):** Interleukins and procalcitonin in early diagnosis of severe bacterial infection in critically ill neonate. *Plugers Arch*;2(5 Suppl.)R:72-74.
- Caplan, E., Oed, L. and Kassel, R.(2005):** Cytokines serum level in neonatal sepsis. *Ann Surg.* 15: 210.
- Carrigan SD, Scott G and Tobrizian M.(2004):** Toward resolving the challenges of sepsis diagnosis. *Clinical Chemistry* ; 50: 1301-1314.

- Chiesa C, Panero A, Osborn JF, et al.(2004):*** Diagnosis of neonatal sepsis: A clinical and laboratory challenge. *Clinical Chemistry* ; 50 (2): 279-287.
- Combes A, Figliolini C, Trouillet JL, et al.(2002):*** Incidence and outcome of polymicrobial ventilator-associated pneumonia. *Chest* ;121:1618-1623.
- De Benedetti F, Auriti C, D'urbano LE, et al.(2007):*** Low serum levels of mannose binding lectin are a risk factor for neonatal sepsis. *Pediatr Res* 2007; 61: 325-328.
- Dzwonek AB, Neth OW, Thiebaut R, et al.(2008):*** The Role of Mannose-Binding Lectin in Susceptibility to Infection in Preterm Neonates. *Pediatr Res* ; 63: 680-685.
- Edward, M., Jackson, C. and Baker, C.(2001):*** Twin pregnancies, incidence of group- B streptococcal colonization and disease. *J. Pediat.* 97: 635.
- Edwards M.S. and Baker C.J. (2004):*** Sepsis in the newborn. In *Diseases of children*, 11th ed: 545 – 61. Edited by Gershon A. A., Hotez P.J., and Katz S.L. By Mosby.
- Ehl S, Gering B, Bartmann P and Pohlandt F.(1997):*** C-reactive protein is a useful marker for guiding duration of antibiotic therapy in suspected neonatal bacterial infection. *Pediatrics* ; 99 (2): 216-221.

EL-Kerdani T, Abd-ELWahed MA, ALy GS, et al. (2001): The value of neutrophil CD11b expression, interleukin-6 and soluble receptors of tumor necrosis factor in early diagnosis of neonatal sepsis. *The Egyptian Journal of Neonatology*; 2(2) :115-125.

Engle, W., Rosenfield, C. and Mouzinho, A. (2005): Circulating neutrophil in septic preterm neonates. *Pediatrics*. 99: 10, 2005.

Eskander R.M., Badrawi N.H., El-Atter S. et al. (2003): Epidemiological study of neonatal sepsis in a neonatal intensive care unit. M.D. Thesis. *Exp Mol Pathol*; 75 (1): 45 – 52.

Fanaroff, A., Korores, S. and Wright, L. (2005): Incidence, presenting features, risk factors and significance of late onset septicemia in very low birth weight infants. *Pediatr. Infect. Dis*. 17: 598.

Ferrieri, P. (2006): Neonatal susceptibility & immunity to major bacterial pathogens. *Rev. Infect. Dis*. 4: 5394.

Franz A R, Steinbach G, Korn M, and Pohlandt F. (2001): "Interleukin-8: a valuable tool restrict antibiotic therapy in newborn infants," *Acta paediatrica*, vol. 90, no. 9, pp. 1025-1032.

Freij B.J. and McCracken G.H. (1999): Acute infections. In the textbook of *Neonatology: pathophysiology and Management of the Newborn*, 5th ed: 1189 – 230. Edited by Avery G.B., Fletcher M.A., and

MacDonald M.G. By Lippincott Williams & Wilkins.

Garcia-parts J(2000): The critically ill neonate with infection: management considerations in the term and preterm infant. *Pediatr infect Dis* : 11 : 4.

Gerdes JS(2004): Diagnosis and management of bacterial infections in the neonate. *Pediatr Clin N Am* ; 51: 939-59.

Ghosh, A., Mittal, M and Jaganthen, G(2004): Early diagnosis Of neonatal sepsis. *Inian J. Med. Sci.* 55: 500.

Gomella TL, Cunningham MD, Eyal FG and Zenk KE. (2004):Neonatal sepsis. In: *Lange Clinical Manual of Neonatology, 5th ed., Lange Medical Books/McGraw-Hill* ; 68: 434-442.

Gottoff S.P. (2000): Infections of the newborn infant. In *Nelson Textbook of Pediatrics, 16th edition: 538 – 52.* Edited by Behrman R.E., Kliegman R.M., and Jenson H.B. By W.B. Saunders.

Guida, J., Runing, A. and Leaf, K(2003): Platelet count and sepsis, is there an organism specific response. *Pediatrics.* 11:315.

Hague, K(2002): Infection & immunity in the newborn. *Lancet. Infect. Dis.* 12: 316.

Hebra, A., Diello, C. and Russil, A(2006): Necrotizing enterocolitis. *JAMA.* 5: 874.

Hengst JM.(2003): The role of C-reactive protein in the evaluation and management of infants with suspected sepsis. *Adv neonatal Care* ; 3 (1): 3-13.

Horisberger T,Harbarth S,Nadal D,et al.(2004):G-CSF and IL-8 for early diagnosis of sepsis in neonates and critically ill children -safety and cost effectiveness of a new laboratory prediction model :study protocol of a randomized controlled trial,*Critical care*,Vol .8,no.6,pp.443-450.1

Hsu KK, Pelton SI and Shaprio DS.(2003): Detection of group B streptococcal bacteremia in simulated intrapartum antimicrobial prophylaxis. *Diagn Microbiol Infect Dis* ; 45: 23-7.

Imam SS, Saafan HA and Amin SK.(2004): A new modality for early diagnosis and severity prediction in necrotizing enterocolitis. *The Egyptian Journal of Neonatology* ; 5 (1): 10-13.

Jacobson, J.A., Chester, T.J. and Fraser, D.W.(2006): Bacterial infection of newborn. *Clin. Inf. Dis.* 26: 187.

Jerome, O (2001):Bacterial sepsis and meningitis. *J. pediatr.* 5: 966, 2001.

Kemp, A. and Campbell, D.(2006): The neonatal immune system. *J. Pediatr.* 101: 1025.

Klein J. O. and Remington J.S. (2001): Current concepts of infections of the fetus and newborn infant. In Infectious Diseases of the fetus and Newborn infant 5th ed: 1 – 24. Edited by Remington J.S and Klein J.O. By W.B. Saunders.

Klein OJ.(2001): Bacterial sepsis and meningitis.In:Remington Js, Klein OJ,eds. Infectious disease of the faetus and Newborn Infant 5 th ed .philadelphia : W.B. Saunders ;943-98.

Klein, J.(2005): Bacterial sepsis and meningitis. Med. Clin. N. Am. 4: 90.

Kocabas E, Sarikcioglu A, Aksaray N, et al.(2007):"Role of procalcitonin, C-reactive protein, interleukin-6, interleukin-8 and tumor necrosis factor – α in the diagnosis of neonatal sepsis," Turkish Journal of Pediatrics, vol. 49, no. 1, pp. 7-20.

Kumar Y, Qunibi M, Neal TJ and Yoxal CW.(2001): Time to positivity of neonatal blood cultures. Arch Dis Fetal Neonatal Ed ; 85: F182-F186.

Laborada G,Rego M,Jain A, et al.,(2003):Diagnostic value of cytokines and C-reactive protein in the first 24 hour of neonatal sepsis. American Journal of perinatology, volume 20,Number 8,pp 491-501.

- LedurAF, Fitting C, David B, et al., (1995):** Variable estimation of cytokine levels produced by commercial ELISA Kits results using international cytokine standards. *J. Immunol. methods*, 186: 171-179.
- Lee, C. Tang, R and Ching, R (2005):** Evaluation of different blood culture media in neonatal sepsis. *J. Microbiol. Immunol. Infect.* 33: 165.
- Levi M, Tencate L, H, Vander Poll, et al. (2000):** Pathogenesis of thrombocytopenia and disseminated intravascular coagulation in sepsis. *Journal of American Medical Association*, 270: 975-9.
- Lewis, D. (2005):** Developmental immunology and role of host defense in neonatal susceptibility to infection. *J. pediatr.* 108: 11.
- Linda, B (2002):** Neonatal sepsis, the scientific basis for practice. *Neonatal network.* 10: 40, 2002.
- Lock, G (2001):** Infection, antibiotics and preterm delivery. *Clin. Infect. Dis.* 25: 309.
- Lott, W. (2006):** Neonatal bacterial sepsis, *crit. Care. Nurse.* *Clin. North Am.* 15: 65.
- Luck S, Torny M, d'Agapeyeff K, et al. (2002):** Horizontal transmission of *Candida parapsilosis* candidemia in a neonatal intensive care unit. *J Clin Microbiol* 2002; 40 (7): 2363-69.

M. Pamela Griffin and J. Randall Moorman (2001):Lessons from E coli : from basic research to clinical aspects. Villars sur Ollon, Septembre2005; DOI: 10.1542/peds.107.1.97 Pediatrics ;107;97-104.

Maisels, M., Khare, S. and Kring, P.(2002): Risk of sepsis in newborn with severe hyperbilirubinemia. Pediatrics. 90: 741.

Malik A, Hui CPS, Pennie RA, Kirpalani H (2003): Beyond the complete blood cell count and C-reactive protein. A systematic review of modern diagnostic tests for neonatal sepsis. Arch. Pediatr. Adolesc. Med. 157: 511-516.

Manucha V, Rusia U, sikka M,et al.(2002): "Utility of haematological parameters and C-reactive protein in the decetion of neonatal sepsis, "Journal of paediatrics and child Health, vol. 38, no. 5, pp. 459 – 464.

Martin H, Olander B and Norman M(2001): Reactive hyperemia and interleukin-6, interleukin-8 and tumor necrosis factor- α in diagnosis of early onset neonatal sepsis.Pediatrics;108(4):E61.

Martin H,O Lander B,Norman M,(2001):Reactive hyperemia and Interleukin 6,Interleukin 8,Tumor Necrosis Factor $-\alpha$ In the Diagnosis of Early-onset Neonatal sepsis .pediatrics vol.108 No.4,pp1-6.

- Mustafa S, Farooqui S, Waheed S and Mahmoud K. (2005):***
Evaluation of C-reactive protein as early indicator of blood culture positivity in neonates. Pak J Med Sci ; 21 (1): 69-37.
- Ng PC.(2004):*** Diagnostic markers of infection in neonates. Arch Dis Child Fetal Neonatal Ed ; 89(3): F229-F235.
- Noya, F(2002):*** Prevention of GBS infection. Infect. Disease. Clin. North Am. 6: 41.
- Nupponen I,Andersson S,Jarvenpaa AL, et al. (2001):***
Neutrophil CD11b expression and circulating interleukin-8 as diagnostic markers for early onset neonatal sepsis.Pediatrics;108(1):E12.
- Ohlsson A. and Lacy J.B., (2004):*** Intravenous immunoglobulin for suspected or subsequently proven infection in neonates. Cochrane Database Syst Rev; (1) CD 001239.
- Osrin D,vergnano S,CostelloA(2004):***Serious Bacterial Infections In Newborn Infants In developing countries.Curropin Infect dis 2004;17:217-224.
- Ottolini, M., Lungren, K. and Cason, S.(2003):*** Utility of complete blood count and blood culture screening in the diagnosis of neonatal sepsis Pedat. Infect. Dis. J. 17: 235.

- Persson, E., Trollfors, B. and Brendberg L.(2002):*** Septicemia and meningitis in neonates and during infancy. Acta, Pediatr. 10: 92.
- Polin, R(2003):*** The "Ins and Outs" of neonatal sepsis. J. Pediatr. 143: 8.
- Powel, K.(2005):*** Laboratory aids for diagnosis of neonatal sepsis. J. Clin. Invest. 4: 1240.
- Procianoy and Silveira RC.(2004):*** The role of sample collection timing on interleukin-6 levels in early-onset neonatal sepsis. J Pediatr (Rio J) ; 80 (5): 407-10 .
- Ragharen, M., Mandal, G. and Bhat, B.(2002):*** Prenatal risk factors in neonatal infections. Ind. J. Pediatr. 3: 335.
- Rennie J.M., and Roberton N.R., (2002):*** Infection. InA Manual of Neonatal Intensive Care 4th ed: 223 – 67. By Arnold and member of the Holder Headline Group.
- Rubin LG, Sanchez PJ, Siegel J, Levine G, Saiman L, Jarvis WR and Pediatric Prevention Network (2002):*** Evaluation and treatment of neonates with suspected late onset sepsis: A survey of neonatologist practice. Pediatrics; 110: 1-7.
- Rudnick C.M. and Hoekzema G.S. (2002):*** Neonatal Herpes Simplex Virus Infections, Am Fam Physician. 2002 Mar 15; 65 (6): 1138 – 42.

Saizous, C., Famoux, C. and Bingen, E(2001): Severe neonatal bacterial infections. Arch Pediat. 4: 725.

Santana RC, Garcia M F, Reyes D et al.,(2003): Role of Cytokines (and Interleukin -1 β , 6, 8, Tumor Necrosis Factor α ,and Soluble Receptor of Interleukin -2 and C-reactive protein in the Diagnosis of Neonatal sepsis. Acta paediatr ;92:221-227.

Santana Reyes C, Garcí'a-Muñoz F, Reyes D, et al.(2003): Role of cytokines (interleukin-1 β , 6, 8, tumor necrosis factor- α , and soluble receptor of interleukin-2) and C-reactive protein in the diagnosis of neonatal sepsis. Acta Pædiatr ; 92: 221–227. Stockholm. ISSN 0803-5253.

Seaward, P., Hannah, M. and Myhr, T.(2006): International multicenter term PROM study: Evaluation of infection born to patients with premature rupture of membranes at term Premature rupture of membranes. Obstet. Gynecol. 635: 9.

Sheperd, C., Fisher, M. and Rosentein, N(2003): Neonatal meningococcal disease in the United States. Pediat. Infect. Dis. J. 20: 31.

Sinha A, Yokoe D and Platt R.(2003): Epidemiology of neonatal infections: experience during and after hospitalization. Pediatr Infect Dis J ; 22: 244-251.

Soman, M., Green, B. and Daling, J.(2005): Risk factors for early neonatal sepsis. Am. J. Epidemiol. 121: 712.

Spear ML, Stefano JL, Fawcett P, Proujansky R.(1995):
Soluble interleukin-2 receptor as a predictor of
neonatal sepsis. J Pediatr ; 126: 982–5.

Stangno, S. (2004): Cytomegalovirus. In Nelson textbook of
pediatrics, 17th edition: 1066 – 9 edited by Behrman
R.E. Kliegman R.M., and Jenson H.B. by W.B.
Saunders.

Stoll B(2008): Infections in neonates: Etiology of fetal and
neonatal infection. In Nelson text book of pediatrics,
18th ed., Kliegman, Beharman, Janson & Stanton
(eds); Sanders; Philadelphia, USA..Chapter 109. pp
794 – 811.

Stoll BJ.(2008): Infections of the Neonatal Infant. In: Kliegman
RM, Behrman RE, Jenson HB and Stanton BF (eds).
Nelson Textbook of Pediatrics, 18th ed. Elsevier ;
109: 794-811.

Suri M., Van de Ven C., and Cairo M.S. (2003):
Immunotherapy in the prophylaxis and treatment of
neonatal sepsis, Curr Opin Pediatr; 15: 155-60.

Taylor, PC (2001) : Mol. Biotechnol 19(2)153-168.

Weinberg GA, Powel KR (2001): Laboratory aids for diagnosis
of neonatal sepsis. Remington, JS, Klein Jo eds.
Infectious diseases of the fetus and newborn infants.
P: 1327-1344. WB Saunders, Philadelphia.

WHO (2003): Recommendations for first line empiric therapy
of serious neonatal infections.

Yoder, M.(2001): Clinical Gram-positive sepsis: does it fundamentally differ from Gram-negative bacterial sepsis. J. Pediatr. 18: 340.

Zaidi A.K., Huskins W.C., Thaver D., Bhutta Z. et al., (2005): Hospital – acquired neonatal infections in developing countries. Lancet; 365: 1175 – 88.