

REFERENCES

Abyliolm, F.E., Bergland, O. and Semb, G. (1981): Secodary bone grafting of alveolar clefts. Scand.J. Plastic Reconsttr. Surg. 15:127-140.

Abyliolm, Borcbgrevink, H.H.C. and Eskeland, G. (1979): Palatal fistulae following cleft palate surgery. Scandinavian J. of plastic and Reconstructive surgery, 13: 295-300.

Adams, D.A. (1987): The causes of deafness. In Scott-Brown's Otolaryngology, 5th ed., volume 6, chapter 4, pp. 35-53. London : Butterworths.

Albrektsson, T. and Linder, L. (1981): Intravital, long term follow-up of autologous experimental bone grafts. Archives of Orthopedic and Traumatic Surgery, 98: 189-193.

Azzam, N.A. and Kuchn, D.P. (1978): The morphology of the musculus uvulae. Cleft palate J., 15: 337-348.

Basset, CA.L. (1972): Clinical implications of the cell function in bone grafting. Clinical Ortopedics, 87: 49-59.

Bellantone, M., Coleman, N. J., and Hench, L. L. (2000): Bacteriostatic action of a novel four-component bioactive glass. J. Biomed Mater Res. Sep 5: 51(3): 484-490.

References

Bernstein, L. (1991): Maxillofacial clefts. In Paparella Otolaryngology, 3rd ed., volume 3 (A), chapter 13, pp. 983-1994. Philadelphia : W.B. Saunders Company.

Binderman, I., Fin N. (1990): Bone substitutes-organic, inorganic, and polymeric, cell material interactions, CRC handbook of Bioactive Ceramics (eds. T. Yamamuro., L. Hench, and J. Wilson), CRC Press, Boca Raton., pp.45-51.

Bluestone, C.D. and Klein, J.O.(1995): Physiology, pathophysiology, and pathogenesis of otitis media in infants and children, 2nd ed., chapter 3, pp. 17-37. Philadelphia, W.B. Saunders Company.

Borchgrevink, H.H.C. (1986): Cleft palate repair, In plastic and reconstructive surgery, ch. 4, by Muir, I.F.k. The baillire Tindall Co., London, p. 64-91.

Brescia, N.J.(1971): Anatomy of lip and palate, dental and speech aspects, by Grabb, W.c.; Rosenstein, S.w. and Bzocks (eds.) Littlebrown Co., Boston, pp. 12-20.

Broader, H. and Stauss, P.R. (1989): Self cocept of early primary school children with visible or invisible defects. Cleft palate J. 26: 114.

Broude, D.J. and Walte, D.E. (1974): Secondary closure of alveolar defects. J. Oral Surgery, 37: 829-840.

References

Bumsted, R.N. (1986): Cleft lip and palate. In otolaryngology- head and neck surgery volume II, part III, (clinical entities), Section A (congenital disorders), ch. 63, by Schuller, D.E. (ed.). The Mosby, C.V. Co, Saint Louis, P.1129-1167.

Burdi, A.R. (1968): Distribution of midpalatine cysts: A reevaluation of human palatal closure mechanisms. J. oral Surgery, 26 : 41-45.

Burdi, A.R. and Silvery, R.G.(1969): Sexual differences in closure of the human palatal shelves. Cleft palate J.,6:1.

Bzoch, K.R. (1965): Articulation in cleft palate. Cleft Palate J., 2: 340-345.

Caddy, C.M. and Reid, CD. (1985): An atraumatic technique for harvesting cancellous bone for secondary alveolar bone grafting in cleft palate. British J. Plastic Surgery, 24: 197-204. (Q.A. Jafek, et al., 1981).

Clifford, E. (1969): Cleft palate and the person: psychological studies of its impact. Cleft palate J., 6: 221.

Converse, J.M. and Knize, Dm. (1977): Bone grafting in the cleft palate patients. In reconstructive plastic surgery, 2nd edition, vol. IV, ch.48, by Converse, J.M. (ed.). Saunders, W.B. Co., Philadelphia. P. 2205-2212.

References

Costello BJ, Ruiz RL(2004): cleft lip and palate :comprehensive treatment and primary repair. In Milaro M, Larsen PE, Waite PD,Ghali G(eds): Petreson's oral and maxilla facial surgery. Hamilton, Ontario,BC Decker, pp.839-858.

Croft, C.B., Shprintzen, R.J. and Daniller, A.(1978): The occult Submucous cleft palate and the muscular uvulae. Cleft Palate J. 15: 150-154.

Croft, C.B., Shprintzen, R.J. and Ruben(1981): Hypernasal speech following adenotomylectomy. Otolaryngology Head and Neck Surgery, 89:179.

Damien, J. C., Parson, J. R. (1991): Bone graft and bone graft substitutes: A review of current technology and applications J App Biomaterials 2: 187-208.

Deep, ME., Messer, L.B., Lehnert, M.W., Hebda, T,w. and Waite, D.E. (1982): Canine eruption into grafted bone in maxillary alveolar cleft defects. Cleft palate J., 19: 9-16.

Dickson, D.R. (1986): The normal velopharyngeal mechanism. J. Childhood Community, Disord. 10 (special issue): 5-16.

Dingman, R.O. and Grabb, W.C. (1962): A new mouth gag. Plastic Reconstr. Surg., 29: 208.

References

Doyle, W.J., Reilly, J.s., Jardini, I. And Rovnak, S. (1986): Effect of palatoplasty on the function of the Eustachian tube in children with cleft palate. Cleft palate J., 23:63.

Edwards, M.(1980): Speech and language disability. In advances in the management of cleft palate, pp. 83-96. Edinburgh: Churchill Livingstone.

El-Ghannam, A., Ducheyne, P., and Shapiro, I. (1995): The effect of serum protein adsorption on osteoplast adhesion to bioactive glass and hydroxyapatite ceramic. 21st Ann. Meet. Soc. Biomat. Proc., vol XVIII, 406.

Eskeland, G., Bergland, O., Borchgrevink, H., Semb, G. and Abyhlom, F.E. (1985): Management of the cleft alveolar arch. In recent advances in plastic surgery, No. 3, by Jackson, I.T. and Sornmerland, B.c. (eds.). Churchill livingstone co., Edinburgh, pp.14-32.

Evan B, Kartzel, B.A. Patrick Basile, M.D., Peter F. koltz, M.D., John ,A. Gritto, M.D(2009): Current surgical practices in cleft care: Plast Reconstr Surg. September. pp899-906.

Furlow, L.T. (1980): Double reversing Z-plasty for cleft palate. In alveolar and palatal deformities, volume 3, by Millard , D.R. (ed.). Little Brown, Boston, pp.519.

References

Girgis, I.H. (1966): the blood supply of the uvula and its surgical importance. J. laryngol. Otol., 130/4: 397-402.

Gan, J. C., Ducheyne, P., Verilovic, E., and Shapiro, I.M. (2000): Bioactive glass serves as a substrate for maintenance of phenotypes of nucleus pulposus cells of the intervertebral disc. J. Biomed. Mater. Res. Sep 15: 51(4): 596-604.

Gordon, A.S., Jean-Louis, F. and Morton, R.p.(1988): Late ear sequelae in cleft palate patients. Int. J. Peiatr. Otorhinolaryngol., 15:149.

Grant, H.R., Quiney, R.E., Mercer, D.M. and Lodge, S.(1988): Cleft palate and glue ear. Arch. Dis. Child., 63:176.

Hench, L.(1988): Bioactive ceramics. In: Ducheyne P., Lemmons J.E. (eds.). Bioceramics: Materials Characteristics versus in vivo behavior. Annals of the New York Academy of Sciences. 523:54-71.

Heppenstall, R.(1980): Bone grafting, fracture treatment and healing R. Heppenstall, W. P. Saunders, company, Philadelphia, pp 97-112.

Isshiki, N. and Koyama, H. (1980): Palatoplasty without elevation of mucoperiosteal flap on the cleft side. Ann. Plastic surgery, 4: 457.

Jackson, L.T., Scheker, L.R., Vandervord, J.G. and McLennan, J.G. (1981): Bone Marrow grafting in the secondary closure of alveolar-palatal defects in children. British J. Plastic Surgery, 34:422-425.

References

Jolleys, A. and Robertson, R.E. (1972): A study of the effects of early bone grafting in complete clefts of the lip and palate. Five years study. British J. Plastic Surgery, 25:229.

Kaufmann, E.A., Ducheyne, P., Randin, S., Bonnell, D.A., and Composto., R.(2000): Initial events at the bioactive glass surface in contact with protein-containing solutions. J. Biomed Mater Res. Dec. 15: 52(4): 825-830.

Kim, C. Y., Clark, A.E., and Hench, L.L. (1989): Early stages of calcium phosphate layer formation in bioglasses, J. Non Cryst Sol 113: 195-202.

Kinnunen, I., Aitasalo, K., Pollonen, M., and Varpula, M.(2000): Reconstruction of orbital floor fractures using bioactive glass. J. Maxillofac Surg Aug. 28(4): 229-234.

Krause, C.J., Tharp, R.F. and Moris, H.L. (1976): A comparative study of the results of the VonLangenbeck and V-Y. pushback palatoplasties. Cleft palate J., 13:11.

Laczka, M., Colewa-Kowalska, K., Laczka-Osyczka, A., Tworzydło, M., and Turyna, B. (2000): Geld-Derived materials of a CaO-P(2)O(5)-SiO(2) system modified by boron, sodium, magnesium, aluminium, and fluorine compounds. J. Biomed Mater Res. Dec 15: 52(4): 601-612.

La Rossa, D., Vaniver, K. and Randall, P. (1986): The effect of preoperative antibiotics on postoperative morbidity in patients with

References

cleft lip and palate. A retrospective study. Presented at the American cleft palate association meeting, New York.

Last, R.J. (1984): Anatomy regional and applied, 7th edition. The English Society and Churchill Livingstone, Edinburgh, and London, P. 407-424.

Lees, V.C. and Piggot, R.W. (1992): Early postoperative complications in primary cleft lip and palate surgery-how soon may we discharge patients from hospital? British J. of Plast. Surg. 45: 232-234.

Le Gros, R.(1988): Calcium phosphate materials in reconstructive dentistry: A review, Advances in Dental Res. 21:219-236.

Lindsay, W.K.(1974): Von Langenbech-simple closure palatoplasty. In symposium on management of cleft lip and palate and associated deformities, volume 8, ch. 25, by Georgiade, N.G. (ed.). The Mosby C.V. Co., Saint Louis, pp. 161-164.

Mason, R.M. and Warren, D.W. (1980): Adenoid involution and developing hypernasality in cleft palate. J. of Speech and Hearing disorders, 49:469.

McCarthy, I.J. (1980): Plastic Surgery, Vol. 4, pp.2697-746. Philadelphia: W.B. Saunders Co.

McMinn, R.M.H., Hutching, R.T. and Logan, B.M. (1981): Head and neck anatomy. Year Book publishers, Chicago.

References

McWilliams, B.J., Shelton, R.L. and Moris, H.L. (1984): Cleft palate speech. Philadelphia: B.C. Deckers.

Menyuk, R. (1980): Effect of persistent otitis media on language development. Ann. Otol. Rhinol. Laryngol., 89:257.

Moore, L.K. (1982): The developing human. Clinically oriented embryology. Philadelphia: WB Saunders.

Nakakita, M.D., Maeda, M.D., Ojimi, M.D. and Maekawa, M.D. (1991): The modified buccal Musclomucosal flap method for cleft palate surgery. Plast. And Reconstr. Surgery: 421-426.

Nylen, B., Korlof, B., Amander, C., Leanderson, R., Barr, B. and Nordin, KE. (1974): Primary, early bone grafting in complete clefts of the lip and palate. A follow up study of 53 cases. Scandinavian J. plastic surgery, 8:79.

Onishi, H, Kushitani, S., Yasukawa, E., Iwaki, H., Hench, L.L., Wilson, J., Tsuji, and Sugihara, T. (1997): Pariticulate Bioglass compared with hydroxyapatite bas a bone graft substitute. Clinical Orthopedics and related Res. 334: 316-325.

Paradise, J.L., Bluestone, C.D. and Felder, H.(1969): The universality of otitis media in 50 patients with cleft palate. Pediatrics, 44: 35-42.

Peltola, M, Suonpaa, J., Aitasalo, K., Maattanen, H., Andresson, O., and Laippala, P.(2000): Experimental follow-up model for clinical

References

frontal sinus obliteration with bioactive glass. *Acta Otolaryngol Suppl*, 543: 167-169.

Parven K Patel. Rev. article(2006): Craniofacial cleft palate repair. eMedicine June 14.

Randall, P. and Grabb, W. (1968): plastic surgery little brown Co. Boston, p.169.

Randall, P., Larossa, D.D., Solman, M. and Cohen, M. (1986): Experience with Furlow double reversing Z-plasty for cleft palate repair. *Plastic Reconstr. Surgery*, 77: 569-574.

Ricci, J.L., Blumenthal N.C., Spivak, J. M., and Alexander, H. (1992): Evaluation of a low calcium phosphate particulate implant material: physical chemical properties and vivo bone response, *J Oral Maxillofacial Surg.* , 969-978.

Richman, L.C and Eliason, M.(1984): Type of reading disabilities related to cleft type and neuropsychological patterns. *Cleft Palate J.*,21:1.

Ripamonti, U.(1990): Inductive bone matrix and porous hydroxyapatite composites in rodents and non-human primates, *CRC Handbook of Bioactive Ceramics* (Eds. T. Yamamuro, L. L. Hench, and J. Wilson), CRC Press, Boca Raton, 1990, pp. 245-253.

References

Robinson, P.J., Lodge, S., Jones, B.M., Walker, C.C. and Grant, H.R. (1992): The effect of palate repair on otitis media with effusion. *Plast. Reconstr. Surg.*, 89: 640-645.

Romanes, G.J. (1977): Cunningham's textbook of anatomy. 14th edition, Oxford University Press, London, p.293 and 401-403.

Rood, S.R. and Stool, S.E. (1981): Current concepts of aetiology, diagnosis and management cleft palate related otopathologic disease. *Otolaryngologic Clinics of north America*, 14: 865-884.

Ross, R.B. (1987): Treatment variables affecting facial growth in complete unilateral cleft lip and palate. *Cleft palate J.* 24 : 5-77.

Schepers, E.J.G., Ducheyne, P., Babier, L.(1993): Bioactive glass particles of narrow size range: a new material for the repair of bone defects, *Int. Dentistry*, 2,3:151-156.

Seibert, R.W. and Bumsted, R.M. (1993): Cleft lip and palate. In Cummings Otolaryngology-Head and neck Surgery, 3rd edition ., vol3, chapter 66, pp. 1128-1164. Saint Louis : Mosby year book.

Sell, D. (1987): Disorders of speech. In Scott-Brown's Otolaryngology, 5th edition , volume 6, chapter 11, pp. 136-158. London: Butterworths.

Sicher, n. and Du Brill, E.L.(1975): Oral anatomy, 6th edition. The Mosby C.V. Co, Saint Louis, pp.330-346.

References

Shapoff, C.A., Alexander, D.C. and Clark, A.E. (1997): Clinical use of a bioactive glass particulate in the treatment of human osseous defects. *Compendium*. 18: 352-363.

Simpson, R.K. and Austin, A.A. (1972): A cephalometric investigation of velar stretch. *Cleft palate J.*, 9: 341-351.

Smith, H.W.(1983): The atlas of cleft lip and cleft palate surgery, by Lee, K.J. (ed.) Grune and Stratton, New York.

Stanley, H.R., Clark, A.E., Hench L.L. and Berte, J.J. (1997): Using 45S5 Bioglass cones as endosseous ridge maintenance implants to prevent alveolar ridge resorption: A 5-year evaluation, *The international journal of oral and maxillofacial implants*, 12: 95-101.

Stark, R.B. (1977): Cleft palate. In *Reconstructive plastic surgery*, 2nd edition, vol. 4, ch. 45, by Converse, J.M. (ed.) Saunders W.B. Co., Philadelphia, pp.2090-2115.

Thompson, M.D. and Haerwood-Nash, D.(1972): The fate of the infractured hamuli. *Plastic Reconstructive Surgery* 50: 354.

Tisza, V.B. and Gumpertz, E. (1962): The parents reaction to the birth and early care of children with cleft palate. *Pediatrics*, 30:86.

References

Trasler, D.G., Fraser and F.C (1977): time position relationship. In Wilson JG, Fraser, EG (eds.). Handbook of teratology. Vol. (2). New York: Plenum 271.

Trier, W.C. (1986): Surgery for congenital cleft palate. Advances in plastic and reconstructive surgery, Vol. 2, by Habal, M.B. and Morain, W.D. (eds.). Year Book Medical Publisher Inc, Chicagoc , London, pp. 192-226.

Van Demark, D.R., Hardin, A.M. and Morris, L.H. (1988): Assessment of velopharyngeal incompetence. A long term process. Cleft palate J., 25(4): 362.

Vanarasdall, R.L and Coru, H.(1977): Soft tissue management of labially positioned unerupted teeth. American J. of Ortodontics, 72:53-64.

Von Langenbech, B. (1861): Die uranoplastik mittels abschloung des mokos-periostalen gaum enuberzuges. Arch. Klin. Chir., 2:205 (Q.A. Bumsted, 1986).

Waite, D.E. and Kersten, R.B. (1980): Residual alveolar and palatal clefts. In surgical correction of dentofacial deformities, Vol.5, ch. 7, Bell, W.H., Profit, W.R., and White, R.p. (eds.) Saunders W.B. Co., Philadelphia, pp. 1329-1367.

References

Walker, B.E. and Fraser, F.C. (1956): Closure of the secondary palate in three strain of mice. J. Embryol, Exp. Morphol. 4:176.

Wardill, W.E.M. (1937): Technique for cleft palate. British J. Surgery, 25:117. (Q.A. Smith, 1983).

Warwich, R. and Williams, P.I. (1973): Gray's anatomy 35th edition. Longman Group Ltd., pp.630 and 1206-1209.

Wilson, J., Clark, A.E., Hall, M.et al. (1993): Tissue response to bioglass endosseous ridge maintenance implants. J. Oral Implantol. (19):295.

Wilson, J., Pigott, G.H. and Hench L.L.(1981): Toxicology and biocompatibility of bioglasses. Journal of Biomedical Materials Res. 15: 805-817.

Xynos, I.D., Edgar, A.J. , Buttery, L.D., Hench,L.L., and Polak J., M.(2000 a): Ionic products of bioactive glass dissolution increase proliferation of human osteoblasts and induce insulin like growth factor II mRNA expression and protein synthesis. Biochem Biophys Res Commun. Sep. 24;276(2): 461-465.

References

Xynos, I.D., Buttery, L.D., L.L., and Polak J., M. (2000 b): Bioglass 45S5 stimulates osteoblast turnover and enhances bone formation in vitro: implications and application or bone tissue engineering. *Calcif Tissue Int* Oct., 67(4): 321-329.

Yules, R.B., Chase, R.A., Blocksma, R. and Long, R.B. (1971): Secondary techniques for correction of palatopharyngeal incompetence. In *cleft lip and palate*, by Grabb, W.C.: Rosenstien, S.W. and Bzoch, K.R. (eds.). Little Brown Co., Boston, P. 451-466.