

Reference

Amy J. Wagers and Irving L. Weissman (March 5, 2004):

Plasticity of Adult Stem Cells, *Cell*, Vol. 116, 639–648.

Avital I, Inderbitzin D, Aoki T et al. (2001): Isolation, characterization, and transplantation of bone marrow-derived hepatocyte stem cells. *Biochemical and Biophysical Research Communications*, 288, 156–164.

Axtell J. (1999):

The School Upon a Hill: Education and Society in Colonial New England. In: New Haven: Yale University Press.

Barbash I M, Chouraqui P, Baron J et al. (2003):

Systemic delivery of bone marrow-derived mesenchymal stem cells to the infarcted myocardium: Feasibility, cell migration, and body distribution. *Circulation*, 108, 863–868.

Bisgaard HC, Nacy P, Ton PR et al (1994):

Modulation of keratin 14 and a-fetoprotein during hepatic oval cell proliferation and liver regeneration. *J Cell Physiol*;159:475-484.

Blakolmer K, Jaskiewicz K, Dunsford HA et al (1995):

Hematopoietic stem cell markers are expressed by ductal plate and bile duct cells in developing human liver. *Hepatology*;21:1510–1516.

Bosch FX, Ribes J, Diaz M, et al (Nov 2004):

Primary liver cancer: worldwide incidence and trends. *Gastroenterology*. ;127(5 Suppl 1):S5-S16.

Brill S, Holst PA, Zvibel L, Fiorino A, Sigal SH, et al (1994):

Liver Biology and Pathobiology. 3rd ed. New York: Raven Press, :869 – 897.

Brockes JP, and Kumar A (2002):

Plasticity and reprogramming of differentiated cells in amphibian regeneration. *Nat. Rev. Mol. Cell Biol.* 3, 566–574.

Reference

Broelsch CE, Frilling A, Malago M (Apr 2004):

Hepatoma--resection or transplantation. *Surg Clin North Am*;84(2):495-511, x.

Brustle O, Choudary K, Karram K, et al. (1998):

Chimeric brains generated by intraventricular transplantation of human brain cells into embryonic rats. *Nat Biotech* 16: 1040–1044

Bruix J, Sherman M (Nov 2005): Management of hepatocellular carcinoma. *Hepatology*. ;42(5):1208-36.

Burkert J, Wright NA, Alison MR (2006):

Stem cells and cancer: an intimate relationship. *J Pathol*;209:287–297.

Busuttil RW, Farmer DG (Sep 1996):

The surgical treatment of primary hepatobiliary malignancy. *Liver Transpl Surg*. ;2(5 Suppl 1):114-30.

Cai J, Zhao Y, Liu Y, Ye F, Song Z, Qin H, Meng S, Chen Y,

Zhou R, Song X, Guo Y, Ding M, and Deng H. (2007):

Directed differentiation of human embryonic stem cells into functional hepatic cells. *Hepatology* 45, 1229-39.

Calabrese, C., et al., (2007):

“A perivascular niche for brain tumor stem cells,” *Cancer Cell.*, 11:69-82.

Campard D, Lysy PA, Najimi M, Sokal EM(2008):

Native umbilical cord matrix stem cells express hepatic markers and differentiate into hepatocyte-like cells. *Gastroenterology*; 134: 833-848.

Cantz T, Sharma AD, Jochheim-Richter A, et al (2004):

Reevaluation of bone marrow-derived cells as a source for hepatocyte regeneration. *Cell Transplant* 13:659–666.

Reference

Cho C H, Parashurama, N, Park E Y, Suganuma K, Nahmias Y, Park J, Tilles A W, Berthiaume F, and Yarmush M L. (2007):

Homogeneous differentiation of hepatocyte-like cells from embryonic stem cells: applications for the treatment of liver failure. *Faseb J*.

Christa L, Simon MT, Flinois JP, Gebhardt R, Brechot C, Lasserre C (1994):

Overexpression of glutamine synthetase in human primary liver cancer. *Gastroenterology*; 106:1312 -1320.

Cirillo L A, Lin F R, Cuesta I, Friedman D, Jarnik M, and Zaret K S. (2002):

Opening of compacted chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4. *Mol Cell* 9, 279-89

Clotman F, Jacquemin P, Plumb-Rudewiez N, Pierreux C E, Vander Smissen P, Dietz HC, Courtoy P J, Rousseau G G, and Lemaigre F P. (2005):

Control of liver cell fate decision by a gradient of TGF beta signaling modulated by Onecut transcription factors. *Genes Dev* 19, 1849-54.

Cormier JN, Thomas KT, Chari RS, et al (May 2006):

Management of hepatocellular carcinoma. *J Gastrointest Surg.* ;10(5):761-80.

Crawford JM (2005):

Liver and biliary tract. In: Kumar V, Abbas AK, Fausto N, eds. *Robbins and Cotran Pathologic Basis of Disease*. 7th ed. Philadelphia, Pa.: Elsevier Saunders, :877-938.

Curley SA, Izzo F, Ellis LM, et al (Sep 2000):

Radiofrequency ablation of hepatocellular cancer in 110 patients with cirrhosis. *Ann Surg.* ;232(3):381-91.

Reference

Dabeva MD, Shafritz DA (1993):

Activation, proliferation and differentiation of progenitor cells into hepatocytes in the D-galactosamine model of liver regeneration. *Am J Pathol*;143:1606-1620.

Dalakas E, Newsome PN, Harrison DJ, Plevris JN (2005):

Hematopoietic stem cell trafficking in liver injury. *FASEB J.* 19, 1225–1231.

Dalerba, P., et al (2005):

“Cancer Stem Cells: Models and Concepts,” *Annu. Rev. Med.*, 58:267-284.

D’Amour K A, Agulnick AD, Eliazer S, Kelly O G, Kroon E, and Baetge E E. (2005):

Efficient differentiation of human embryonic stem cells to definitive endoderm. *Nat Biotechnol* 23, 1534-41.

David A Axelrod, Dirk J van Leeuwen (Sep 18, 2008):

Hepatic carcinoma. *eMedicine specialities, General surgery, Abdomen.*

Dhawan A, Mitry RR, Hughes RD, Lehec S, Terry C, Bansal S, et al (2004):

Hepatocyte transplantation for inherited factor VII deficiency. *Transplantation*; 78:1812–1814.

Di Bisceglie AM, Carithers RL Jr, Gores GJ (Oct 1998):

Hepatocellular carcinoma. Hepatology. ;28(4):1161-5.

Díaz R, Kim JW, Hui JJ, Li Z, Swain GP, Fong KS, Csiszar K, Russo PA, Rand EB, Furth EE, Wells RG, (2008):

Evidence for the epithelial to mesenchymal transition in biliary atresia fibrosis. *Hum. Pathol.* 39, 102–115.

Doherty DG, Norris S, Madrigal-Estebas L, McEntee G, Traynor O, Hegarty JE, et al. (1999):

The human liver contains multiple populations of NK cells, T cells, and CD3-CD56-natural T-cells with distinct cytotoxic activities and Th1, Th2, and Th0 cytokine secretion patterns. *J Immunol*; 163: 2314-2321.

Reference

Don C. Rockey (February 2005):

Antifibrotic therapy in chronic liver disease ; Clinical Gastroenterology and Hepatology.

Duffield JS, Forbes SJ, Constandinou CM, et al (2005):

Selective depletion of macrophages reveals distinct, opposing roles during liver injury and repair. *J Clin Invest*; 115: 56-65.

Eiges R, Schuldiner M, Drukker M, Yanuka O, Itskovitz-Eldor J, and Benvenisty N (2001) :

Establishment of human embryonic stem cell-transduced clones carrying a marker of undifferentiated cells. *Curr. Biol.* 11, 514–518.

Eiseman, E. (2000):

Quick response: use of fetal human tissue in federally funded research. In: *Ethical issues in Human Stem Cell Research, Vol. II. Commissioned Papers. National Bioethics Advisory Commission, US Government Printing Office: Rockville, MD, pp C1 – C7.*

El-Serag HB, Mason AC (Nov 27 2000):

Risk factors for the rising rates of primary liver cancer in the United States. *Arch Intern Med.* ;160(21):3227-30.

Emilio Bajetta, Giuseppe Procopio, Andrea Colombo, Valentina Guadalupi, Elena Verzoni, Filippo Pietrantonio et al (2009):

Sorafenib in Hepatocellular Carcinoma. *Clinical Medicine: Therapeutics*:1 277–287.

Fabrikant JI (1968):

The kinetics of cellular proliferation in regenerating liver. *J Cell Biol*;36:551–565.

Fahrner J, Labruyere WT, Gaunitz C, Moorman AF et al (1993):

Identification and functional characterization of regulatory elements of the glutamine synthetase gene from rat liver. *Eur J Biochem*; 213:1067 – 1073.

Reference

Farley MA. (2000):

Roman Catholic views on research o research involving human embryonic stem cells. In: Ethical Issues in Human Stem Cell Research. Vol. 111. Religious Perspectives. National Bioethics Advisory Commission, US Government Printing Office. Rockville, MD, pp D1-D5.

Fausto N (2000):

Liver regeneration. J Hepatol. 32:19-31.

Fausto N. (2004):

Liver regeneration and repair: hepatocytes, progenitor cells, and stem cells. Hepatology;39:1477–1487

Fiegel HC, Gluer S, Roth B, Rischewski J, von Schweinitz D, Ure B, et al., (2004):

Stem-like cells in human hepatoblastoma. J Histochem Cytochem;52:1495–1501.

Fox IJ, Chowdhury JR, Kaufman SS, Goertzen TC, Chowdhury NR, Warkentin PI, et al (1998):

Treatment of the Crigler– Najjar syndrome type I with hepatocyte transplantation. N Engl J Med;338:1422–1426.

Friedman S, Schiano T (2004):

Cirrhosis and its sequelae. In: Goldman L, Ausiello D, eds. Cecil Textbook of Medicine. 22nd ed. Philadelphia, Pa.: Saunders:936-44.

Fujii H, Hirose T, Oe S, et al. (2002):

Contribution of bone marrow cells to liver regeneration after partial hepatectomy in mice. Journal of Hepatology, 36, 653–659.

Gardener R., Lanza R., Gearhart J., Hogan B., Melton D. and Pedersen R. (2006):

Present presepective and future challenges: In Essehtials of stem cell biology, 1st ed(Ed.by)Gardener R., Lanza R., Lanza R.,Gearhart J., Hogan B., Melton . and Pedersen R., Elsevier academic press. PP:1-10.

Reference

Jang YY, Collector MI, Baylin SB, Diehl AM, Sharkis SJ (2004):

Hematopoietic stem cells convert into liver cells within days without fusion. *Nat Cell Biol*; 6: 532-9.

Jiang Y, Jahagirdar BN, Reinhardt RL, et al. (2002):

Pluripotency of mesenchymal stem cells derived from adult bone marrow. *Nature* 418: 41–49.

Jonas S, Herrmann M, Rayes N, et al (Nov-Dec 2001):

Survival after liver transplantation for hepatocellular carcinoma in cirrhosis according to the underlying liver disease. *Transplant Proc.* ;33(7-8):3444-5.

Kakinuma S, Tanaka Y, Chinzei R, Watanabe M, Shimizu-Saito K, Hara Y, et al (2003):

Human umbilical cord blood as a source of transplantable hepatic progenitor cells. *Stem Cells*; 21: 217-27.

Keith E Stuart, Zsofia K Stadler (Jan 12, 2009):

hepatic carcinoma, primary . eMedicine specialities, oncology, carcinomas of the gastrointestinal tract.

Lagasse E, Connors H, Al-Dhalimy M, Reitsma M, Dohse M, Osborne L, Wang X, et al (2000):

Purified hematopoietic stem cells can differentiate into hepatocytes in vivo. *Nat Med*; 6: 1229-34.

Lake JR (2001):

Evaluation and management of prospective liver transplant candidates. In: Norman DJ, Turka LA, eds. *Primer on Transplantation*. 2nd ed. Mt. Laurel, N.J.: American Society of Transplantation:523-8.

Lavon N, Yanuka O and Benvenisty N (2004):

Differentiation and isolation of hepatic-like cells from human embryonic stem cells. *Differentiation* 72, 230-8.

Reference

Lee KD, Kuo TK, Whang-Peng J, Chung YF, Lin CT, Chou SH, Chen JR, Chen YP, Lee OK (2004):

In vitro hepatic differentiation of human mesenchymal stem cells. *Hepatology*; 40: 1275-1284

Lee OK, Kuo TK, Chen WM, Lee KD, Hsieh SL, Chen TH (2004):

Isolation of multipotent mesenchymal stem cells from umbilical cord blood. *Blood*; 103: 1669-75.

Lemaigre F, Zaret KS (2004):

Liver development update: New embryo models, cell lineage control, and morphogenesis. *Curr Opin Genet Dev*;14:582–590.

Leung TW, Tang AM, Zee B, et al (Jan 15 2002):

Factors predicting response and survival in 149 patients with unresectable hepatocellular carcinoma treated by combination cisplatin, interferon-alpha, doxorubicin and 5-fluorouracil chemotherapy. *Cancer*;94(2):421-7.

Liu ZC, & Chang, TM (2006):

Transdifferentiation of bioencapsulated bone marrow cells into hepatocyte-like cells in the 90% hepatectomized rat model. *Liver Transplantation*, 12, 566–572.

Llovet JM, Fuster J, Bruix J (Feb 2004):

The Barcelona approach: diagnosis, staging, and treatment of hepatocellular carcinoma. *Liver Transpl.* ;10(2 Suppl 1):S115-20.

Lo CM, Ngan H, Tso WK, et al (May 2002):

Randomized controlled trial of transarterial lipiodol chemoembolization for unresectable hepatocellular carcinoma. *Hepatology*.;35(5):1164-71.

Reference

Lucey MR, Brown KA, Everson GT, Fung JJ, Gish R, Keefe EB, et al (1998):

Minimal criteria for placement of adults on the liver transplant waiting list: a report of a national conference organized by the American Society of Transplant Physicians and the American Association for the Study of Liver Diseases. *Transplantation*;66:956-62

Lyra AC, Soares MB, da Silva LF, et al., (2007):

Feasibility and safety of autologous bone marrow mononuclear cell transplantation in patients with advanced chronic liver disease. World J Gastroenterol; 13: 1067-73.

Macdonald JM, Xu ASL, Hiroshi K, LeCluyse E, Hamilton G, Liu H, Rong YW, Moss N, Lodestro C, Luntz T, Wolfe SP, Reid L. (2002):

In: Lanza WL, Langer R, Vacanti J, eds. Methods of Tissue Engineering. San Diego: Academic Press;151 – 201.

Marrero JA, Fontana RJ, Barrat A, et al (Apr 2005):

Prognosis of hepatocellular carcinoma: comparison of 7 staging systems in an American cohort. Hepatology.;41(4):707-16.

Mathurin P, Raynard B, Dharancy S, et al (May 15 2003):

Meta-analysis: evaluation of adjuvant therapy after curative liver resection for hepatocellular carcinoma. Aliment Pharmacol Ther. ;17(10):1247-61.

Maurizio Muraca, Ginevra Galbiati, Maria Teresa Vilei, Aline Sueli Coelho Fabricio, Maddalena Caruso (2006):

The future of stem cells in liver diseases ; Annals of Hepatology; 5(2): April-June: 68-76

Michael Oertel, Anuradha Menthena, Yuan-Qing Chen, David A. Shafritz (October 2006):

Properties of Cryopreserved Fetal Liver Stem/Progenitor Cells That Exhibit Long-Term Repopulation of the Normal Rat Liver. Stem Cells Vol. 24 No. 10, pp. 2244 -2251

Michalopoulos GK, Khan Z (2005):

Liver regeneration, growth factors, and amphiregulin. Gastroenterology;128:503-506.

Mulholland, Michael W.; Lillemoe, Keith D.; Doherty, Gerard M.; Maier, Ronald V.; Upchurch, Gilbert R (2006):

Greenfield's Surgery: SCIENTIFIC PRINCIPLES AND PRACTICE, 4th Edition. Lippincott Williams & Wilkins

Newsome PN, Johannessen I, Boyle S, Dalakas E, McAulay KA, Samuel K, Rae F, et al (2003):

Human cord blood-derived cells can differentiate into hepatocytes in the mouse liver with no evidence of cellular fusion. Gastroenterology; 124: 1891-900.

Nierhoff D, Ogawa A, Oertel M, Chen YQ, Shafritz DA (2005):

Purification and characterization of mouse fetal liver epithelial cells with high in vivo repopulation capacity. Hepatology;42:130-139.

Nomoto M, Ychikosi Y, Kajikazawa N et al (1992):

Appearance of hepatocyte like cells in the interlobular bile ducts of human liver in various liver disease states. Hepatology;16:1199-1205.

Odorico JS, Kaufman DS and Thomson JA (2001):

Multilineage Differentiation from Human Embryonic Stem Cell Lines. Stem Cells. 19, 193-204

Okumoto K, Saito T, Hattori E et al. (2005): Expression of Notch signalling markers in bone marrow cells that differentiate into a liver cell lineage in a rat transplant model. Hepatology Research, 31, 7-12.

Reference

Ong SY, Dai H, Leong KW (2006):

Hepatic differentiation potential of commercially available human mesenchymal stem cells. Tissue Eng; 12: 3477-3485.

Pan XN, Shen JK, Zhuang YP, Chen XL, Li YX, Chen LJ, Yang H (Jul 2008):

Autologous bone marrow stem cell transplantation for treatment terminal liver diseases. Haematology; 2008 Jul;28(7):1207-9

Parashurama N, Nahmias Y, Cho CH, van Poll D, Tilles AW, Berthiaume F and Yarmush ML (2007) :

Activin Alters the Kinetics of Endoderm Induction in Embryonic Stem Cells Cultured on Collagen Gels. Stem Cells

Pawlik TM, Delman KA, Vauthey JN, et al (Sep 2005):

Tumor size predicts vascular invasion and histologic grade: Implications for selection of surgical treatment for hepatocellular carcinoma. Liver Transpl.;11(9):1086-92.

Petersen BE, Bowen WC, Patrene KD, Mars WM, Sullivan AK, Murase N, Boggs SS, Greenberger JS, Goff JP (1999):

Bone marrow as a potential source of hepatic oval cells. Science; 284:1168-70.

Poon RT, Fan ST, Lo CM, et al (Jul 2001):

Improving survival results after resection of hepatocellular carcinoma: a prospective study of 377 patients over 10 years. Ann Surg;234(1):63-70.

Preston, MR Alison, SJ Forbes, NC Direkze, R Poulson, and NA Wright (2003 April):

The new stem cell biology: something for everyone; Molecular Pathology, Journal of clinical pathology; 56(2): 86-96.

Radisky DC, (2005):

Epithelial-mesenchymal transition. J. Cell. Sci. 118, 4325- 4326.

Rambhatla L, Chiu CP, Kundu P, Peng Y and Carpenter MK (2003):
Generation of hepatocyte-like cells from human embryonic stem cells.
Cell Transplant 12, 1-11.

Ratajczak MZ, Machalinski B, Wojakowski W, Ratajczak J, Kucia M (2007).
"A hypothesis for an embryonic origin of pluripotent Oct-4(+) stem cells in adult bone marrow and other tissues" . Leukemia 21 (5): 860-7

Reubinoff BE, Pera MF, Fong CY, Trounson A and Bongso A (2000):
Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. Nat. Biotechnol. 18, 399-404

Riehle KJ, Campbell JS, McMahan RS, Johnson MM, Beyer RP, Bammler TK, et al (2008):
Regulation of liver regeneration and hepatocarcinogenesis by suppressor of cytokine signaling 3. J Exp Med;205:91-103.

Robertson H, Kirby JA, Yip WW, Jo Riehle KJ, Campbell JS, McMahan RS, Johnson MM, Beyer RP, Bammler TK, et al (2008):
Regulation of liver regeneration and hepatocarcinogenesis by suppressor of cytokine signaling 3. J Exp Med;205:91-103

Robey PG (2000):
Stem cells near the century mark. J. Clin. Invest. 105, 1489-1491.

Roskams TA, Theise ND, Balabaud C, Bhagat G, Bhathal PS, Bioulac-Sage P, et al (2004):
Nomenclature of the finer branches of the biliary tree: canals, ductules, and ductular reactions in human livers.
Hepatology;39:1739-1745.

Rossi JD., NR. Hogan , BL. Zaret, KS. (2001):
Distinct mesodermal signals, including BMPs from the septum transversum mesenchyme, are require in combination for hepatogenesis from the endoderm. Genes Dev 15, 1998-2001.

Rougier P, Mitry E, Barbare JC, et al (Apr 2007):

Reference

Hepatocellular carcinoma (HCC): an update. Semin Oncol;34(2 Suppl 1):S12-20.

Rubin H (2002):

The disparity between human cell senescence in vitro and lifelong replication in vivo. Nat Biotechnol 20:675-681

Sandhu JS, Petkov PM, Dabeva MD and Shafritz DA (2001):

Stem cell properties and repopulation of the rat liver by fetal liver epithelial progenitor cells. Am. J. Pathol. 159:1323-1334.

Sarraf C, Lalani E-N, Golding M et al (1994):

Cell behavior in the actelyaminofluorene-treated regenerating rat liver. Light and electron microscopic observations. Am J Pathol;145:1114-1126.

Sato Y, Araki H, Kato J et al. (2005):

Human mesenchymal stem cells xenografted directly to rat liver are differentiated into human hepatocytes without fusion. Blood, 106, 756-763.

Schaffner F, Popper H (1963):

Capillarization of hepatic sinusoids in man. Gastroenterology;44:239-242.

Sekhon SS, Tan X, Micsenyi A et al (2004):

Fibroblast growth factor enriches the embryonic liver cultures for hepatic progenitors. Am J Pathol;164:2229-2240.

Sei Kakinuma, Hiromitsu Nakauchi and Mamoru Watanabe (2009):

Hepatic stem/progenitor cells and stem-cell transplantation for the treatment of liver disease; J Gastroenterol; 44:167-172

Sell S (1990):

Is there a liver stem cell? Cancer Res. 50,3811-3815

Sell S (1993):

Reference

The role of determined stem cells in the development of hepatocellular carcinoma. Int J Develop Biol;37:189-201

Reference

Shamblott, M.J., Axelman, J., Wang, S., Bugg, E.M., Littlefield, J.W., Donovan, P.J., Blumenthal, P.D., Huggins, G.R., and Gearhart, J.D. (1998):

Derivation of pluripotent stem cells from cultured human primordial germ cells. Proc. Natl. Acad. Sci. U. S. A. 95, 13726–13731.

Sharma AD, Cantz T, Richter R, Eckert K, Henschler R, Wilkens AJR, Arseniev L (2005):

Human cord blood stem cells generate human cytokeratin 18-negative hepatocyte-like cells in injured mouse liver. Am J Pathol; 167: 555-64.

Sherwood RI, Jitianu C, Cleaver O, Shaywitz DA, Lamenzo JO, Chen AE, Golub TR and Melton DA (2007):

Prospective isolation and global gene expression analysis of definitive and visceral endoderm. Dev Biol 304, 541-55.

Sjogren A, Hardarson T, Andersson K, et al (2004):

Human blastocysts for the development of embryonic stem cells. Reprod Biomed Online 9:326–329

Snorri S. Thorgeirsson (1996):

Hepatic stem cells in liver regeneration; FASEBJ. 10,1249-1256

Soto-Gutierrez A, Navarro-Alvarez N, Zhao D, Rivas -Carrillo JD, Lebkowski J, Tanaka N, Fox I J and Kobayashi N (2007):

Differentiation of mouse embryonic stem cells to hepatocyte-like cells by co-culture with human liver nonparenchymal cell lines. Nat Protoc 2, 347-56.

Stadtfeld M, Nagaya M, Utikal J, Weir G, Hochedlinger K (2008 Sep 25):
Induced Pluripotent Stem Cells Generated without Viral Integration. Science.

Stefania Lorenzini, Stefano Gitto, Elena Grandini, Pietro Andreone, Mauro Bernardi (2008 August 7):

Reference

Emergence of intraembryonic hematopoietic precursors in the pre-liver human embryo. Development. 126, 793–803.

Tee LBG, Kirilak Y, Huang W-H et al (1996):

Dual phenotypic expression of hepatocytes and bile ductular markers in developing and preneoplastic rat liver.17:251-259.

Terai S, Ishikawa T, Omori K et al (2006):

Improved liver function in liver cirrhosis patients after autologous bone marrow cell infusion therapy. STEM CELLS;24:2292–2298.

Theise ND, Saxena R, Portmann BC, Thung SN, Yee H, Chiriboga L, et al (1999):

The canals of Hering and hepatic stem cells in humans. Hepatology;30:1425–1433.

Thomas P. Zwaka (20 January 2006):

Use of genetically modified stem cells in experimental gene therapy; Science: Vol. 311. no. 5759, p. 335.

Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS and Jones JM (1998):

Embryonic stem cell lines derived from human blastocysts. Science. 282, 1145–1147.

Vassilopoulos G, Wang PR, Russell DW (2003):

Transplanted bone marrow regenerates liver by cell fusion. Nature; 422: 901-4.

Vogels BA, Maas MA, Bosma A et al. (1996):

Significant improvement of survival by intrasplenic hepatocyte transplantation in totally hepatectomized rats. Cell Transplant, 5, 369–378.

Wang X, Foster M, Al-Dhalimy M, Lagasse E, Finegold M, Grompe M (2003):

Reference

The origin and liver repopulating capacity of murine oval cells. Proc Natl Acad Sci USA;100(suppl 1):11881-11888.

Weimann JM, Johansson CB, Trejo A and Blau HM (2003):

Stable reprogrammed heterokaryons form spontaneously in Purkinje neurons after bone marrow transplant. Nat. Cell Biol. 5, 959-966.

Weissman IL. (2000):

Translating stem and progenitor cell biology to the clinic: barriers and opportunities. Science 287:1442-1446

Wilhelm A, Leister I, Sabandal P et al. (2004):

Acute impairment of hepatic microcirculation and recruitment of nonparenchymal cells by intrasplenic hepatocyte transplantation. Journal of Pediatric Surgery, 39, 1214-1219.

Willenbring H, Bailey AS, Foster M, Akkari Y, Dorrell C, Olson S, Finegold M, Fleming WH, Grompe M (2004):

Myelomonocytic cells are sufficient for therapeutic cell fusion in liver. Nat Med; 10: 744-748

Wilmut I and Paterson L (2003):

Somatic cell nuclear transfer. Oncol Res; 13:303-307.

Wobus AM and Boheler KR (2005):

Embryonic stem cells: prospects for developmental biology and cell therapy. Physiol Rev 85, 635-78.

Xu A, Luntz T, Macdonald J, Kubota H, et al (2000):

In: Lanza R, Langer R, Vacanti J, eds. Principles of Tissue Engineering. 2nd ed. San Diego: Academic Press.

Yamada Y, Nishimoto E, Mitsuya H, Yonemura Y (2006):

In vitro transdifferentiation of adult bone marrow Sca-1+ cKit cells cocultured with fetal liver cells into hepatic-like cells without fusion. Exp Hematol; 34: 97-106

Reference

Zeisberg M, Yang C, Martino M, Duncan MB, Rieder F, Tanjore H, Kalluri R (2007):

Fibroblasts derive from hepatocytes in liver fibrosis via epithelial to mesenchymal transition. J. Biol. Chem. 282, 23337–23347

Zhu AX, Blaszkowsky LS, Ryan DP, et al (April 2006):

Phase II study of gemcitabine and oxaliplatin in combination with bevacizumab in patients with advanced hepatocellular carcinoma. J Clin Oncol. ;24(12):1898-903.