

References

- A**lagarsamy S, Chhabra M, Gudavalli M , et al,(2005).Comparison of clinical criteria with echocardiographic findings in diagnosing PDA in preterm infants. *J Perinat Med*;33:151-'J'64.
- Allan LD and Sharland GK. (2001).** "The echocardiographic diagnosis of totally anomalous pulmonary venous connection in the fetus"
- Anderson RH and Weinberg.(2005)** The clinical anatomy of tetralogy of Fallot. *Cardiol Young*.
- Andrews RE, Tulloh RMR,(2004):** Interventional cardiac catheterization in congenital heart disease. *Arch Dis Child* 89:7168-1773.
- B**akr AF and Habib HS. (2005).Combining pulse oximetry and clinical examination in screening for congenital heart disease . *Pediatr Cardiol*; 26 :832 – 835.
- Bhatikar SR, DeGross C and Mahajan RL.(2005).** A classifier based on the artificial neural network approach for cardiologic auscultation in pediatrics. *Artif Intell Med*;33:251,-260.
- Biancaneello T.(2005)** .Innocent murmurs. *Circulation*:111:e20-+22.
- Bjørnstad P. (2006).** "Is interventional closure the current treatment of choice for selected patients with deficient atrial septation?"
- Blecha and Matthew J. (2005).** "General Surgery ABSITE and Board Review (Pearls of Wisdom)"
- Brion LP and Soll RF. (2008).** Diuretics for respiratory distress syndrome in preterm infants. *Cochrane Database Syst Rev*.
-

Brown KL , Ridout DA and Hoskote A.(2006). Delayed diagnosis of congenital heart disease worsens preoperative conditions and outcome of surgery in neonates . ; 92 : 1298 – 1302 *oung* **16** (1): 3–10. doi:10.1017/S1047951105002027. PMID 16454871

Burch TM, Mizuguchi KA, Wesley MC, Swanson TM and Dinardo JA.(2008). Echocardiographic features of pulmonary atresia with intact ventricular septum. *Anesth Analg.*

Chang RK, Gurvitz M and Rodriguez S. (2008) Missed diagnosis of critical congenital heart disease. *Arch Pediatr Adolesc Med* 162:969–974

-Comber JT, Lopez BL(1996): Evaluation of pulse oximetry in sickle cell anemia patients presenting to the emergency department in acute

Cotran, R., V. Kumar, and N. Fausto (2005). Robbins Pathologic Basis of Disease, 7th Ed.. W.B. Saunders

Dasgupta C, Martinez AM, Zuppan CW, Shah MM, Bailey LL, Fletcher WH(2001). "Identification of connexin43 (alpha1) gap junction gene mutations in patients with hypoplastic left heart syndrome by denaturing gradient gel electrophoresis (DGGE)"

de Kort E, Thijssen JM, Daniels O, et al(2006). Improvement of heart function after balloon dilation of congenital valvar aortic stenosis: a pilot study with ultrasound tissue Doppler and strain rate imaging. *Ultrasound Med Biol.*

de Wahl Granelli A , Wennergren M , Sandberg K et al ,(2009). Impact of pulseoximetry screening on the detection of duct dependent congenital heart disease: a Swedish prospective screening study in 39 821 newborns. *BMJ*; 338 : a3037(7) .

Edelist **G(1995):** Acrylic nails and pulse oximetry [letter]. *Anesth Analg*, 81:882-891.

Frommelt **MA and Frommelt PC(1999):.** Advances in echocardiographic diagnostic modalities for the pediatricran. *Pediatr Clin North Am*;46:427-439.

Fu YC, Bass J, Amin Z, et al. (2006). Transcatheter closure of perimembranous ventricular septal defects using the new Amplatzer membranous VSD occluder: results of the U.S. phase I trial. *J Am Coll Cardiol.* ;47(2):319-25

Galindo **A, Nieto O, Villagra S, Graneras A, Herraiz I, Mendoza A(2009).** Hypoplastic left heart syndrome diagnosed in fetal life: associated findings, pregnancy outcome and results of palliative surgery. *Ultrasound Obstet Gynecol.*

Gatzoulis MA, Webb GD, Daubeney PE. (2005). Diagnosis and Management of Adult Congenital Heart Disease. Churchill Livingstone, Philadelphia.

Giardini A, Moore P, Brook M, et al (2005). Effect of transcatheter atrial septal defect closure in children on left ventricular diastolic function. *Am J Cardiol*;95:125-125.

Gill HK, Splitt M, Sharland G, et al(2003): Patterns of recurrence of congenital heart

Griebsch I , Knowles RL , Brown J et al .(2007). Comparing the clinical and economic effects of clinical examination, pulse oximetry, and echocardiography in newborn screening for congenital heart defects: a probabilistic cost-effectiveness model and value of information analysis .*Int J Technol Assess Health Care* ; 23 : 192 – 204

Hanslik A, Pospisil U, Salzer-Muhar U, Greber-Platzer S, Male C. (2006). Predictors of spontaneous closure of isolated secundum atrial septal defect in children: a longitudinal study. *Pediatrics*;118(4):1560-5.

Hirayama-Yamada K, Kamisago M, Akimoto K, et al(2005). Phenotypes with GATA4 or NKX2.5 mutations in familial atrial septal defect. *Am J Med Genet A.* ;135(1):47-52.

Hoffman JI and Kaplan S.(2002). The incidence of congenital heart disease. *J Am Coll Cardiol.* Jun 19;39(12):1890-900. PMID 12084585

Hohn AR, Dwyer KM and Dwyer JH (2004): Blood pressure in youth from four ethnic groups: The Pasadena Prevention Project.,*f Pediatr* 199 4;125 :368-37 3.Pelech AN: The physiology of cardiac auscultation. *Pediatr Clin North Am*;51:1515-1535.

Holzer R, Balzer D, Cao QL, et a)2004): Device closure of muscular ventricular septal defects using the Amplatzer muscular ventricular septal defect occluder: Immediate and mid-term results of a U.S. registry. *J Am Coll Cardiol*;43:1257-1 263.

Horer J, Schreiber C, Dworak E, et al.,(2007).Long-term results after the Rastelli repair for transposition of the great arteries. *Ann Thorac Surg.*

Jonas RA. (2004) :Atrial septal defect. In: *Comprehensive Surgical Management of Congenital Heart Disease.* :225-41.

Jubran A.(1998): Pulse oximetry. In: Tobin MJ (ed). *Principles and Practice of Intensive Care Monitoring.* New York: McGraw Hill, Inc;, :261-287

Jubran A, Tobin MJ(1996): Monitoring during mechanical ventilation. *Clin Chest Med*, 17:453-473.

Khoo NS , van Essen P , Richardson M et al.,(2008). Effectiveness of prenatal diagnosis of congenital heart defects in south Australia: A population analysis 1999-2003 . *Australian and New Zealand Journal of Obstetrics and Gynecology* ; 48 : 559 – 563

Khoshnood B , De Vigan C , Vodovar V et al.,(2005) . Trends in prenatal diagnosis, pregnancy termination and prenatal mortality of newborns with congenital heart disease in France, 1983 – 2000: a population based evaluation. *Pediatrics* ; 115 : 95 – 10

Knauth AL, Lock JE, Perry SB, et al,(2004): Transcatheter device closure of congenital and postoperative residual ventricular septal defects. *Circulation*;110:501-507.

Konduri GG.(2004): New approaches for persistent pulmonary hypertension of newborn. *Clin Perinatol*;31:591-611.

Koppel RI , Druschel CM , Carter T et al.,(2003). Effectiveness of pulse oximetry screening for congenital heart disease in asymptomatic newborns .Pediatrics; 111 : 451-455]

Kreutzer J(2001.): Transcatheter intervention in the neonate with congenital heart disease C. /iz Perinatol:28:1.37-517 .

Langer A and John J(2009) . Neugeborenen screening im Spannungsfeld der Gesundheits ö konomie . Monatsschr Kinderheilk ; 157 : 1230 – 1236

Lilly LS (editor) (2003). Pathophysiology of Heart Disease (3rd ed.). Lippincott Williams & Wilkins.

Mahle WT , Newburger JW , Matherne GP et al.(2009) . Role of pulse oximetry in examining newborns for congenital heart disease: a scientific statement from the American Heart Association and American Academy of Pediatrics . Circulation ; 120 : 447 – 458(15] .

Marino BS, Bird GL, Wernovsky G, (2008): Diagnosis and management of the newborn with suspected congenital heart disease. Clin Perinatal; 28: 91- 136

Maslen CL.(2004). Molecular genetics of atrioventricular septal defects. Curr Opin Cardiol 2004 ; 1,9:205 – 270 disease: An analysis of 6,640 consecutive pregnancies evaluated by detailed fetal echocardiography., J Am Coll Cardiol;42:923-929

Masura J, Gavora P, Podnar T.(2005). Long-term outcome of transcatheter secundum-type atrial septal defect closure using Amplatzer septal occluders/. J Am Coll Cardiol;45:505-507

McMillen IC and Robinson JS(2005): Developmental origins of the metabolic syndrome: Prediction, plasticity and programming. *Physiol Rev*; 85: 525-77-433.

Meberg A , Brüggemann-Pieper S , Due R Jr et al.(2008) . First day of life pulse oximetry screening to detect congenital heart defects . *J Pediatr* ; 152 : 761 – 765

Mellander M , Sunnegårdh J .(2008). Failure to diagnose critical heart malformations in newborns before discharge — an increasing problem? *Acta Paediatr* 2006 ; 95 : 407 – 413. Wren C, Reinhardt Z, Khawaja K. Twenty-year trends in diagnosis of life-threatening neonatal cardiovascular malformations. *Arch Dis Child Fetal Neonatal Ed*; 93: F33 – F35 [7] .

Mower WR, Sachs C, Nicklin EL, et al.,1995): Effect of routine emergency department triage pulse oximetry screening on medical management. *Chest*, 108:1297-1302.

Nienaber CA, Rehders T C and Fratz S(1999): Detection and assessment of congenital heart disease with magnetic resonance techniques *J Cardiovasc Magn Reson*; 1,; 1,69-784

Nourusis MJ (1997): Statistical package of social science (SPSS) for windows 8 advanced statistics Chicago, IL: SPSS.

Ohlsson A, Walia R, Shah S. (2008). Ibuprofen for the treatment of patent ductus arteriosus in preterm and/or low birth weight infants. *Cochrane Database Syst Rev*.

Payne RM, Johnson MC, Grant JW, Strauss AW (1995). Toward a molecular understanding of congenital heart disease. *Circulation* 91:494–504

Porter CJ, Feldt RH, Edwards WD, et al. (2001). Atrial septal defects. In: *Heart Disease in Infants, Children, and Adolescents*. Vol 1. 6th ed. :603-17.

Quiros-Lopez R, Garcia-Alegria J (2007). "A medical mystery -- high blood pressure". *N Engl J Med* 356 (25): 2630

Rajdev S, Nanda NC, Patel V, et al.(2006). Live/real-time three-dimensional transthoracic echocardiographic assessment of combined valvar and supra-valvar aortic stenosis. *Am J Geriatr Cardiol*.

-Rao PS.(2009). Diagnosis and management of cyanotic congenital heart disease: part I. *Indian J Pediatr*.

Reamon-Buettner SM, Hecker H, Spanel-Borowski, et al. (2004). Novel NKX2-5 mutations in diseased heart tissues of patients with cardiac malformations. *Am J Pathol*; '16, 4 :21,1 7- 2125.

Reich JD , Miller S , Brogdon B et al ., (2003) The use of pulse oximetry to detect congenital heart disease . *J Pediatr*; 142 : 268 – 272 19] . 56-Richmond S, Reay G and Abu Harb M (2002) Routine pulse oximetry in the asymptomatic newborn. *Arch Dis Child* 87:F83–F88

Reich DL, Timcenko A, Bodian CA, et al., (1996): Predictors of pulse oximetry data failure. *Anesthesiology*, 84:859-864

Richmond S and Wren C . (2001) .Early diagnosis of congenital heart disease . *Semin Neonatal* ; 6 : 27 – 35] .

-Riede FT , D ä hnert I , Woerner C et al.(2009) . Verkleinerung der diagnostischen L ü cke bei kritischen angeborenen Herzfehlern durch Pulsoxymetriescreening. *Monatsschr Kinderheilk* ; 157 : 896 – 902 [22] .

Roguin N, Du ZD, Barak M, Nasser N, Hershkowitz S and Milgram E.(1995). High prevalence of muscular ventricular septal defect in neonates. *J Am Coll Cardiol* November 15;26(6):1545-8

Rosati E , Chitano G , Dipaola L et al .(2005). Indications and limitations for a neonatal pulse oximetry screening of critical congenital heart disease . *J Perinat Med*; 33: 455 – 457

Rudolph AM.(2001): *Congenital Diseases of the Heart: Clinical-Physiological Considerations*, 2nd ed. New York, Futura,.

Sahn DJ and Vick GrJT III.(2001): Review of new techniques in echocardiography and magnetic resonance imaging as applied to patients with congenital heart disease *Heart* ; 86(Suppl2):II41-II53.

Saito S, Fukura H, Shimada H, Fujita T(1995): Prolonged interference of blue dye "patent blue" with pulse oximetry readings. *Acta Anaesthesiol Scand* , 89:268-269.

Saxena A (2003) Issues in newborn screening. *Genet Test*7:131–134

-Sekar KC and Corff KE. (2008);Treatment of patent ductus arteriosus: indomethacin or ibuprofen?*J Perinatol*.

-Sendelbach DM , Jackson GL , Lai SS et al.,(2008) . Pulse oximetry screening at 4hours of age to detect critical congenital heart defects . Pediatrics; 122 : e815 – e820(24] .

Steinberger J, Moller JH, Berry JM, et al.(2000): Echocardiographic diagnosis of heart disease in apparently healthy adolescents. Pediatrics;105:815-818.

Stoll and Kliegman RM,(2004):The fetus and neonatal infant .In Behrman RE, Kliegman RM and Jenson HB(eds).Nelson textbook of pediatrics. Saunders Elseviers Science.Philadelphia. P519 – 23.

Sullivan ID,(2002): Prenatal diagnosis of structural heart disease: Does it make a difference to survival? Heart 2002;87 :405-406.

Talner NS, Carboni MP(2000): Chest pain in the adolescent and young adult. Cardiol Rev;8:49-56.

Tautz 1 , C. Merkel 1 , F. Loersch 1 , O. Egen 1 , F. Hägele 2 , H.-M. Thon 3 , T. Schaible 1 Affiliations 1.(2010). Universitätsmedizin Mannheim, Department of Pediatrics, Mannheim, Germany

Todros T,(2000): Prenatal diagnosis and management of fetal cardiovascular malformations. Curr Opin Obstet Gynecol 2000;72:105-109.

Tsien CL, Fackler JC(1997): Poor prognosis for existing monitors in the intensive care unit. Crit Care Med, 25:614-619.

Valmari P . (2007). Should pulse oximetry be used to screen for congenital heart disease? Arch Dis Child Fetal Neonatal Ed ; 92 92: F 219– F 224.

Van Overmeire B, Smets K, Lecoutere D, et al.(2000). A comparison of ibuprofen and indomethacin for closure of patent ductus arteriosus. *N Engl J Med*.

Volkl TM, Degenhardt K, Koch A, Simm D, Dorr HG, Singer H (2005). "Cardiovascular anomalies in children and young adults with Ullrich-Turner syndrome the Erlangen experience". *Clin Cardiol* **28**.

Webb GD, Smallhorn FS, Therrien J, Redington AN.(2005): Congenital heart disease. In Zies DP. Libby P, Bonow RO, Braunwald E (ed). *Braunwald heart disease*. 7th edition. Elsevier Saunders. Philadelphia. P1489 – 547 .

Weeks B and Friedman AH.(2004): Training pediatric residents to evaluate congenital heart disease in the current era. *Pediatr Clin North Am* 51,:1641.-1.651.

Wilson W, Taubert KA, Gewitz M, et al. (2007) Prevention of infective endocarditis: guidelines from the American Heart Association: a guideline from the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation*. ;116(15):1736-54.

Wolfson BJ.(2001): Radiologic interpretation of congenital heart disease. *Clin Perinatol* 28:7L-89.

Wren C, Birrell G, Hawthorne G. (2003) Cardiovascular malformations in infants of diabetic mothers. *Heart* .

References

Wu MH, Wang JK, Lin MT, et al.,(2006). Ventricular septal defect with secondary left ventricular-to-right atrial shunt is associated with a higher risk for infective endocarditis and a lower late chance of closure. *Pediatrics*. Feb ;117(2):e262-7. .

Yetman AT, Rosenberg HC and Joubert GI(1995). Progression of asymptomatic aortic stenosis identified in the neonatal period. *Am J Cardiol*. Mar 15.
