

IX- References

- Agapova OA, Kaufman PL, Lucarelli MJ: Differential expression of matrix metalloproteinases in monkey eyes with experimental glaucoma or optic nerve transection. *Brain Res* 2003;967(1-2): 132-43.
- Airaksinen PJ, Heijel A: Visual field and retinal nerve fiber layer in early glaucoma after optic disc haemorrhage. *Acta Ophthalmol (Copenh)* 1983; 61:186.
- Ajtony Csilla ,Zolat Balla ,Szabolcs Somoskeoy and Balint Kovacs :The Relationship Between Visual Field Sensitivity and Retinal Nerve Fiber Layer Thickness as measured by Optical Coherence Tomography .*IVOS* 2007;258-263.
- Allingham RR : Visual fields and their relationship to the optic nerve. In Epstein DL, Allingham RR, Schuman JS (eds): *Chandler & Grants Glaucoma*. 4th edition. Williams and Walkins (A Waverly Company), Baltimore, Boston, Chicago.1997 Chapter 1p 3-8.
- American Academy of Ophthalmology preferred practice patterns committee glaucoma panel. Preferred practice patterns. Primary open angle glaucoma. San Francisco: American Academy of Ophthalmology; 2001:1-38.
- American Academy of Ophthalmology .Primary open –angle glaucoma , Preferred practice pattern .San Francisco :American Academy of Ophthalmology ; 2005.

- Anderson DR, Pattella VM .Automated static perimetry ,2nd edition .st. louis :Mosby ;1999:10-35..
- Anton A, Moreno –Montanes J ,Blazques F ,Alvarez A, Martin B: Usefulness of Optical Coherence Tomography parameters of the optic disc and the retinal nerve fiber layer to differentiate glaucomatous ,ocular hypertensive and normal eyes .J Glaucoma 2007, 16(1):1-8.
- Asano E, Mochizuki K, Sawada A ,Nagasaka E, Kondo Y, Yamamoto T: Decreased nasal –temporal asymmetry of the second order kernel response of multifocal electroretinogram in eyes with normal tension glaucoma .Jpn J Ophthalmol 2007;51(5):379-89.
- Asman P, Heijl A.: Diffuse visual field loss and glaucoma .Acta Ophthalmol (Copenh) ,1994;72:303-8.
- Aydin A , Wollesten G , Price L L , Fujimoto JG, Shumann JS: Optical Coherence Tomography Assessment of Retinal Nerve Fiber Layer Thickness Changes after Glaucoma Surgery : Ophthalmology .2003.
- Badala F, Nouri-Mahdavi K, Roof DA, Leeperchanon N, Law K, Caprioli J: Optic disc and nerve fiber layer imaging to detect glaucoma. AJ Ophthalmol 2007;144(5):724-32.
- Baga H, Fever WJ, Greenfield DS: Detection of psychophysical and structural injury in eyes with glaucomatous optic neuropathy and normal standard automated perimetry .Arch Ophthalmol 2006;124(2) :169-76.

- Bearse Jr MA, Sutter EE, Smith DN: Ganglion cell components of the human 51 multifocal ERG are abnormal in optic nerve' s atrophy and glaucoma. IVOS 1995; 36 (suppl): S444.
- Bearse Jr MA, Sim D, Sutter EF: stamper R, Leiberman M. Application of the multifocal ERG to glaucoma. IOVS 1996; 37(suppl): S511
- Bearse, M. A., Jr., & Sutter, E. E. Imaging localized retinal dysfunction with the multifocal electroretinogram. Journal of the Optical Society of America A, 1996., 13, 634-640. [PubMed].
- Bearse MA, Sutter EE, Sim D, Stamper, R : Glaucomatous dysfunction revealed in higher order components of the electroetinogram. Vision Science and Its Applications, OSA Technical Digest Series 1, 1996., 104-107 Optical Society of America Washington, DC.
- Bearse, M. A., Sutter, E. E. and Palmowski, A. M. : New developments toward a clinical test of retinal ganglion cell function. In Vision Science and Its Applications , OSA Technical Digest Series, 1, pp. 1997., 280-283. Optical Society of America, Washington, DC.
- Bearse, M. A. and Sutter, E. E. : Contrast dependence of multifocal ERG components. In Vision Science and its Applications, OSA Technical Digest Series, 1, pp.1998., 24-27. Optical Society of America, Washington, DC.
- Bearse MA, Jr, Stamper RL, Sutter EE : Detection of functional abnormalities in glaucoma using a new multifocal ERG paradigm; IVOS 41 (4), 2000., S103 Abstract nr 536.

- Bearse MA, Sutter EE, Stanoer RL : Detection of glaucomatous dysfunction using a global flash multi focal electroretinogram (mERG) paradigm OSA Technical Digest Series, 14-17 Optical Society of America Washington, DC. Vision Science and Its Applications, 2001., Vol. 1.
- Bengetsson B, Heijl A, Olsson J: Evaluation of a new threshold visual field strategy, SITA, in normal subjects. *Acta Ophthalmol Scand* 1998.,76:165.
- Bengetsson B, Heijl A, : Intersubject variability and normal limits of SITA standard, SITA Fast, and Humphery full threshold computerized perimetry strategies,SITA STAPAC. *Acta Ophthalmol Scand* ,1999., 77:125.
- Bill A.: Vascular physiology of the optic nerve. In: *The optic nerve in glaucoma*, Varma R., Spaeth GL., (eds.) JB Lippincott, Philadelphia 1993, p. 37.
- Brigell M, Bach M, Barber C, Moskowitz A, Robson J. :Guidelines for calibration of stimulus and recording parameters used in clinical electrophysiology of vision. *Doc Ophthalmol* 2003;107: 185-93.
- Bruncato Rosario ; Lumbraso Bruno: Guide to optical coherence tomography interpretation . *Innovation-News_Comunication* 2004 ;71-75.
- Budenz DL : Glaucomatous visual field loss. In *Atlas of Visual Fields*. Lippincott Raven Publishers, Philadelphia, New York.1997; Chapter 5, p. 143-193.

- Budenz L Donald ,Robert T chang ,Xinagrum ,Robert W Kington and James M Tielesh : Reproducibility of Retinal Nerve Fiber Thickness measurements using the Stratus OCT in Normal and Glaucomatous eyes .IVOS 2005 ;46:2440-2443.
- Budenz DL ,Michael A ,Chaang RT ,McSoley J,Katz J :Sensitivity and Specificity of the Sratus OCT for perimetric glaucoma ; Ophthalmology 2005,112(1):3-9.
- Budenz DL,Fredette MJ ,Feur WJ,Anderson DR:Reproducibility of peripapillary retinal nerve fiber thickness measurements with stratus OCT in glaucomatous eyes ;.Ophthalmology 2007,115(4):661-666.
- Butt Z, McKillop G, O'Brien C, et al: Measurement of ocular blood flow velocity using color Doppler imaging in low tension glaucoma. Eye 1995;9: 29-33.
- Caprioli .1, Kitano S. Morgan JE: Hyperthermia and hypoxia increase tolerance of retinal ganglion cells to anoxia and excitotoxicity. IVOS 1996;37:237d- X I.
- Chan, HL. and Brown, B. : Multifocal ERG changes in glaucoma. Ophthal. Phy vio/.1999; Opt. 19, 306-316.
- Chan HL. Brown B. Pilot study of the multifocal electroretinogram in ocular hypertension. Br J ophthalmol 2000; 84 147-1153.
- Chan HL: Detection of glaucomatous damage using multifocal ERG. Clinical and experimental Optometry 2005; 88:6:410-414.
- Chaumman Bc. The relationship between intraocular Pressure and visual field progression in glaucoma. In: Drance SM, editor. Update to

glaucoma, blood flow, and drug Treatment Amsterdam: Kugler, 1995. p.I-6.

- Chaumman BC, LeVatte TL, Jollimore CA, et al: Model of endothelin - 1- induced chronic optic neuropathy in rat. WOS 2004;45: 144 52.
- Chen HY,Wang TH ,Lee YM, Hung TJ ,Retinal Nerve Layer Thickness measured by Optical Coherence Tomography and its Correlation with visual field defects in early glaucoma .J Formos Med Assoc.2005 Dec ;104(12):927-34.
- Chihara and Sawada A.: A typical nerve fiber layer defects in high myopes with high-tension glaucoma. Arch Ophthalmol 1990. 108: 228-232.
- Cholpin NT, Edwards RP : Basic Principles of Visual Field Interpretation In Visual Field Testing with Humphrey Field Analyzer. SLACK Incorporated, USA.1995; Chapter 5, p. 69-98.
- Chu PH,Chan H H L ,Brown B: Luminance –modulation adaptation of global flash mfERG:Fellow eye losses in asymmetric glaucoma ,IVOS 2007,48;2626-2633.
- Chu PH,Chan H H L ,Brown B:Glaucoma detection is facilitated by luminance modulation of the global flash multifocal electroretinogram .IVOS 2006;47:929-937.
- Cioffi GA. And Van Buskrik EM.: Anatomy of the optic nerve. In Textbook of Ophthalmology. Wright KW.(ed.). William and Wilkins, Baltimore, 1997, p. 591.

- Ciula TA ,Regillo CD,Harris A:Retina and Optic Nerve Imaging .LIPPINCOTT WILLIAMS &WILKINS ,2003 213-230.
- Costa VP, Harris A, Stefansson E, et al: The effects of antiglaucoma and systemic medications on ocular blood flow. Prog Retin Eye Res 2003;22:769-805.
- Drance S, Anderson DR, Schulzer M: Risk factors for progression of visual field abnormalities in normal-tension glaucoma. Am J Ophthalmol 2001; 131 :699-708.
- Dreyer EB: A proposed role for excitotoxicity in glaucoma. J Glaucoma 1998;7:627.
- Eid TM and Speath GL :Pathogenesis of glaucomatous optic nerve damage . IN:The glaucoma concepts and fundamentals Lippincort Philadelphia,2000,P.63-71.
- EL Beltagi TA, Christopher Bowd ,Catherine Boden ,et al :Retinal Nerve FiberLayer Thickness Measured with Optical Coherence Tomography Is Related to Visual Function in Glaucomatous Eyes .Ophthalmology 2003;volume110,Number11,2185-2191.
- EL Sanbary Z,Osman L ,Kamal M and Hassan N: Comparison between optical coherence tomography and scanning laser phthalmoscopy in evaluating retinal nerve fibre layer thickness in glaucoma patients .Bull .ophthalmol soc Egypt ,1999;677-681.
- Flammer J, Eppler E_ Niesel P: Quantitative perimetry in glaucoma patient without local visual field defects: Graefes Arch Clin Exp Ophthalmol ,1982; 219:92

- Flammer J.: Normal values in computerized perimetry. In : computerized visual fields. Walen WR. And Spaeth G., (eds.). Slack Inc, thorofare NJ, 1984,p. 159-164.
- Flammer .1, Orgul S, Costa VP, et al. The impact of ocular blood flow in glaucoma. *Prog Retin Eye Res* 2002; 21 :359-93.
- Fortune B, Cioffi, GA, Johnson CA, Kondo Y, Mochizuki K, Kitazawa Y : The relationship between multifocal electroretinogram and standard automated perimetry findings in normal tension glaucoma Weinreb, RN Krieglstein, GK Kitazawa, Y eds. *Glaucoma in the 21st Century*, 2000; 73-78 Harcourt Publishers London.
- Fortune B, Cioffi GA, Johnson CA : Glaucoma causes selective reduction of an oscillatory component in multifocal ERG responses from the temporal retina. *IVOS* 2001; 42(4), S147 Abstract nr 783.
- Fortune B, Johnson CA, Cioffi, GA : The topographic relationship between multifocal electroretinographic (MERG) and behavioral perimetric measures of function in glaucoma. *Optom Vis Sci.*2001; 78, 206-214
- Fortune B, Cull G, Wang L, Van Buskirk EM, Cioffi GA :Factors affecting the use of multi focal electroretinography to monitor function in a primate model of glaucoma. *IVOS*,2001
- Fortune Brad, Marcus A, Bearse Jr, George A, Cioffi and Chris A. Johnson : Selective loss of an oscillatory component from temporal retinal multi focal ERG responses in glaucoma. *IVOS* 2002, 43: 2638-2647.

- Frishman LJ, Saszik S, Harweth RS, et al. : Effects of experimental glaucoma in macaques on the multifocal ERG: sin laser-induced glaucoma. *Doc Ophthalmol* 2000, 231-251.
- Fukuchi T, Sawaguchi S, Hara H, et al. Extracellular matrix changes of the optic nerve lamina cribrosa in monkey eyes with experimentally chronic glaucoma. *Graefes Arch Clin .p Ophthalmol* 1992;230:421.
- Funaki S., Shirakashi M., and Abe H.:Relation between size of the optic disc and thickness of retinal nerve fiber layer in normal subjects. *Br J Ophthalmol* 1998. 82:1242-1245.
- Galassi F, Sodi A, Ucci F, Harris A, Chung HS: Ocular haemodynamics in glaucoma associated With high myopia. *Int ophthalmol* 1998; 22: 299-305.
- Galassi F. Sodi A, Ucci F. et al: Ocular hemodynamics and glaucoma II" Prognosis: ocular Doppler imaging study. *Arch Ophthalmol* 2003;121:1711-5.
- Garway-Health DF,Holder GE,Fitzke FW, et al .: Relation between electrophysiologic ,psychophysical and anatomic measurements in glaucoma .*IVOS* 2002;43:2213-2220.
- Glass E,Schaumberger M,Lachenmayer BJ ;Simulations for FASTPAC and the standerd 4-2 dB full threshold strategy of the Humphery field analyzer ;*IVOS* 1995,36-1847.
- Goldstic BJ,Weinreb RN: The effect of refractive error on automated global analysis program G-1 *Am J Ophthalmol* 1987;104-229.
- Graham SE, Drance SM: Nocturnal hypotension: role in glaucoma, - *Progress. Surv Ophthalmol* 1999;43(SupplI): SI 0-6.

- Grodum K., Heijj A. ,Bengtsson: Refractjve error and glaucoma. Acta ophthalmol Scand 2001;79: 560-6.
- Haefliger IO, Flammer :J : Fluctuation of the differential light threshold at the border of absolute scotomas. Comparison between glaucomatous visual field, defects and blind spots. Ophthalmology.1991; 98:1529.
- Hare W A, Ton H, Ruiz G, et al : Characterization of retinal injury using ERG measures obtained with both conventional and multifocal methods in chronic ocular hypertensive primates. IVOS ;2001; 42, 127-136 .
- Harewerth RS,Carter-Dawson L,Smith ELIII Brances G ,Hot WF,Crawford ML:Neural losses correlated with visual losses in clinical perimetry .IVOS 2004;45:3152-3160.
- Harewerth RS,Vilupuru AS,Rangaswamy NVand Smith EL:The relationship between nerve fiber layer thickness and perimetry measurements .IVOS 2007;48:763-773.
- Harrington DO : The visual Fields. A textbook and Atlas of Clinical Perimetry 5th ed,1981; - St Louis, CV Mosby.
- Hasegawa, S., Takagi, M., Usui, T., Takada, R. and Abe, H. : Waveform changes of the first-order multifocal electroretinogram in patients with glaucoma. IVOS. 2000; 41, 1597-1603.
- Hayreh S, Zummennan ME, Podhajsky P, el al: Nocturnal arterial hypotension and its role in optic nerve head and ocular ischemic disorders. Am J Ophthalmol 1994;11 7:603-24.

- Hayreh SS, Podhajsky p, Zimmerman MB: Beta-blocker eyedrops and nocturnal arterial hypotension. *Am J Ophthalmol* 1999;128:301-9.
- Hee MR, Izatt JA, Swanson EA, Huang D, Schuman JS, Lin CP, Puliafito CA, Fujimoto JG: Optical coherence tomography of the human retina. *Arch Ophthalmol* 1995; 113:325-32.

- Heijl ,A : The implications of the results of computerized perimetry in normals for the statistical evaluation of glaucomatous visual fields. In Krieglstein GK (ed.): *Glaucoma update III*, Berlin. Springer-Verlag.1987. Quoted in Stamper RL,: Lieberman MF, Drake MV : *BeckerShaffer's Diagnosis and Therapy of the Glaucomas*. 7th edition, Section III. Mosby Inc. St Louis, Baltimore, Boston, Chicago.1999; Chapter 11p. 144:171
- Heijl A, Bengtsson B: The effect of perimetric experience in patients with glaucoma, *Arch Ophthalmol*.1996; 114: 19.
- Hernandez MR, Pena JD. The optic nerve head in glaucomatous optic neuropathy. *Arch Ophthalmoi* 1997;115:38995.
- Hernandez MR. Pena .ID, Selvidge 1A: Hydrostatic pressure stimulates synthesis of elastin in cultured opticnerve head astrocytes. *Cilia* 2000;32: 12236.
- Hitzenberger CK. Optical measurement of axial eye length by laser Doppler interferometry. *IVOS*, 1991; 32:616-624.

- Holder GE, Brigell M, Hawlina M, Meigen T, Vaegan, Bach M. Standard for pattern electroretinography. *Doc Ophthalmol* 2007;114:111-
- Hong S,Ahn H ,HaSJ ,Yeom HY ,Seong GH ,HongYJ: Early glaucoma detection using the Humphery Matrix Perimeter ,GDX,VCC,stratus OCT and retinal nerve fiber layer photography .*Ophthalmology* 2007;114(2):210-5.
- Hood DC, Seiple, W, Holopigian, K, Greenstein, V. : A comparison of the components of the multifocal and full-field ERGs. *Vis Neurosci* 1997;14, 533-544
- Hood, D. c., Greenstein, V., Frishman, L., Holopigian, K., Viswanathan, S., Seiple, W., Ahmed, J. and Robson, J. G. : Identifying inner retinal contributions to the human multifocal ERG. *Vis. Re.* 1999; 39, 2285-2291.
- Hood, D. c., Frishman, L J., Viswanathan, S., Robson, J. G. and Ahmed, J. : Evidence for a ganglion cell contribution to the primate electroretinogram (ERG): effects of TTX on the multifocal ERG in macaque. *Vis. Neuro.* 1999; 16, 411--416.
- Hood DC. Assessing retinal function with the multifocal technique. *Prog Retin Eye Res* 2000; 19: 607-646.
- Hood, D. c., Frishman, L J., Saszik, S., Viswanathan, S. and Robson, J. G. : The variations in the waveform of the primate cone ERG with retinal eccentricity depend upon inner and outer retinal contributions. *IVOS* 2000; 41(suppl), S496.

-
- Hood DC, Greenstein VC, Holopigian K, et al., : An attempt to detect glaucomatous damage to the inner retina with the multifocal ERG. *IVOS* .2000;41, 1570-1579
 - Hood DC, Bearnse MA, Sutter EE, Viswanathan S, Frishman LJ : The optic nerve head component of the monkey's (*Macaca mulatta*) multifocal electroretinogram (mERG). *Vision Res* .2001;41, 2029-2041.
 - Hood DC, Frishman LJ, Viswanathan S. Retinal origin of the primate multifocal ERG : implications for the human response .*IVOS* 2002;43:1673-1685.
 - Hood DC , Harizman FN , Kandani FN , et al : Retinal nerve fibre thickness measured with optical coherence tomography accurately detects confirmed glaucomatous damage .*BJO*,2006;91:905-907.
 - Hoskins HD Jr, Migliazzo D : Development of visual field screening test using a Humphrey Visual Field Analyser, *Doc Ophthalmol* 1985; 42:85.
 - Hougaard JL, Heijl LA , Bengtsson B : Glaucoma detection By Stratus OCT .*J Glaucoma* ,2007 ;16(3):302 -6.
 - Huang O, Swanson EA, Lin CP, Schuman JS, Stinson WG, Chang W, Hee MR, Flotte T, Gregory K, Puliafito CA, Fujimoto JG. Optical coherence tomography. *Science* 1991; 254:1178-1181.
 - Ji JZ, Elyaman W, Yip HK, et al: CNTF promotes survival of retinal ganglion cells after induction of ocular hypertension in rats: the

possible involvement of STAT3 pathway. *Eur J Neurosci* 2004; 19: 265-72.

- Johnson CA, Keltner JL : Optimal rates of movement for kinetic perimetry, *Arch Ophthalmol*.1987; 105: 73
- Johnson CA, Adams AJ, Lewis RA: Evidence for a neural basis of age-related visual field loss in normal observers. *IVOS* 1989;30: 2056.
- Johnson CA, Nelson-Quigg JM : a prospective three-years study of response properties of normal subjects and patients during automated perimetry, *ophthalmology*.1993; 100:269.
- Johnson LN : Effects of intermittent versus continuous patient monitoring on reliability indices during automated perimetry, *Ophthalmology*.1993; 100: 76.
- Johnson EC, Deppnieicr LM, Wentzien SK, et al. Chronology of optic nerve head and retinal responses to elevated intraocular pressure. *IVOS* 2000;41:431-42.
- Jonas JB., Schmidt AM., Muller – Eberg JA.,Schlotzer UM., and Naumann GO.: Human optic nerve count and optic disc size. *IVOS*. 1992.33: 2012- 2018.
- Jonas JB, Dichtl A. Optic disc morphology in myopic Primary open angle glaucoma. *Gates Arch Clin Exp ophthalmol* 1997;235:627-33.
- Jonas JB,Budde WM,Panda Jonass: Ophthalmoscopic evaluation of the optic nerve head .*Surv Ophthalmol* 1999;43:293.
- Kanamori Akiyasu ,Makoto Nakamura ,Michael F.TEscano ,Ryu Seya ,Hidetaka Maeda .Evaluation of the Glaucomatous damage on retinal

nerve fiber layer thickness measured by Optical Coherence Tomography .Am J of ophthalmology .2003 ;volume 135,issue 4,513-520.

- Kass MA, Kolker AE, Becker B : Prognostic factors in glaucomatous visual field loss. Arch. Ophthalmol 1976;. 94: 1274.
- Katz j,Congdon N,Frideman DS .Methodological variations in estimating apparent visual field loss in clinical trials of glaucoma treatment ,Arch .Ophthalmol .1999;117:1137-42.
- Keltner JL and Johnson CA:Baseline visual field profile of optic neuritis ,Arch Ophthalmol 1993;111-312.
- Kerrigan BLA, Lack D.I, Quigley HA:, et al. TUNEL. positive ganglion cells in human primary open-angle glaucoma. Arch Ophthalmol 1997;115:H»15.
- Kerrigan –Baumrind LA , Quigley HA, Pease ME ,et al .Number of ganglion cells in glaucoma eyes compared with threshold visual field tests in the same persons .IVOS 2000;41:741-748.
- Kim Tae –Woo,Un-Chul Park ,Ki-HO Park ,and DongMyung Kim :Ability of Stratus OCT to identify localized retinal nerve fiber layer defects in patients with normal standerd Automated Perimetry results. IVOS 2007;48:1635-1641.
- Klistorner AI, Graham SL, Martins A : Multifocal pattern electroretinogram does not demonstrate localized field defects in glaucoma. Doc Ophthalmol 2000; 155-165.

- Kondo M, Miyake Y, Horiguchi M, Suzuki S, Tanikawa A. Clinical evaluation of multifocal electroretinogram. *IVOS* 1995; 376: 2146-2150
- Kondo M, Miyake Y, Horiguchi M, Suzuki S, Tanikawa A. Recording multifocal electroretinogram on and off responses in humans. *IVOS* 1998 39:574-580
- Kondo, M., Miyake, Y., Piao, C., Tanikawa, A., Horiguchi, M. and Terasaki, H. :Amplitude increase of the multifocal electroretinogram during light adaptation. *IVOS* 2000;40, 2633-2637.
- Kretschmann U, Bock M, Gockeln R, Zrenner E : Clinical applications of multifocal electroretinography. *Doc. Ophthalmol* .2000;100,99-113.
- Kuehn, Markus H., Fingert, John H., Kuron, Young H.: Retinal ganglion cell death in glaucoma : mechanisms and neuroprotective strategies. *Ophthalmology clinics of North America*. 2005; 441-473.
- Lee M, Zulauf M, Caprioli J: The influence of patient reliability on visual field outcome, *Am J Ophthalmology* .1994;117:756-61.
- Leung Christopher Kai-shun, Wai –manchan, Kelvin Kan –long chong, et al: Comparative Study of Retinal Nerve Fibre Layer Measurement by Stratus OCT GDX VCC, I: Correlation Analysis in Glaucoma: *IVOS*, 2005, volume .46 No.9, 3214-3219.
- Leung Christopher Kai –Shun, Arthur Chak Kwan Cheng, Kelvin Kam Lung Chang, et al : optic disc Measurements in myopia with Optical Coherence Tomography and Confocal Scanning Laser Ophthalmoscopy. *IVOS* 2007, 48:3178-3183.

- Leung CKL, Medeiros FA, Zanguill LM, Sample PA, et al : American Chinese glaucoma imaging study, A comparison of the optic disc and retinal nerve fibre layer in detecting glaucomatous damage. *IVOS* 2007; 48: 2644-2652.
- Lindenmuth KA, Skuta GL, Rabbani R, Musch DC : Effects of papillary constriction on automated perimetry in normal eyes. *Ophthalmology* 1989; 96:1298.
- Lynn JR, Fellman RL,Starita RI : Principles of perimetry .In :Ritch R,Shields MB,Krupin T,eds. *The glaucomas* ,2nd ed. St.Louis ;Mosby –Year Book;1996:491-521.
- Maeda H, Tanaka Y, Nakamura M, Yamamoto m:: Blue-on-yellow perimerty using an armaly glaucoma screening program *Ophthalmologica*.1999; 213:71.
- Manassakorn A,Nouri –mahadavi K,Caprioli J: Comparison of retinal nerve fibre layer thickness and optic disc algorithms with optical coherence tomography to detect glaucoma; *Am J Ophthalmol* 2006;141(1) :105-115.
- Marmor MF, Holder GE, Seeliger MW, Yamamoto S: International Society for Clinical Electrophysiology of Vision. Standard for clinical electroretinography (2004 update). *Doc Ophthalmol*. 2004;108:107-14.
- Martin KR, Quigley IIA, Zack DJ: et al. Gene therapy with brain-derived neurotrophic factor as a protection: retinal ganglion cells in a rat glaucoma model. *IVOS* 2003;44: 4357 65.

- Maruyama I, Nakazawa M, Ohguro H: Autoimmune, mechanisms in molecular pathology of glaucomatous optic neuropathy. *Nippon Ganka Gakkai Zasshi* 2001; 105:205- 12 .
- Mattox C.: Clinical examination of the nerve fiber layer. In: *Imaging in Glaucoma*. Schuman JS (ed.). Slack, Thirifare NJ., 1999, p. 63-80.
- McLeod SO, West S. Quigley HA, et al: A longitudinal study of the relationship between intraocular pressure and blood pressure. *IVOS* 1990; 31 :2361-6.
- Mickelberg FS, ,Schulzer M, Dtance SM, Lau W : The rate of progression of scotomas in glaucoma. *Am J Ophthalmol*.1986; 01: 1.
- Mickelberg F., Drance S., Schluzer M., Yidegiligne H., and West H.: The normal human optic nerve axon- diameter distribution. *Ophthalmology* 1989. 96: 1325- 1328.
- Morrison C John ,Pollack P Irnin : *Glaucoma Science and Practice* 2003.80-89.
- Naithani Parashant ,Ramanjiti Shiota ,Tanuji daola , et al: Evaluation of Optical Coherence Tomography and Heidelberg retinal Tomography Parameters in Detecting Early and moderate glaucoma, *IVOS* ,2007 ;48:3138-3145.
- Nagatomo, A., Nao-i, N., Mariuiwa, F., Arai, M. and Sawada, A. : Multifocal electroretinograms in normal subjects. *Jpn. J. aptha*.1998; 42, 129-135.

- Nasemann IE, Carl T, Pamer S, et al: Perfusion time of the central retinal artery in normal pressure glaucoma: initial results. *Ophthalmologie* 1994; 91: 595-601.
- Nicolela MT, Buckley AR, Walman BE, et al: A comparative study of the effects of timolol and latanoprost on blood flow velocity of the retrobulbar vessels. *Am J Ophthalmol* 1996; 122:784-9.
- Okisaka S, Murakami A, Mizukawa A, et al. Apoptosis in retinal ganglion cell decrease in human glaucomatous eyes. *Jpn J Ophthalmol* 1997;41:84 S.
- Orgul. S. and Cioffi G.: Embryology, anatomy, and histology of the optic nerve vasculature. *J Glaucoma* 1996. 5:285.
- Ortega Julio E Deleon ,Stella N Arthur ,Gerald MC Gwin Jr ,et al :Discrimination between Glaucomatous and non glaucomatous eyes using quantitative imaging devices and subjective optical nerve head assessment .*IVOS* 2006 ;47:3374-3380.
- Ortega Julio E Deleon ,Stella N Arthur ,Gerald MC Gwin Jr et al :Effect of Glaucomatous Damage on Reproducibility of confocal Scanning Laser Ophthalmoscope ,Scanning Laser polarimetry and Optical Coherence Tomography .*IVOS* 2007;48;1156-1163.
- Osborne NN, Wood JP, Chidlow G, Bae JH, Melena J, Nash MS : Ganglion cell death in glaucoma: what do we really know? *Br J Ophthalmol* .1999; 83, 980-986.

- Osman ZM: Glaucomatous Optic Neuropathy ; Pathophysiology . Glaucoma News 2003; vol 3; No2 ; 1-8.
- Palmowski AM, Allgayer R, Heinemann-Vernaleken B : The multifocal ERG in open angle glaucoma: a comparison of high and low contrast recordings in high- and low-tension open angle glaucoma. Doc Ophthalmol 2000; 101, 35-49 .
- Palmowski AM, Wolf Aja M, Todorova M, Orguel S, Flammer J, Brigell : An attempt to analyse the sensitivity of two global flash , mfERG to detect glaucomatous dysfunction in normal tension glaucoma and high tension (POAG) patients . Documenta Ophthalmologica 2007; 114, 9-19.
- Parks, S., Keating, D., Evans, A., L., Williamson, T. H., Jay, J. L., & Elliott, A. T. : Comparison of repeatability of the multifocal electroretinogram and Humphrey perimeter. Documenta Ophthalmologica, 1997; 92, 281-289.
- Paunescue A, Leila , Joel S Schuman , Lori lyn Price , Paul C Stark , Siobahn Beaton , Hiroshi Ishikawa , Gadi Wollsteiu , and James G Fujimoto : Reproducibility of nerve fiber thickness , macular thickness and optic nerve head measurements using stratus OCT . IVOS 2004 ; 45: 1716-1724.
- Pearson HE, Stoffler D.I: Retinal ganglion cell degeneration following loss of postsynaptic target neurons in the dorsal lateral geniculate nucleus of the adult cat. Exp Neural 1992; 116: 163-71.

- Pena JD, Varela HJ, Ricard CS: Enhanced tenascin expression associated with reactive astrocytes in human optic nerve heads with primary open angle glaucoma. *Exp Eye Res* 1999;68:29-40.
- Pennebaker GE, Stewart WC: Temporal visual field in glaucoma: a reevaluation in the automated perimetry era. *Graefes Arch Clin Exp Ophthalmol* 1992;30:103-111.
- Puliafito CA, Hee MR., Shuman JS and Fujimoto JC: Optical Coherence Tomography of ocular Diseases. Slack Thorofare NJ, 1996, p.291-344.
- Quaranta L, Manni G, Donato F, et al: The effect of increased intraocular pressure on pulsatile ocular blood flow in low tension glaucoma. *Surv Ophthalmol* 1994;38(Suppl):S 177-81.
- Quigley H, Addicks E: Regional differences in the structure of lamina cribrosa and their relation to glaucomatous optic nerve damage. *Arch Ophthalmol* 1981; 99: 137.
- Quigley HA, Addicks EM, Green WR, et al: Optic nerve damage in human glaucoma. II. The site of injury and susceptibility to damage. *Arch Ophthalmol* 1981;99:635.
- Quigley HA, Hohman RM, Addicks EM, et al. Morphologic changes in the lamina cribrosa correlated with neural loss in open-angle glaucoma. *Am J Ophthalmol* 1983;95:673.
- Quigley HA: Early detection of glaucomatous damage. II. Changes in the appearance of the optic disc. *Surv Ophthalmol* 1985; 30: 111-126.

- Quigley H., Sanchez R , dunelberger G., and Bangniski T.: Chronic glaucoma selectively damages larger optic nerve fibers. *IVOS* 1987. 28: 913-920.
- Quigley H., et al.: The size and shape of the optic disc in normal human eyes. *Arch Ophthalmol* 1989. 108:51-57.
- Quigley H. Calemar Al: and Dorman – pease ME.: Larger optic nerve heads have more nerve fiber in normal monkey eyes. *Arch Ophthalmol* 1991: 1441-1443.
- Quigley HA, Nickells RW, Ken-igan LA: Retinal ganglion cell death in experimental glaucoma and after axotomy occurs by apoptosis. *Invest ophthalmol Vis Sci* 1995;36:774-86.
- Quigley HA : Neuronal death in glaucoma. *Prog Retinal Eye Res.*1999; 18, 39-57
- Rangasawy NV, Weil Zhou ,Ronald Sharwerth and Laura J.frihman : Effect of Experimental Glaucoma in primates on oscillatory potentials of the slow –sequence mfERG . *IVOS* 2006;47:735-767.
- Ravalico G, Pastori G, Toffoli G, et al: Visual and blood flow responses in low-tension glaucoma. *Surv Ophthalmol* 1994;38(suppl):S 173-6.
- Raz D,Perlman I,Percicot CL, Lambrou GN, Ofri R:Functional damage to inner and outer retinal cells in experimental glaucoma ;*IVOS* 2003,44(8)3675-84.
- Read R and Spaeth G.: Partical clinical appraisal of the optic disc in glaucoma. The natural history of glaucomatous cupping. *Trans Am Acad ophthalmol Otolaryngol* 1974. 78 :225.

- Rhee JD (2003):Color Atlas and Synopsis of clinical Ophthalmology. MCCraw –Hill ,78-103.
- Richard CS, Kobayashi S, Pena JD: Selective expression of neural cell adhesion molecule (NCAM)-ISO in optic nerve head astrocytes exposed to elevated hydrostatic pressure in vitro. Brain Res Mol Brain Res 2000;81(1-2):62- 79.
- Roh S., Noecker RJ.: and Schuman JS.: Evaluation of coexisting optic nerve head drusen and glaucoma with optical coherence tomography. Ophthalmology 1997. 104 :1138 – 1144.
- Roh S., et al.: Effect of optic nerve head drusen on nerve fiber layer thickness. Ophthalmology 1998.105: 878-885.
- Schuman JS ., Kloizmam TP . Hertzmark F., Hee MR and Puliafito CA : Rproducibility of nerve fiber layer thickness measurements using optical coherence tomography .ophthalmology 1996. 103:1889 – 1898.
- Schuman J, Orgul S, Gugleta K, et al: Interocular difference V in progression of glaucoma correlates with interocular differences ,in retrobulbar circulation. Am J Ophthalmol 2000; 129:728-33.
- Schuman J,Wolstein g,Fara T et al : Comparison of optic nerve head measurement obtained by optic coherence tomography and confocal scanning laser ophthalmoscopy .AM J Ophthalmol .2003;135 :504-512.
- Schwartz B: cupping and pallor of the optic disc. Arch Ophthalmol 1973; 89:272.

- Seeliger, M. W., Kretschmann, U. H., Apfelstedt-Sylla, E. and Zrenner, E.: Implicit time topography of multifocal electroretinograms. *IVOS* (1998a). 39,718-723.
- Sehi Mitra ,Stephen Uime ,David S :Greenfield and advanced imaging in glaucoma study group :Scanning Laser Polarimetry with enhanced corneal compensation and optical coherence tomography in normal and glaucomatous eyes .*IVOS* ;2007 ,3146-3153.
- Shields MB,Allingham RR,Damji KF ,Freedman S, Moroi SE; Shafranof G: Shields textbook of glaucoma ; LIPPINCOT WILLIAMS &WILKINS 2005,99-135.
- Shiota R,Sony P,Gupta V,Data T,Singh R. 2006:Diagnostic capability of optical coherence tomography in evaluating the degree of glaucomatous retinal nerve fiber damage .*IVOS* 2006,47(5):2006-10.
- Shiota R , Sony P , Cupta V ,DaDa T and Singh R :Diagnostic Capability of Optical Coherence Tomography in Evaluating the Degree of Glaucomatous Retinal Nerve Fiber damage .*IVOS* 2006;47:2006-2010.
- Shorestein NH, Dawson WW, Sherwood MB: Mid peripheral pattern electrical retinal responses in normals, glaucoma suspects and glaucoma patients. *Br. J. ophthalmol*1.1999, 1: 15.
- Sieving, P. A., Murayama, K. and Naarendorp, F.: Push-pull model of the primate photopic electroretinogram: a role for hyperpolarizing neurons in shaping the b-wave. *Neuro.vci*. 1994 11, 519-532.

-
- Sommer A, Tielsch IM, Katz J, et al. Racial differences in the cause-specific prevalence of blindness in East Baltimore, N Eng! J Med 1991;325:1412-7.
 - Spaeth GL, Varma R (1987): Assessment of glaucomatous patients. Eye. 1: 29.
 - Spaeth GL (2007): The colored glaucoma graph : A useful aid in the diagnosis and management of patients with glaucoma . Highlights of ophthalmology volume 35 ,number 3,17-26.
 - Spalton J David ,Hilshings A Roger(1994) :Atlas of clinical Ophthalmology ,17.2-17.27.
 - Stamper RL., Liberman MF., and Drake MV.: Optic nerve anatomy and pathophysiology. In: Diagnosis and therapy of the Glaucoma : Mosby, St Louis, 1999, p 177-190.
 - Stewart WC,Shields MB,Ollie AR:peripheral visual field testing by automated kinetic perimetry in glaucoma .Arch Ophthalmol 1988;106-202.
 - Sutter EE : The fast m-transform: fast computation of cross correlations with binary m-sequences Society of Industrial and Applied Mathematics Journal on Computing .1991; 20,686-694.
 - Sutter, E. E., & Tran, D. : The field topography of ERG components in man—I. Documenta Ophthalmologica, 1992; 95, 73-90.
 - Sutter EE, Tran D : The field topography of ERG components in man. II: the photopic luminance response. Vision Res.1992; 32, 433-446 .
 - Sutter, E. E. and Bearse, M. A. Extraction of a ganglion cell component from the corneal response. In Vision Science and its

-
- Applications, OSA Technical Digest Series. Washington DC Optical Society of America, 1995; 1:310-3
- Sutter EE, Bearnse Jr MA. The optic nerve head component of the human ERG. *Vision Res* 1999; 39: 419-436.
 - Sutter EE, Shimada Y, Li Y, Bearnse MA : Mapping inner retinal function through enhancement of adaptive components in the mERG OSA Technical Digest Series 1, 52-55.1999; Optical Society of America Washington, DC. *Vision Science and Its Applications*, Vol. 1.
 - Sutter EE, Bearnse MA, Shimada Y, Li Y : A multi focal ERG protocol for testing retinal ganglion cell function [AROV Abstract]. *IVOS1999*; 40(4), S 15 Abstract nr 79.
 - Sutter, E. E. : Imaging visual function with the multifocal m-sequence technique. *Vision Research*,2001; 41, 1241-1255.[PubMed]
 - Sutter EE, Bearnse MA, Stamper RL, et al : Monitoring retinal ganglion cell function with the mERG: recent advances OSA Technical Digest Series, 10-13 Optical Society of America Washington, DC. *Vision Science and Its Applications*,2001; Vol. 1.
 - Swanson EA, Huang D, Hee MR, Fujimoto JG, Lin CP, Puliafito CA: High-speed optical coherence domain reflectometry. *Opt Lett* 1992; 17:151-153.
 - Swanson EA, Izatt JA, Hee MR, Huang D, Lin CP, Schuman JS, Puliafito CA, Fujimoto JG. In vivo retinal imaging by optical coherence tomography. *Opt Lett* 1993; 18:1864-6.

- Takada K, Yokohoma I, Chida K, Noda J: New measurement system for fault location in optical waveguide devices based on an interferometric technique. *Appl Opt* 1987; 26:1603-1606.
- Tbrien E, Murphy J, Tyndall A, et al: Twenty-four hour ambulatory Blood pressure in men and women . *J Hypertens* 1991,9.355.
- Tezel G, Wax MB: Hypoxia-inducible factor 1 alpha in the chronic moderately elevated intraocular pressure. *IVOS* 1999;40:2884-91.
- Tezel G, Edward DP, Wax MB: Serum autoantibodies to optic nerve head glycosaminoglycans in patients with glaucoma. *Arch Ophthalmol* 1999; 17: 917-24.
- Trew DR, Smith SE.: Postural studies in pulsatile ocular blood flow, II: chronic open-angle glaucoma. *Br J Ophthalmol* 1991;75:71-5.
- Tsai CS., Ritch R., Shin DH., War JY., and Chi T.: Age related decline of disc rim area in visually normal subjects, *Ophthalmology* 1992.99:29- 35.
- Vaegan and Buckland, L. : The spatial distribution of ERG losses across the posterior pole of glaucomatous eyes in multifocal recordings. *NZ J. Ophthal* ;1997; 14(SuppI2), 28-31.
- Van Adel BA, Kostic C, Deglon N, et al: Delivery of ciliary neurotrophic factor via lentiviral-mediated transfer protect axotomized retinal ganglion cells for an extended period of time. *Hum Gene Ther* 2003; 14: 103-15.

- Varma R. and Minckler DS.: Anatomy and Pathophysiology of the retina and optic nerve. In: The Glaucoma. Ritch R., Shields MB., and Kupin T., (eds.). Mosby, St Louis, 1996. p. 139-176.
- Varma R. Skaf M., and Barron E.: Retinal nerve fiber layer thickness in normal human eyes. *Ophthalmology* 1996. 103: 2114 – 2119.
- Ventura LM, Okac NS, Santos RDL, Feuer WJ, Porciatti WJ: The relationship between retinal ganglion cell function and retinal nerve fiber thickness in early glaucoma ; *IVOS* 2006;47,3904-3911.
- Verdon, W. A., & Haegerstrom-Portnoy, G. :. Topography of the multifocal., Dietrich, T. J., Selig, B., Hofmann, C., & Schiller, J. (1999). Angioscotoma detection with fundus-oriented perimetry. A study with dark and bright stimuli of different sizes. *Vision Research*, 1999;39, 1897-1909.
- Verdon, W. A. and Haegerstrom-Portnoy, G. : Topography of the multifocal electroretinogram. *Doc. Ophthalmol* 1998; 95, 73-90.
- Wax MB: Is there a role for the immune system in glaucomatous optic neuropathy'. *curr Opin Ophthalmol* :2000;! 1:145 -50.
- Weber A.J., Kaufman L., Hubbard W.C.: "morphology of single ganglion cells in the glaucomatous primate retina". *IVOS* 1998;39:2304-2320
- Weinreb RN., Shakiba S., and Sample PA.: Association between quantitative nerve fiber measurement and visual field loss in glaucoma. *Am J Ophthalmol* 1995. 120: 732-738.

- Wolfs, Arend O, Sponsel WE, et al: Retinal heino-dynamics using scanning laser ophthalmoscopy and hemorrheology in chronic open-angle glaucoma. *Ophthalmology* 1993; 100: 1561-6.
- Yan DB, Coloboma FM, Metheetairut A,et al: Deformation of the lamina cribrosa by elevated intraocular pressure .*B J Ophthalmol* 1994;102-1693.
- Yan X, Tezel G, Wax MB, et al.: Matrix metalloproteinases and tumor necrosis factor alpha in glaucomatous optic nerve head. *Arch Ophthalmol* 2000;1'18:666-73.
- Youngquist Rc, Carr S, Davies DEN: Optical coherence-do main reflectometry: a new optical evaluation technique. *Opt Lett* 1987; 12:158-160.
- Zeyen TG, Caprioli J. Progression of disc and field damage in early glaucoma. *Arch Ophthalmol* 1993;111:62.