



REFERENCES

- Abu-Alfa, AK strous FH, Montag Ag, (1994):** An Immunohistochemical study of thyroid Hurthle cells and their neoplasms. The roles of S-100 and HMB-45 proteins. *Mod pathol* 7: 529-532.
- Akslen L A, Livolsi V A, (2000):** Prognostic Significance of Histologic Grading Compared with subclassification of papillary thyroid carcinoma. *Cancer* 88 (8): 1902-1908.
- Akslen LA, (1993):** Prognostic importance of histologic grading in papillary thyroid carcinoma. *Cancer* 72: 2680-5.
- Alessandri AJ, Goddard KJ, Blair GK et al., (2000):** Age is a major determinant of recurrence in pediatric differentiated thyroid carcinoma. *Med-pediat oncol* 35(1): 41-6.
- American Joint Committee on Cancer ,(1997):** Thyroid gland. Manual for staging of cancer 5th ed. Philadelphia: JB Lippincott, 47-50.
- Anwar F, Emond MJ, Schmdt RA et al., (2000):** Retinoblastoma expression in thyroid neoplasms. *Mod pathol* 13 (5): 562-9.
- Aogi K, Kitahara K, Urqundi V et al., (1999):** Comparison of Telomerase and CD44 expression as diagnostic tumor markers. *Clin Cancer Res.* 5 (10): 2790-7.
- Apel RL, Alpert LC, Rizzo A et al., (1994):** A metastasizing composite carcinomas of the thyroid with distinct medullary and papillary components. *Arch pathol Lab Med* 118: 1143-1147.
- Apel RL, Asa SL, Livolsi VA (1995):** Papillary Hurthle cell carcinoma with Lymphocytic stroma (warthin- tumor) of the thyroid. *Am J surg pathol* 19:810.



- Ashfaq R, Vuitch F, Delgado R et al., (1994):** Papillary and follicular thyroid carcinomas with an insular component. *Cancer* 73: 416-423.
- Avantaggiati ML, Orgyzko V, Gardner K et al., (1997):** Recruitment of P 300/CBP in P53 dependent signal pathways. *Cell* 89: 1175.
- Bakiri F, Djemli FK, Mokrane LA, (1998):** The relative roles of endemic goitre and socioeconomic development status in the prognosis of thyroid carcinoma. *Cancer* 82 (6): 1146-53.
- Basolo F, Caligo M, Pinchera A et al., (2000):** Cyclin D1 over expression in thyroid carcinomas: relation with clinicopathological parameters, retinoblastoma gene product and ki67 labeling index. *Thyroid* 19 (6): 192-5.
- Beckner ME, Heffess CS, Ortel JE, (1995):** Oxyphilic papillary thyroid carcinomas. *Am J Clin pathol* 103: 280-287.
- Bocklage. T, Ditomasso JP, Ramzy I et al., (1997):** Tall cell variant of papillary thyroid Carcinoma: Cytologic features and differential diagnostic considerations. *Diagn Cytopathol* 17: 25-29.
- Bohm J, Niskanen L, Tamimi R et al., (2000):** Hyaluronan expression in differentiated thyroid carcinoma. *J pathol.* 196 (2): 180-5.
- Bohm JP, Niskanen LK, Pirinen RT, (2000):** Reduced CD44 standard expression is associated with tumor recurrence and unfavorable outcome in differentiated thyroid carcinoma. *J. Pathol.* 192: 321-327.
- Boise LH, Gonzalez-GM, Postema CE et al, (1993):** Bcl-x, a Bcl-2 related gene that functions as a dominant regulator of apoptotic cell death. *Cell.* 74: 597-608.

- Cady B, Rossi R, (1988):* An example view of risk group definition in differentiated thyroid carcinoma. *Surgery* 104(6): 947-53.
- Callendr D, Sherman S, Gagel R et al., (1996):* Cancer of the thyroid. *Cancer of the head and neck*. In: Sue J, Myers E, (eds) 3rd ed philadelphia: Saunders, : 422.
- Carla BJ, Natalia T, Haart et al., (1999):* Cyclin D1 gene amplification and over-expression are present in ductal carcinoma in situ of the Breast. *J. Pathol* 187: 279-84.
- Cetani F, Tonachera M, Pinchera A et al., (1999):* Genetic analysis of the TSH receptor gene in differentiated thyroid carcinomas. *J endocrinol Invest*. 22 (4): 273-8.
- Chao DT, Korsmeyer SJ, (1998):* BCL-2 family: regulators of cell death. *Annu Rev Immunol* 16: 395.
- Chen H, Lin RJ, Schills L et al., (1997):* Nuclear receptor coactivator. ACTR is a novel histone acetyltransferase and forms a multimeric activation complex. *Cell* 90: 569.
- Cheng EH, Kirsch DG, Clem RJ et al., (1997):* Conversion of bcl-2 to a bax-like death effector by caspases. *Science* 278: 1966-96.
- Chetty R, Beydow R, Livolsi A, (1994):* Paraganglioma-like (Hyalinizing trabecular) adenoma of the thyroid. *Surgery* 18: 1131.
- Chetty R, Mills A, Livolsi V, (1993):* Anaplastic carcinoma of the thyroid with sclerohyaline nodules. *Endocr Pathol*. 4: 110-114.
- Chieng DC, Ross JS, Mckenna BJ, (1997):* CD44 immunostaining of thyroid fine needle aspiration differentiates thyroid papillary carcinoma from other lesions with nuclear grooves and inclusions. *Cancer Cytopathol* 81: 157-62.



- Dejong JS, Vandiest PJ, Baak JP, (2000):** Number of apoptotic cells as a prognostic marker in invasive breast cancer. *British J. Cancer.* 82(2), 368-373.
- Demeter JG, Dejong SA, Laurence AM et al., (1991):** Anaplastic thyroid carcinoma: Risk factors and outcome. *Surgery* 110: 956-963.
- Derwahi M, (1999):** Mutations in the thyrotropin receptor gene in the pathogenesis of toxic thyroid adenomas, toxic goitre nodules and autosomal dominant hyperthyroidism. *Zarzti, Fortbild Qualitat Ssich* 93 (1): 25-8.
- Derwahi M, Manole D, Sobke A, (1998):** Pathogenesis of toxic thyroid adenomas and nodules: relevance of activating mutations in the TSH-receptor and Gs-alpha gene. The possible role of iodine deficiency and secondary TSH-independent Molecular mechanisms. *EP. Clin Endocrinol-Diabetes* 106 (suppl 4): 56-9.
- De-Vathaire F, Hardiman C, Shausalol A et al., (1999):** Thyroid carcinomas after irradiation for a first cancer during childhood. *Arch. Intern Med* 159 (22): 27 13-9.
- Devita VT, Hellman S, Steven A, (1997):** Cancer-principles, practice of oncology. Lippincott.Raven, Pp. 1617-1631.
- Dillin A, Rine J, (1998):** Roles for ORC in M phase and S phase. *Science* 279: 1733.
- Donghi R, Longoni A, Pilotti, S et al., (1993):** Gene p53 mutations are restricted to poorly differentiated and undifferentiated carcinomas of the thyroid gland. *J. Clin. Invest.* 91: 1753-1760.
- Donnellan R, Chetty R, (1998):** Cyclin D1 and human neoplasia. *Mol pathol* 51 (1): 1-7.
- Dupetrez L, Parma J, Vansande J, et al., (1999):** J pediatri endocrinol metab. 12 (suppl 1): 295-302.



- Dynlacht BD, (1997):** Regulation of transcription proteins that control the cell cycle. *Nature* 389: 149.
- El-Bolkainy MN, (2000):** Endocrine tumors. In : Topographic pathology of cancer. NCI, Cairo University Ch (11) P. (89).
- Eng C, Mulligan LM, Healey CS, et al., (1996):** Heterogenous mutations of the RET proto-oncogene in sub-populations of medullary thyroid carcinoma. *Cancer Res.* May 1, 56 (6): 2167-70.
- Eng C, Thomas GA, Neubreg DS et al., (1998):** Mutation of the RET proto-oncogene is correlated in RET immunostaining in such population of cells in sporadic medullary thyroid carcinomas. *J-Clin-Endocrinol Met* 83 (12): 4310-3.
- Erickson LA, Long J, Goellner JR et al., (2000):** Pathologic, proliferative activity, and CyclinD1 expression in Hurthle Cell Neoplasms of the thyroid. *Modern Pathology* 13: 186-192.
- Ermak G, Gerasimov G, Troshinia K et al., (1995):** Deregulated alternative splicing of CD44 messenger RNA transcripts in neoplastic and non neoplastic lesion of the human thyroid. *Cancer Res.* 55 (20): 4594-98.
- Esapa CT, Johnson SJ, Kendall TP, (1999):** Prevalence of Ras mutations in thyroid neoplasia. *Clin Endocrinol oxf* 50 (4): 529-35.
- Farahati J, Bucsk YP, Pahloarsky T, (1997):** Characteristics of differentiated thyroid cancer in children and adolescents with respect to age, sex and histology. *Thyroid* (supplement I) Abstract.
- Farahati J, Demidchick EP, Biko J et al., (2000):** Inverse association between age at the time of radiation exposure and extent of diseases in cases of radiation induced childhood thyroid carcinoma in Belarus. *Cancer* 88 (6): 1470-6.



- Green D, Bissonnette R, Gotter T, (1999):** Apoptosis and cancer. In important advances in oncology. Vincent TD, Samuel Helmian (eds) 3rd ed. Lippincott company, Philadelphia.
- Gu J, Daa T, Kashima K et al., (1998):** Exprepsion of splice variants of CD44 in thyroid neoplasms derived from follicular cells Pathol-Int. 48 (3) 184: 190.
- GU W, Shi XL, Roedr RG, (1997):** Synergistic activation of transcription by CBD and P53. Nature 387: 819.
- Hamel PA, Hanley HJ, (1997):** G1 Cylclins and control of the cell division cycle in normal and transformed cells. Cancer Invest. 15 (2): 143-52.
- Hankard F, Cezard JP, Igrany et al, (1998):** CD44 variant expression in inflammatory colonic mucosa is not disease but associated with increased crypt cell proliferation. Histopathology 32: 317-321.
- Hanna NN, Mcgrath PC, Sloan A et al., (1999):** Advances in pathogenesis and treatment of thyroid cancer. Curr, opin, oncol 11 (1): 42-7.
- Hansen HS, Terring H, Goodballe C, et al., (2000):** Is thyroidectomy necessary in RET Mutations carriers of the familial medullary thyriod carcinoma syndrome? Cancer 89(4): 862-67.
- Harach H, Williams E, (1995):** Childhood thyroid cancer. Br.J Cancer. 72 (3): 777-783.
- Hay I, Bergstralh E, Goellner J, (1993):** Predicting outcome in papillary thyroid carcinoma: development of a relieable prognostic scoring system in a cohort of 1779 patients surgically treated at one institution during 19 40 through 1989. Surgary, 119: 1050.
- Hedinger C, Williams, DW, Leslie H,(1989):** The WHO Histological classification of thyroid tumors: A commentary on the second Edition Cancer. 63(5): 908-911.



- Hicks Dg, Livolsi VA, Puck JM et al., (1990):** Clonal analysis of solitary follicular nodules in the thyroid. *Am J Pathol* 137: 553-62.
- Hinze R, Gimmo R, Taubert H, (2000):** Regulation of proliferation and apoptosis in sporadic and hereditary medullary thyroid carcinoma and their putative precursor-lesions. *Virchows-Arch* 437 (3): 256-63.
- Hirvikoski P, Tammi R, Kumpulainen E et al., (1999):** Irregular expression of hyaluronan and its CD44 receptors is associated with metastatic phenotype in laryngeal squamous cell carcinoma. *Virchows Arch* 434: 37-44.
- Hoda SA, Reed RJ, (1993):** S-100 proteins Immununo reactivity differentiate follicular from papillary neoplasms of thyroid (abstract) *Lab Invest* 6: 38A.
- Holland JF, Frei E, Bast RC, et al., (1993):** *Cancer medicine* 3rd ed. Vol. (1) Lea, Febiger. Philadelphia. London. 857-859.
- Hoyt MA, (1997):** Eliminating all obstacles: regulated proteolysis in the eukaryotic cell cycle [comment]. *Cell* 91: 149.
- Hundahl SA, Fleming ID, Fremgen AM et al., (1998):** A national cancer database report on 53, 586 cases of thyroid carcinoma treated in the U.S. 1985-1995. *Cancer*, 83: 2638-48.
- Hundhal, SA, Cady B, Mazzaferri M et al., (2000):** Initial results from a prospective cohort study of 5583 cases of thyroid carcinoma treated in the United States during 1996. *Cancer* 89: 202-217.
- Hurban RH, Huvos AG, Traganos F et al., (1990):** Follicular neoplasms of the thyroid in men older than 50 years of age. A DNA flow cytometric study. *Am J clin pathol* 94: 527-32.



- Komminoth R, Roth J, Sarem AP et al., (1994):** Polysialic acid of the neural adhesion molecule in the human thyroid. A marker for medullary thyroid carcinomas and primary C-Cell hyperplasia. *Am J Surg. pathol* 18:339-411.
- Komoriya A, Packard Pz, Brown MJ et al., (2000):** Assessment of caspase activities in intact apoptotic thymocytes using cell-permeable. Fluorogenic caspase substrates. *Journal of Experimental Medicine* 191: 1819-1828.
- Krajewski S, Krajewska M, Shabiak A et al., (1994):** Immunohistochemical analysis of in vivo distribution of bax, a dominant inhibitor of bcl-2. *Am J. Pathol.* 145: 1323-1326.
- Kroemer G, (1997):** The protooncogene Bcl-2 and its role in regulating apoptosis. *Nature Med.* 3: 614-620.
- Krude T, Jachman M, Pines J et al., (1997):** Cyclin/CdK dependent initiation of DNA replication in a human cell free system. *Cell* 88: 109.
- Lax SF, Behan A, Kronberger SG et al., (1994):** Coexistence of papillary and medullary carcinoma of the thyroid gland-mixed or collision tumor? Clinicopathological analysis of three cases. *Virchows Arch.* 424: 441-447.
- Leite RP, Carwo M, Cabecadas J et al., (1995):** Differential expression of cell adhesion molecules in functional compartments of lymph nodes and tonsils. *J clin Pathol: Mol Pathol* 48: 93-100.
- Leung CS, Haytwick Rw, Bedard YC, (1993):** Correlation of cytologic analysis of Histologic features in variants of papillary carcinoma of the thyroid. *Acta Cytol* 37: 645-650.
- Levine AJ, (1997):** P53, the cellular gate keeper for growth and division. *Cell* 88: 323.

- Li M, Rosai J, Carcangiu ML, (1995):** Hyalinizing trabecular adenoma of the thyroid. A distinct tumor type or a pattern of growth ?? Evaluation of 28 cases [abstract] Mod pathol 8: 54A. 1995.
- Livolsi V, Asa A, (1994):** The demise of follicular carcinoma of the thyroid. Thyroid 4: 233.
- Livolsi VA, Baloch ZW, (1999):** Determining the diagnosis and prognosis of thyroid neoplasms: Do special studies help? Hum Pathol. 30: 885-886.
- Louis H, Royseptions S, Kihany W, (1999):** Cancer of the thyroid gland In: Head and neck cancer: A multidisciplinary approach. Ch.33 3rded. P 799 Lippincott-Raven Publishers Philadelphia New York.
- Lui Q, Guntukus, Cui xs, (2000):** Chk1 is an essential kinase that is regulated by ATR and required for the G2/M DNA damage-checkpoint. Genes and Development 14: 1448-55.
- Lukas J, Herzinga T, Hansen K et al., (1997):** Cyclin E-induced S-phase. Without activation of the PRb/E 2F pathway. Genes Dev 11: 1479.
- MacDonald J, Haller D, Mcdaugall R, et al, (2000):** Endocrine system In: Clinical oncology. Martin DA. James OA, Alleu SL, John. EN (eds) 2nd ed. Chinchill Livingstone USA.
- Malenchenko AF, (1997):** Role of ecological factors in the pathogenesis of thyroid cancer in regions of byelarus exposed during the Chernobyl accident. Radiats Biol Radio ecol 37 (6): 890-1.
- Martinez PD, Fernandez JC, Hierro GC et al., (1996):** Follicular variant of papillary carcinoma of the thyroid: to what extent is fine needle aspiration reliable? Diagn Cytopathol 15:12-16.



- Massaferri E , (1993):** Thyroid carcinoma: Papillary and follicular. In: Mazzaferri E, Samaan N eds. Endocrine tumors. Boston: Blackwell Scientific publications, 278-333.
- Matias GX, Caiaxas AC, Cabezas RJ ,(1994):** Compound medullary-papillary carcinoma of the thyroid. True mixed tumor. Histopathology 25: 183-184.
- Mazzaferri EL, Jhiang SM, (1994):** Long-term impact of initial surgical and medical therapy on papillary and follicular thyroid cancer. Am J Med. 97: 418-28.
- Minn AJ, Boise LH, Thompson CB , (1996):** Bcl-x_s antagonizes the protective effects of BCL-XL J Biol Chem 271: 6306-6312.
- Miyashita T, Kitada S, Krajewski et al., (1995):** Over expression of Bcl-2 protein increases the half-life of P21 (Bax). J Bio Chem 270: 26049-26052.
- Mizukami Y, Michigishi T, Noumoure A et al., (1994):** Autonomously functioning (hot) nodule of the thyroid gland. A clinical and histopathologic study of 17 cases. Am J clin pathol 101: 29-35.
- Mizukami Y, Nogoushi M, Michi GT et al., (1992):** Papillary thyroid Carcinoma in Kanazawa, Japan: Prognostic significance of histological types. Histopathology 20: 243-250.
- Mochizuki M, Saito K, Kanazawa K ,(1992):** Benign follicular thyroid nodule composed of signet ring cells with PAS-negative thyroglobulin accumulation in dilated rough endoplasmic reticulum. Acta pathol Jpn 42: 111-114.
- Modigliani E, Cohen R, Campos JM et al., (1998):** Prognostic factors for survival and for biochemical cure in medullary thyroid carcinoma. Results in 899 patients. Clin Endocrinol 48: 265-73.



- Mokhtar N, (1998):** Molecular Pathology of cancer 1st ed. National Cancer institute, Cairo University 525-531.
- Molberg K, Albores SJ, (1999):** Hyalinizing trabecular adenoma of the thyroid gland. Hum pathol 25: 192-97.
- Moore D, Ohene D, Garcia B et al., (1998):** Apoptosis in Thyroid Neoplasms: Relationship with P53 and Bcl-2 expression. Histopathology, 32: 35-42
- Morikawa Y, Ishihara Y, Kawano I et al., (1995):** Cystic Squamous cell carcinoma of the thyroid. A possible new subgroup of intrathyroidal epithelial thymoma. Endocr pathol 6: 77-81.
- Muller H, Moroni MC, Vigo E et al., (1997):** Induction of S-phase entry by E2F transcription factors depends on their nuclear localization. Mol. Cell Biol 17: 5508-20.
- Muller-Hocker J, (1999):** Immunoreactivity of p53 Ki-67, and Bcl-2 in oncocytic adenomas and carcinomas of the thyroid gland. Hum. Pathol. 30: 926-933.
- Nikiforov Y, Gnepp DR, (1994):** Pediatric thyroid cancer after the chernobyl disaster. Cancer 74: 748-766.
- Nishida T, Katayama S, Tsujimoto M, (1996):** Clinicopathological significance of poorly differentiated thyroid carcinoma. Am. J. Pathol. 23: 205-211.
- Nordzig MA, Vansteenbrugge GJ, Verkai NS et al., (1997):** The prognostic value of CD44 isoforms in prostate cancer patients treated by radical prostatectomy. Clin. Cancer Res. 3: 805-815.
- Nuydens R, Dispersyn G, Gerdvan Ret al., (2000):** Bcl-2 protects against apoptosis-related microtubule alterations in neuronal cells. Apoptosis 5: 43-51.



- O'Connor L, Strasser A, O'Rielly LA et al., (1998):** A novel member of bcl-2 family that promotes apoptosis. *EMBO Journal* 17: 384-395.
- O'Riordan D, O'Brien T, Crotty T et al., (1995):** Multiple endocrine neoplasia type 2B: More than an endocrine disorder. *Surgery* 118 (6): 936-942.
- Ogilvys MD, Print CG et al., (1999):** Constitutive bcl-2 expression throughout the hemopoietic compartment affects multiple lineages and enhances progenitor cell survival. *Proceedings of National academy of sciences of the USA.* 96: 14943-48.
- Oren M, (1999):** Regulation of the P53 tumor suppressor protein. *Journal of biological Chemistry* 274: 36031-36034.
- Orlandi F, Caraci P, Mussa A et al., (2001):** Treatment of medullary thyroid carcinoma: An update. *Endocr-Relat-Cancer.* 8 (2): 135-47.
- Oyama T, Vickery AL, preffer FI et al., (1994):** A comparative study of flow cytometry and histopathological findings in thyroid follicular carcinomas and adenomas. *Human pathol* 25: 271-275.
- Pacini F, Romi C, Paolo M, (1995):** Early treatment of hereditary medullary thyroid carcinoma after attribution of multiple endocrine neoplasia type 2 gene carrier status by screening for the RET gene mutations. *Surgery* 118 (6): 1032.
- Pacini F, Vorontsova T, Molinavo E et al., (1999):** Thyroid consequences of the Chernobyl accident. *Acta Paediatr suppl.* 88 (433): 23-27.
- Papothi M, Micca MF, Favero A et al., (1993):** Poorly differentiated thyroid carcinomas with primordial cell component. A group of aggressive lesions sharing insular, trabecular and solid patterns. *Am. J. Surg. Pathol.;* 17: 291-301.



- Paulovich AG, Toczyski DP, Hatmell LH , (1997):** When checkpoints fail. *Cell* 88: 315.
- Peter A, Macsween R, Lowe D, (1997):** Recent advances in histopathology, No. 17 Churchill Livingstone 93-99.
- Peters G, (1994):** The D-type cyclins and their role in tumorigenesis. *J cell sci suppl*, 18: 89-96.
- Peters P, (1996):** Perturbation of cell cycle regulators in human cancer. *Cancer surv* 27: 351-67.
- Pierotti MA, Vignen P, Bongaszone I , (1998):** Rearrangements of RET and VTRKI tyrosine kinase receptors in papillary thyroid carcinomas. *Recent Results Cancer Res.* 154: 237-47.
- Pisarchik AV, Ermak G, Fomicheya , (1998):** The Ret/PTC1 rearrangement is a common feature of Chernobyl-associated papillary thyroid carcinomas from Belarus. *Thyroid* 8 (2): 133-9.
- Pittas AG, Adler M, Fazzari M et al., (2000):** Bone metastases from thyroid carcinoma: Clinical characteristics and prognostic variables in one hundred forty six patients. *Thyroid* 10 (3): 261-8.
- Pontecorvi A, Danese D, (1999):** Molecular genetics of benign thyroid disease. *Rays*, 24 (2): 197-214.
- Ravi R, Bedi H, Fuchs EJ et al., (1998):** CD 95 (Fas) induced caspase mediated proteolysis of NF-KB cancer Res 58: 882.
- Raymond K, Tan BS, Robert K et al., (1995):** Anaplastic Carcinoma of the thyroid: A 24-year. Experience. *Head, NECK* 41-47.
- Read W, Mangano J, (1995):** Thyroid cancer in the USA since accident at Chernobyl. *BMJ* 311 (7003): 511.



- Ries L, Miller B, Hankey B et al., (1994):** SEER cancer statistic review. National Institute of Health No. 94-2789.
- Rieu M, Lame MC, Richard A et al., (1995):** Prevalence of sporadic medullary thyroid carcinoma: the importance of routine measurements of serum calcitonin in the diagnostic evaluation of thyroid nodules. Clin. Endocrinol oxf 42 (5): 453-60.
- Roque L, Serpa A, Castedo S et al., (1999):** Significance of trisomy 7 and 12 in thyroid lesions with follicular differentiation: a cytogenetic and in situ hybridization study. Lab Invest 79 (4): 369-78.
- Rosai J, (1996):** Ackerman's surgical pathology. 8th ed. Ch9 P511. Mosby- Year book, Inc. Library of congress.
- Rosai J, Luisac M, Ronald AD, (1992):** Tumors of the thyroid gland in: Atlas of tumor pathology. ARMED FORCES Institute of Pathology Washington, D.C.
- Rossi RL, Majils S, Rossi M, (2000):** Thyroid Cancer. Surg-Clin-North-Mnn. 80 (2): 571-80.
- Rudzki Z, Jothy S, (1997):** CD44 and the adhesion of neoplastic cells J Clin Pathol: Mol Pathol 50: 57-71.
- Salvesen GS, Dixit VM, (1997):** Caspases: intracellular signaling by proteolysis. Cell 91: 443.
- Sambade C, Franssila K, Cameselles T et al., (1991):** Hyalinising trabecular adenoma: a misnomer for peculiar tumor of the thyroid Endocrinl pathology 26: 429.
- Sanders LE, Cady B, (1998):** Re-examination of Risk group and outcome of treatment. Arch Surg. 133: 419-425.
- Sato N, Oyama N, Koyama Y et al., (1989):** Do the level of nodal disease according to the TNM classification and the number of involved cervical nodes reflect prognosis in patients with differentiated thyroid carcinoma? J Surg Oncol 96 (3): 151-5.



- Schaffler A, Palitzsch KD, Hohnehme S, et al., (1998):** coexistent thyroiditis is associated with lower stage in thyroid carcinoma Eur J clin Invest 28 (10): 838-44.
- Schark C, Fulton N, Yashiro T et al., (1992):** The value of measurement of ras oncogenes and nuclear DNA analysis in the diagnosis of Hurthle cell tumors of the thyroid. World J surg 16: 745-751.
- Schneider A, Ron E, (1996):** Carcinoma of follicular epithelium. In: Braberman LE, Utiger RD (eds). Werner and Ingbar's the thyroid: a fundamental and clinical text 7th ed. Philadelphia: Lippincott-Raven Publishing, 902.
- Schurman G, Mattfeldt T, Feitcher G, (1991):** Flow cytometry and immunohistochemistry of follicular neoplasms of the thyroid. Hum pathol 22: 179-84.
- Scopa CD, Mela CM, Saradopoulon R, et al., (1993):** The significance of the grooved nucleous in thyroid lesions. Mod Pathol 6: 691-694.
- Sedlak TM, Oltrail ZN, Yang E et al., (1995):** Bcl-Xs antagonizes the effects of Bcl-x. Proc. Natl. Acad. Sci. USA 92: 7834-7838.
- Segal K, Arad A, Lubin, E, (1994):** Follicular carcinoma of the thyroid. Head Neck 16: 533-538.
- Sellers WR, Novit BG, Miyale S et al., (1998):** Stable binding to E2F is not required for the retinoblastoma protein to activate transcription, promote differentiation and suppress tumor cell growth. Genes Dev. 12: 95.
- Shaha AR, (2000):** Thyroid cancer: extent of thyroidectomy. Cancer control 7 (3): 229-39.
- Shaha AR, Loree TR, Shah K, (1995):** Prognostic factors and risk group analysis in follicular carcinoma of the thyroid. Surgery 18: 1131.

- Sheaf RJ, Graoutine M, Gordon M et al., (1997):** Cyclin E-CDK2 is a regulator of P27 Ki P1. *Genes Dev* 11: 1464.
- Sherr CJ (1996):** Cancer cell cycles. *Science* 274: 1672.
- Sherr CJ, Roberts JM, (1995):** Inhibitors of mammalian G1 Cyclin dependent kinases. *Genes Dev.* 9: 1149.
- Shi Y, Zou M, Varcondi E et al., (2001):** Cyclin D1 in thyroid carcinomas. *Thyroid* 11 (7): 709-10.
- Shieh SY, Ahn J, Tamai K et al., (2000):** The Human Homolog of chick point kinases Chk1 and cdChk2 (Chk2) phosphorylate P53 at multiple DNA Damage-inducible Sites. *Genes and Development* 14: 289-300.
- Sierra A, Castellsaguex S, Coll T et al., (1998):** Expression of death-related genes and their relationship to loss of apoptosis in T1 ductal breast carcinomas. *Int J cancer* 79: 103-110.
- Soares P, Cameselli TJ, Sobrinho SM, (1994):** Immunohistochemical detection of P53 in differentiated, poorly differentiated and undifferentiated carcinomas of the thyroid. *Histopathology* 24: 205-210.
- Sofer L, Resnitzki D , (1996):** Apoptosis induced by ectopic expression of Cyclin D1 but not Cyclin E. *Oncogene* 13(11): 2431-7.
- Steinmuller T, Kluppy J, Rayers N et al., (2000):** Prognostic factors in patients with differentiated thyroid carcinoma. *Eur-J-surg.* 166(1): 29-33.
- Sternberg S, Donald AA, Darryl C et al., (1999):** Diagnostic surgical pathology Vol. (1) Ch. (15) 3rd (ed.) Library of congress Cataloging in-Publication Data 544-552.
- Stewart Z, Mays D, Jeunfier A, (1999):** Defective G1-S cell cycle checkpoints function sensitizes cells to microtubule inhibitor-induced apoptosis. *Cancer Res.* 59: 3831-37.



- Surgenoya A, Shingug K, Kabayashi S, (1993):** Surgical strategies of differentiated carcinoma of the isthmus. *Head Neck* 15: 158-160.
- Suzuki A, Delapompa JL, Hakem R et al., (1997):** Brca₂ is required for embryonic cellular proliferation in the mouse. *Genes Dev.* 11: 1242.
- Takahashi T, Fujimori K, Simon SL et al., (1999):** Thyroid nodules, thyroid function and dietary iodine in the Marshal Islands. *Int J Epidemiol* 28 (4): 742-9.
- Takegawa K, Sakamori M, Koremoto M et al., (2000):** Large amount of Vitamin A has no major effects on thyroidal hormone synthesis in two-stage rat thyroid carcinogenesis model using N-bis (2 hydroxypropyl) nitrosamine and thiourea. *J. Toxicol-Sci* 25 (2): 67-75.
- Tallini G, Carcangiu ML, Rosai J, (1992):** Oncocytic neoplasms of the thyroid gland. *Acta Pathol Jpn* 42: 305-315.
- Tallini G, Garcia RG, Herrer A, et al, (1999):** Down regulation of p27 kipi and Ki-67/MIB1-labeling index support the classification of thyroid carcinoma into prognostically relevant categories. *Am. J. Surg. Pathol.* 23: 678-685.
- Tallini G, Ladamyi M, Rosai J et al., (1994):** Analysis of nuclear and mitochondrial DNA alterations in thyroid and renal oncocytic tumors. *Cytogenetic Cell Genet* 66: 253-259.
- Tannapfel A, Geissler F, Kockerling F, (1999):** Apoptosis and proliferation in relation to histopathological variables and prognosis in hepatocellular carcinoma. *J. Pathol.* 187: 439-445.
- Taylor SS, Mckeou F, (1997):** Kinetochore localization of murine Bu61 is required for normal mitotic timing and checkpoint response to spindle damage. *Cell* 89: 727.



- Theriot JA, Satterwhite LL, (1997):** New wrinkles in cytokinesis [News Comment]. *Nature* 385: 388.
- Thoreson SO, Akslen LA, Geattre S, et al., (1989):** Survival and prognostic factors in differentiated thyroid cancer: A multivariate analysis of 1.055 cases. *Br. J. Cancer* 59: 231-335.
- Toft NJ, Wintom DJ, Kelly J et al., (1999):** MSH2 status modulates both apoptosis and mutation frequency in the murine small intestine. *Proc Nat Academic USA* (in press).
- Tresler PA, Clark OH, (1997):** Prognostic factor in thyroid carcinoma. *Surg Oncol Clin. North Am* 6: 555-98.
- Vasilopoulos SR, Schultz P Samuel M, (1997):** Differentiated thyroid cancer in children and adolescents. Clinical outcome and mortality after very long term follow-up. *Thyroid* 7 (suppl 1): (Abstract).
- Viglietto G, Chiappetta G, Martiniz T et al., (1995):** RET/PTC oncogene activation is an early event in thyroid carcinogenesis. *Oncogene* 11 (6): 1207-1210.
- Villunger A, O'Reilly LA, Holler N et al., (2000):** FasL, bcl-2, G-CSF and P38 MAPK Regulators of distinct cell death and survival pathways in granulocytes. *J Expe. Med.* 192: 647-657.
- Visona A, Pea M, Bozzola L et al., (1991):** Follicular adenoma of the thyroid gland with extensive chondroid Metaplasia. *Histopathology* 18: 278-279.
- Wang C, Mayo MW, Baldium AJ, (1996):** TNF and cancer therapy-induced apoptosis, potentiation by inhibition of NF-KB. *Science* 274: 784.
- Wang JY, (1998):** Cellular responses to DNA Damage. *Current opinion in cell biology* 10: 240-247.



- Wang S, Lloyd RV, Hutzler MJ et al., (2000):** The role of cell cycle regulatory proteins, Cyclin D1 in the prognosis of thyroid cancer. *Mod. Pahtal* 13 (8): 882-7.
- Wang S, Wu J, Svas L et al., (1998):** The role of cell cycle regulatory proteins, cyclin D1, cyclin E, and P27 in thyroid carcinogenesis *Hum Pathol* 29 (11) 1304-9.
- Webb SJ, Nicholson D, Bubb VJ et al., (1999):** Caspase-mediated cleavage of APC results in amino-terminal fragment with a intact aramadillo repeat domain. *FASEB J* 13: 399-346.
- Weinberg RA, (1995):** The Retinoblastoma Protein and Cell Cycle Control. *Cell* 81: 323.
- Williams ED, (1995):** Mechanisms and pathogenesis of thyroid cancer in animals and man. *Mutat. Res.* 333 (1-2): 123-9.
- Williams GT, Smith CA, (1993):** Molecular regulation of apoptosis: genetic control of cell death. *Cell*.74: 777-779.
- Wingo P, Tong T, Boldens S, (1995):** Cancer statistics. *CA Cancer J. Clin* 45: 8-11.
- Wolowiec D, (1994):** Cyclins and proliferative pathology. *Acta Haematol Pol* 25(3): 195-203.
- Wyllie A, (1997):** Apoptosis, clues in the P53 murder mystery. *Nature* 389: 237.
- Wyllie AH, Bellamy CO, Bubb VJ, et al., (1999):** Apoptosis and Carcinogenesis. *British Journal of Cancer.* 80(1) 34-37.
- Wynford-Thomas D, (1993):** Molecular basis of epithelial tumor genesis: The thyroid model. *Clin Rev oncogenensis* 4:1.



- Yamashita T, Hosoda Y, Kameyawa K, et al., (1992):** Peculiar nuclear clearing composed of microfilaments in papillary carcinoma of the thyroid. *Cancer* 70: 2923-2928.
- Yang J, Lui X, Bhalla K et al., (1997):** Prevention of apoptosis by bcl-2 release of cytochrome C from mitochondria blocked. *Science* 275: 1129.
- Yeager TR, Devries S, Jarrad DF, et al., (1998):** Overcoming cellular senescence in human cancer pathogenesis. *Genes Dev.* 12: 163.
- Zedenius J, Auer G, Backdahl M et al., (1992):** Follicular tumors of the thyroid gland. Diagnosis, clinical aspects and nuclear DNA analysis. *World J surg.* 16: 589-594.
- Zhang P, Liegeois NJ, Wong et al., (1997):** Altered cell proliferation and differentiation in mice lacking P57 Ki P2 indicates a role in Beckwith-wiedemann syndrome. *Nature* 387: 151.
- Zidan J, Kassem S, Kuten A, (2000):** Follicular carcinoma of the thyroid gland: prognostic factors, treatment and survival *Am J Clin oncol* 23 (1): 1-5.
- Zou L, Stillman B, (1998):** Formation of a preinitiation complex by S-phase cyclin CDK-dependent loading of Cdc 45P onto chromatin. *Science* 280: 593.
- Zou M, Shi Y, Farid NR, et al., (1998):** Inverse association between Cyclin D1 over expression and retinoblastoma gene mutation in thyroid Carcinoma. *Endocrine*, 8(1): 61-64.