

REFERENCES

- 1) Adams, H.R. (1975 a).
Direct myocardial depressant effect of gentamycin.
Eur. J. Pharmac., 30 : 272.
- 2) Adams, H.R. (1975 b).
Cardiovascular depressant effects of neomycin and gentamycin in rhesus monkeys.
Br. J. Pharmac., 54 : 453.
- 3) Adams, H.R. and Goodman, F.R. (1975).
Differential inhibitory effects of neomycin on contractile response of various canine arteries.
J. Pharmac. Exp. Ther., 193 : 393.
- 4) Adams, H.R. and Mathew, B.P. (1973).
Cumulative neuromuscular blocking effect of neomycin.
Arch. Int. Pharmacodyn., 210 : 288.
- 5) Adams, H.R., Goodman, F.R. and Weiss, G.B. (1974).
Alteration of contractile function and calcium ions movement in vascular smooth muscle by gentamycin and other aminoglycoside antibiotic.
Antimicrob. Agents and Chemother., 5 : 640.
- 6) Adams, H.R., Goodman, F.R. and Lupean, V.A. (1973).
Effect of neomycin on tension and ⁴⁵Ca movements in rabbit aortic smooth muscle.
Life Sci., 12 : 279.

- 13) Bulbring, E. (1946).
Br. J. Pharmac. Chemother., 1 : 38.
Quoted from Pharmacological Experiments on Isolated Preparations by the Staff of Department of Pharmacology, University of Edinburgh, p. 30.
E. S. Livingstone, Edinburgh and London (1970).
- 14) Bulbring, E. and Burn, J.H. (1942).
Physiol., 101 : 244.
Quoted from Burn, J.H., Finney, D.J. and Goodwin, L.G. Biological Standardization, 2nd edition, p. 352 (1952). Oxford University Press.
- 15) Burn, J.H. (1952).
Quoted from Pharmacological Experiments on Isolated Preparations by the Staff of Department of Pharmacology, University of Edinburgh, p. 38.
E. S. Livingstone, Edinburgh and London (1970).
- 16) Chenoweth, M.B. (1956).
Chelation as a mechanism of Pharmacological action.
Pharmac. Rev., 8 : 57.
- 17) Corrado, A.P. (1963).
Respiratory depression due to antibiotics, calcium in treatment.
Anaesth. Analg., 42 : 1.
- 18) Corrado, A.P., Ramos, A.O. and De Escobar, C.I. (1959).
Neuromuscular blockade by neomycin, potentiation by ether anaesthesia and d-tubocurarine and antagonism by calcium and prostigmine.
Arch. Int. Pharmacodyn. Ther., 121 : 380.

- 19) Cox, C.E. (1970).
Gentamycin.
Med. Clin. North Am., 64 : 1305.
- 20) Davidson, E.W., Modell, J.H., Moya, E. and Farmati,
Q. (1966).
Respiratory depression following use of anti-
biotics in pleural and pseudocyst cavity.
J.A.M.A., 196 : 456.
- 21) Del Castillo, J. and Engbac, E. (1954).
The nature of neuromuscular block by magnesium.
J. Physiol., 124 : 370.
- 22) Del Castillo, J. and Katz, B. (1954).
The effect of magnesium on the activity of
motor end plate.
J. Physiol., 124 : 553.
- 23) Del Castillo, J. and Stark, L. (1952).
The effect of calcium ions on the motor end
plate potential.
J. Physiol., 116 : 507.
- 24) De Rosaryo, M. and Healy, T.E.J. (1978).
Tobramycin and neuromuscular transmission in
the rat isolated phrenic nerve-hemidiaphragm
preparation.
Br. J. Anaesth., 50 : 251.
- 25) Dretchen, K.L., Gergise, S.D., Sokoll, M.D. and Long,
J.P. (1972).
Effect of various antibiotics on neuromuscular
transmission.
Eur. J. Pharmac., 18 : 201.

- 26) Dretchen, K.L., Sokoll, M.D., Gergis, S.D. and Long, J.P. (1973).
Relative effects of streptomycin on motor nerve terminal and end plate.
Eur. J. Pharmac., 22 : 10.
- 27) Drury, P.J. and Healy, T.E.J. (1975).
Neuromuscular transmission and gentamycin. Recent progress in anaesthesiology and resuscitation, edited by A. Arias.
Amsterdam Excerpta Medica, p. 276-280.
- 28) Dunkley, B., Songhvi, I. and Goldstein, G. (1973).
Characterization of neuromuscular block produced by streptomycin.
Arch. Int. Pharmacodyn. Ther., 201 : 213.
- 29) Elmqvist, D. and Josefsson, (1962).
The nature of the neuromuscular block produced by neomycin.
Acta Physiol. Scand., 54 : 105.
- 30) Emery, E.R.J. (1963).
Neuromuscular blocking properties of antibiotics as a cause of postoperative apnea.
Anaesth., 18 : 57.
- 31) Fairhurst, A.S. and Macri, J. (1975).
Aminoglycosides - Ca^{++} interactions in skeletal muscle preparation.
Life Sci., 16 : 1321.
-

- 39) Jindal, M.N. and Deshpande, V.R. (1960).
Neuromuscular block by streptomycin and
dihydrostreptomycin.
Br. J. Pharmac., 15 : 505.
- 40) Kaiser, A.B. and Mc Gee, Z.A. (1975).
Aminoglycoside therapy of gram -ve bacillary
meningitis.
N. Engl. J. Med., 293 : 1215.
- 41) Kaplan, J.M.M., Cracken, G.H. Jr., Thomas, M.L.H.,
Houston, L.J. and Davis, N. (1973).
Clinical pharmacology of tobramycin in newborn.
Am. J. Dis. Childr., 125 : 656.
- 42) Katz, B. and Miledi, R. (1964).
Localization of calcium action at the nerve
muscle junction.
J. Physiol. (Lond.), 171 : 10.
- 43) Katz, B. and Miledi, R. (1965 a).
The effect of calcium on acetylcholine release
from motor nerve terminal.
Proc. R. Soc. B., 161 : 496.
- 44) Katz, B. and Miledi, R. (1965 b).
Release of acetylcholine from a nerve terminal
by electric pulses of variable strength and
duration.
Nature, London, 207 : 1097.
- 45) Kawaguchi, H., Noits, T., Nakagawa, S. and Fujijawa,
K. (1972).
A new semisynthetic aminoglycoside antibiotic.
J. Antibiot. (Tokyo), 25 : 695.

- 46) Keller, H., Krupe, W., Saus, H. and Muchter, H.
(1956 a).
Experiments concerning inhibition of toxicity
of basic streptomycin.
Arzneimittel-Forsch., 6 : 61.
- 47) Keller, H., Krupe, W., Saus, H. and Muchter, H.
(1956 b).
Experiments on the reduction of toxicity of
basic streptomycin antibiotic.
Arzneimittel-Forsch., 6 : 585.
- 48) Koelle, G.B. (1970).
Neuromuscular blocking agents in the Pharmacological Basis of Therapeutics, p. 578.
Ed. by L.S. Goodman and A. Gilman. The
Macmillan Company, New York.
- 49) Kubikowski, P. (1963).
Quoted from Sobek (1982).
- 50) Lawrence, P. Garrod, Harold P. Lambest and Francis
O. Grody (1981).
Antibiotic and Chemotherapy, 5th Edition.
Churchill Livingstone. Medical Division of
Longman Group Limited.
- 51) Lee, B.K., Condon, R.G., Wagman, G.H. and Katz, E.
(1976).
Quoted from Lawrence et al. (1981).
- 52) Line, D.H., Pool, G.W. and Waterworth, P.M. (1970).
Quoted from Lawrence et al. (1981).

- 53) Lüllman, H. and Reuter, H. (1960).
Chemotherapia, 1 : 375.
Quoted from Sobek (1982).
- 54) Martin Dale (1977).
Streptomycin, Tobramycin. The Extrapharmacopia,
27th Edition, pp 1180, 1195. Edited by Aimely
Wade.
- 55) Matsuzoki, M., Nakamura, K., Akustu, S., Sekino, M.,
Hirata, A., Asono, M., Kishimoto, K. and
Fukushima, M. (1975).
Studies on the toxicity of Amikacin (BB, K 8,1),
acute toxicity and subacute toxicity in rats.
Japan. J. Antibiotics, 28 : 4.
- 57) Mc Lean, F.C. and Hastings, A.B. (1934).
A biological method for estimation of calcium
ion concentrations.
J. Biol. Chem., 107 : 337.
Quoted from Pittinger (1970).
- 58) Miller, R.D. (1975).
Factors affecting the action of muscle re-
laxants.
Amsterdam, Excerpta Medica, p. 163.
- 59) Molitor, H., Graessle, O.E., Kuna, S., Mushelt, C.W.
and Silber, R.H. (1946).
Some toxicological and pharmacological proper-
ties of streptomycin.
J. Pharmac. Exp. Ther., 86 : 151.
- 60) Molitor, H., Grasselle, O.E., Kuna, S., Mushelt, C.W.
and Silver, R.H. (1950).
Pharmacology and toxicology of antibiotics.
Pharmac. Rev., 2 : 1.

- 61) Moore, E.W. (1970).
Ionized calcium in normal serum ultrafiltrates
and whole blood determined by ion exchange
electrode.
J. Clin. Invest., 49 : 318.
- 62) Nastuk, W.L. and Liu, J.H. (1966).
Muscle post junctional membrane changes in
chemosensitivity produced by calcium.
Science (Washington), 15 : 266.
- 63) Paradelise, A.G., Triantaphyllidis, C., Fidani, V.
and Logares, G. (1974).
The action of aminoglycoside antitubercular
gentamicin on isolated rat diaphragm.
Arzneim. Forsch., 24 : 1774.
- 64) Paradelise, A.G., Triantaphyllidis, C., Markomichelakis,
J.M. and Logaras, G. (1977).
The neuromuscular blocking activity of amino-
deoxy kanamycin as compared with that of other
aminoglycoside antibiotics.
Arzneim. Forsch., 27 : 141.
- 65) Paradelise, A.G., Triantaphyllidis, C., Markomichelakis,
J.M., Salpigides, G. and Logaras, G. (1978).
Neuromuscular blocking activity of aminoglyco-
side antibiotics interaction with calcium
muscle relaxants and B-adrenergic blocking
agents.
Abstract 957, 7th International Congress of
Pharmacology, Paris, France (1978).

- 66) Paradelise, A.G., Triantaphyllidis, C. and Giala, M.M. (1980).
Neuromuscular blocking activity of aminoglycoside antibiotics.
Meth. Find. Exp. Clin. Pharmac., 2 : 45.
- 67) Pittinger, C.B. (1970).
pH and streptomycin influences upon ionic calcium in serum.
Anaesth. Analg., 49 : 540.
- 68) Pittinger, C.B. and Adamson, R. (1972).
Antibiotic blockade of neuromuscular function.
Ann. Rev. Pharmac., 12 : 169.
- 69) Pittinger, C.B. and Long, J.B. (1958).
Neuromuscular blocking action of neomycin sulphate.
Antibiotic and Chemother., 8 (4) : 198.
- 70) Pittinger, C.B., Erysa, Y. and Adamson, R. (1970).
Antibiotic induced paralysis.
Anaesth. Analg., 49 : 487.
- 71) Prado, W.A., Corrado, A.P. and Marseillan, R.F. (1978).
Competitive antagonism between calcium and antibiotics at the neuromuscular junction.
Arch. Int. Pharmacodyn., 231 : 297.
- 72) Pratt, W.B. (1973).
Fundamental of chemotherapy, p. 63.
Oxford university Press, New York - London - Toronto.

- 73) Pridgeon, J.E. (1956).
Respiratory arrest thought to be due to intra-
peritoneal neomycin.
Surgery, 40 : 571.
- 74) Ream, C.R. (1963).
Respiratory and cardiac arrest after intra-
venous administration of kanamycin with re-
versal of toxic effect with neostigmine.
Ann. Int. Med., 59 : 384.
- 75) Regamey, C., Gordon, R.C. and Kirby, W.M.M. (1973).
Comparative pharmacokinetics of tobramycin and
gentamycin.
Clin. Pharmac. Ther., 14 : 396.
- 76) Renna, G., Siro-Brigioni, G. and Cuomo, V. (1981).
Comparative evaluation of the neuromuscular
blocking activity of three new aminoglycoside
antibiotics in rats.
Toxicology Letters, 9 : 107.
- 77) Rinehart, K.L. Jr. (1969).
J. Infec. Dis., 119 : 345.
Quoted from Lawrence et al. (1981).
- 78) Sabawala, P.B. and Dillon, J.B. (1959).
The action of some antibiotics on the human
intercostal nerve complex.
Anaesthesiology, 20 : 659.
- 79) Sande, M.A. and Mandell, G.L. (1980).
The antimicrobial agents (aminoglycosides).
The Pharmacological Basis of Therapeutics,
p. 1165. (L.S. Goodman and A. Gilman). The
Macmillan Company, N.Y.

- 80) Saurize-Kurtz, G. (1974).
Inhibition of membrane calcium activation by
neomycin and streptomycin in crab muscle fibers.
Pflugers Arch., 349 : 337.
- 81) Schatz, A., Bugie, S. and Waksman, S.A. (1944).
Streptomycin, a substance exhibiting anti-
biotic activity against gram +ve and gram -ve
bacteria.
Proc. Soc. Exp. Biol. Med., 57 : 244.
- 82) Siber, G.R., Cheverria, E., Smith, A.L., Paisly, J.W.
and Smith, D.H. (1975).
Pharmacokinetics of gentamycin in children and
adults.
J. Infec. Dis., 132 : 637.
- 83) Simon, V.K., Mosinger, E.M. and Malerczy, V. (1974).
Pharmacokinetic studies of tobramycin and
gentamycin.
Antimicrob. Agents Chemother., 3 : 454.
- 84) Singh, Y.N., Harvey, A.L. and Marsbol, I.G. (1978).
Antibiotic induced paralysis of the mouse
phrenic-nerve hemidiaphragm preparation and
reversibility by calcium and by neostigmine.
Anaesthesiology, 48 : 418.
- 85) Singh, Y.N., Marshall, I.G. and Harvey, A.L. (1978).
Some effects of aminoglycoside antibiotic
amikacin on neuromuscular and autonomic trans-
mission.
Br. J. Anaesth., 50 : 109.