

## **References**

**Akhtar S, Hameed A (2006):** Versatility of the sural fasciocutaneous flap in the coverage of lower third leg and hind foot defects. *J Plast Reconstr Aesthet Surg.* 59(8):839-45.

**Angrigiani C, Grilli D, Thorne CH (2001):** The adductor flap: a new method for transferring posterior and medial thigh skin *Plast Reconstr Surg* 107: 1725-31.

**Anne M and Keith L(2007):** Lower Limb In:"Essential Clinical Anatomy". 3rd edn. P319. Edited by Anne M and Keith L, Lippincott Williams & Wilkin, Philadelphia.

**Attinger C:(2000):** Plastic surgery techniques for foot and ankle surgery. In: Myerson M, ed. *Foot and Ankle Disorders.* Philadelphia: WB Saunders:627.

**Attinger CE and Ducic I (2007):** Foot and ankle reconstruction. In:"Grabb And Smith's Plastic Surgery".6th edn. Pp689-707. Edited by Thorne CH, Lippincott Williams & Wilkins, Philadelphia..

**Aust MC, Spies M, Guggenheim M, Gohritz A, Kall S, Rosenthal H, Pichlmaier M, Oehlert G, Vogt PM (2008):** Lower limb revascularisation preceding surgical wound coverage e An interdisciplinary algorithm for chronic wound closure. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 61: 925-933.

**Babst R, Khong KS (2007):** Minimally Invasive Surgery. In: "Principles of Fracture Management". 2nd expanded edn.Pp199-211. Edited by Ruedi TP, Buckley RE, Moran CG, Thieme, German, Thieme Verlag.

**Barclay TL, Cardoso DT, Sharpe DT, et al. (1982):** Repair of lower leg injuries with fasciocutaneous flaps. *British Journal of Plastic Surgery.* 35(2): 127-32.

**Basheer MH, Wilson SM, Lewis H, Herbert K (2008):** Microvascular free tissue transfer in reconstruction of the lower limb *Journal of Plastic, Reconstructive & Aesthetic Surgery* 61: 525-528.

## References

---

**Basile A, Stopponi M, Loreti A, Minniti de Simeonibus AU (2008):** Heel coverage using a distally based sural artery fasciocutaneous cross-leg flap: report of a small series. *J Foot Ankle Surg.* 47(2): 112-7.

**Baudet J, Pelissier P, Casoli V, and Caix P (2009):** Soleus flap In: "Flaps and reconstructive surgery". 1st edn., 424-438. Edited by Wei F and Mardini S., Elsevier, China..

**Bhattacharya V, Reddy GR, Goyal S, Bashir SA (2007):** Distally based tenosynovial sheath flap of peroneal tendons for exposed tendo Achilles: Preliminary report of five cases *Journal of Plastic, Reconstructive & Aesthetic Surgery* 60: 1204-1207.

**Bhattacharya V, Watts RK (2005):** Live demonstration of microcirculation in the deep fascia and its implication. *Plast Reconstr.Surg* 115: 458-63.

**Birke-Sorensen H, Malmsjo M, Rome P, Hudson D, Krug E, Berg L, Bruhin A, Caravaggi C, Chariker M, Depoorter M, Dowsett C, Dunn R, Duteille F, Ferreira F, Francos Martínez JM, Grudzien G, Ichioka S, Ingemansson R, Jeffery S, Lee C, Vig S, Runkel N; International Expert Panel on Negative Pressure Wound Therapy [NPWT-EP], Martin R, Smith J (2011):** Evidence -based recommendations for negative pressure wound therapy: Treatment variables (pressure levels, wound filler and contact layer): Steps towards an international consensus. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 64: S1-S16.

**Bitsch M, Saunte DM, Lohmann M, et al. (2005):** Standardised method of surgical treatment of leg ulcers. *Scand J Plast Reconstr Surg Hand Surg* 39: 162-9.

**Blondeel PN, Morris SF, Hallock GG, et al. (2006):** Anatomy of the integument of the lower extremity. In: " Perforator Flaps". Pp 542–577. Edited by Baltimore. QMP.

**Braakenburg A, O bdeijn MC, Feitz R, et al. (2006):** The clinical efficacy and cost effectiveness of the vacuum -assisted closure technique in the management of acute and chronic wounds: a randomized controlled trial. *Plast Reconstr Surg* 118: 390-7.

## References

---

**Bullocks J, Naik B, Lee E, et al (2006):** *Flow-through flaps: a review of current knowledge and a novel classification system. Microsurgery.*26: 439–449.

**Bullocks J, Naik B, Lee E, et al. (2010):** *Flow-through flaps: a review of current knowledge and a novel classification system. Microsurgery.* 26:439–449.

**Carriquiry C, Costa MA, Vasconez LO (1985):** *An anatomical study of the septocutaneous vessel of the leg. Plast. Reconstr. Surg.* 76: 354–361.

**Cavadas PC, Teran-Saavedra PP (2002):** *Combined latissimus dorsi-thora-codorsal artery perforator free flap: the “razor flap”. J Reconstr Microsurg.* 18: 29–31.

**Cervelli V, Gentile P, Grimaldi M (2008):** *Regenerative surgery: Use of fat grafting combined with platelet-rich plasma for chronic lower-extremity ulcers. Aesth Plast Surg;* 33:340–345.

**Chen SL, Chen TM, Chou TD, Chen SG, Wang HJ (2002):** *The distally based lesser saphenous venofasciocutaneous flap for ankle and heel reconstruction. Plast. Reconstr. Surg.* 110:1664 –1672.

**Chen SZ, Li J, Li XY, et al. (2005):** *Effects of vacuum-assisted closure on wound microcirculation: an experimental study. Asian J Surg ;*28:211-7.

**Chen X, Chen H, Zhang G (2010):** *Management of Wounds with Exposed Bone Structures using an Artificial Dermis and Skin Grafting Technique. Journal of Plastic, Reconstructive & Aesthetic Surgery* 63: e512–e518.

**Chen YL, Zheng BG, Zhu JM, et al. (1985):** *Microsurgical anatomy of the lateral skin flap of the leg. Ann. Plast. Surg.* 15: 313-8.

**Chun JK, Marin M (2003):** *Radial artery flow-through flap for distal lower-extremity arterial bypass for limb salvage. Plast Reconstr Surg.* 111: 361–365.

**Coulier B (2002):** *Hyperechogenicity of medial gastrocnemial veins during ultrasound scanning of the calf in sitting patients: a normal variant. Eur. Radio.* 112: 1843–1848.

## **References**

---

**Crowley DJ, Kanakaris NK, Giannoudis PV (2007):** Debridement and wound closure of open fractures: The impact of the time factor on infection rates. *Injury Int. J.* 38: 879-889.

**Darwish AM (2010):** Bipedicled flap in reconstruction of exposed tibia.. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 63: 160-163.

**Davidson JM, Krieg T, Eming SA (2000):** Particle-mediated gene therapy of wounds. *Wound Repair Regen.* 8(6):452-9.

**Demirtas Y, Ayhan S, Sariguney Y, et al. (2006):** Distally based lateral and medial leg adipofascial flaps: need for caution with old, diabetic patients. *Plast Reconstr Surg* . 117: 272–276.

**Demling RH,. Niezgoda JA, Haraway GD, Mostow EN (2004):** Small intestinal submucosa wound matrix and full-thickness venous ulcers: preliminary results. *Wounds;* 16(1):18–22 .

**Dini V, Romanelli M, Piaggese A, Stefani A, Mosca F (2006):** Cutaneous tissue engineering and lower extremity wounds(Part 2). *Int J Low Extrem Wounds.* 5(1): 27–34 27.

**Dong JS, Peng YP, Zhang YX, et al. (2003):** Reverse anterior tibial artery flap for reconstruction of foot donor site. *Plast. Reconstr. Surg.* 112(6): 1604-12.

**Dong JS, Peng YP, Zhang YX, et al. (2003):** Reverse anterior tibial artery flap for reconstruction of foot donor site. *Plast Reconstr Surg* 112: 1604-12.

**Drake L, Vogl W and Mitchell A (2004):** Lower limb In: "Gray's Anatomy for students" Pp468-606. Edited by Drake L, Vogl W and Mitchell A., Churchill Livingstone, New York

**ElGhamry AH (2003):** Peroneal tendofascial flap: a new fascial flap for Achilles tendon coverage, a preliminary report. *British Journal of Plastic Surgery* 56: 284–288.

**Elghamry AH (2011):** The ankle extensor tendon sheath flap: a versatile fascial flap for coverage of the tibia and pretibial area: A preliminary report. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 64: 663-670.

## **References**

---

**El-Khatib HA (2007):** *The split peroneus muscle flap: a new flap for lower leg defects. Journal of Plastic, Reconstructive & Aesthetic Surgery* 60: 898-903.

**Eming SA, Krieg T, Davidson JM (2007):** *Gene therapy and wound healing. Clinics in Dermatology.* 25: 79–92.

**Fix RJ and Heinz TR (2010):** *Reconstruction of the Lower extremity. In: "Plastic Surgery Secrets". 2nd edn. Pp610-615. Edited by Weinzweig J, Mosby, Philadelphia.*

**Foo TL, Chew WY, Tan BK (2011):** *Improving the Reliability of the Distally Based Posterior Tibial Artery Adipofascial Flap With the Great Saphenous Venoneural Network. Ann Plast Surg* 67: 288–293.

**Galiano RD, Mustoe TA (2007):** *Wound care. In: "Grabb and Smith's Plastic Surgery". 6th edn., Pp23-32. Edited by Thorne CH, Lippincott Williams & Wilkins, Philadelphia.*

**Gonzalez MH, Tarandy DI, Troy D, et al (2002):** *Free tissue coverage of chronic traumatic wounds of the lower leg. Plast Reconstr Surg* 109: 592-600.

**Griffin JR and Thoronton JF (2003):** *Lower extremity reconstruction. Selected Readings in Plastic Surgery; 9(37): 1-44.*

**Haertsch PA (1981):** *The blood supply of the skin of the leg: a post mortem investigation. British Journal of Plastic Surgery.* 34: P.470.

**Hallock GG (2002):** *Sagittal Split Tibialis Anterior. Ann Plast Surg* 49: 39–43.

**Hansen SL and Mathes SJ (2006):** *Problem wounds and principles of closure. In: "Plastic Surgery". Pp901-50. Edited by Mathes SJ, Elsevier, New York.*

**Hansen T, Wikstrom J, Johansson LO, et al. (2007):** *The prevalence and quantification of atherosclerosis in an elderly population assessed by whole-body magnetic resonance angiography. Arterioscler Thromb Vasc Biol* 27:649-54.

**Hartmann A, Quist J, Hamm H, Bröcker EB, Friedl P (2008):** *Transplantation of autologous keratinocyte suspension in fibrin matrix*

## **References**

---

*to chronic venous leg ulcers: improved long-term healing after removal of the fibrin carrier. Dermatol Surg. Jul; 34(7):922-9.*

**Hass N, Kimmritz J, Witte W, Halle E (2002):** *Methicillin-resistant staphylococcus aureus (MRSA) sepsis following mesh graft transplantation to treat chronic leg ulcer. Clin Exp Dermatol; 27(4): 322-323.*

**Heitmann C, Khan FN, Levin LS (2003):** *Vasculature of the peroneal artery: an anatomic study focused on the perforator vessels. J. Reconstr. Microsurg. 19: 157-62.*

**Heymans O, Verhelle N, Peters S, Nélissen X, Oelbrandt B (2002):** *Use of the medial adipofascial flap of the leg for coverage of full-thickness burns exposing the tibial crest. Burns 28: 674–678.*

**Hirschowitz B, Moscona R, Kaufman T, Harshai Y (1987):** *External longitudinal splitting of the tibiallis anterior muscle for coverage of compound fractures of the middle third of the tibia. Plast. Reconstr. Surg. 79: 407-14.*

**Hollier L, Sharma S, Babigumira E, Klebuc M (2002):** *Versatility of the sural fasciocutaneous flap in the coverage of lower extremity wounds. Plast. Reconstr. Surg. 110: 1673–1679.*

**Hong JP, Shin HW, Kim JJ, et al. (2005):** *The use of anterolateral thigh perforator flaps in chronic osteomyelitis of the lower extremity. Plast. Reconstr. Surg. 115:142-7.*

**Hong JP, Shin HW, Kim JJ, et al. (2005):** *The use of anterolateral thigh perforator flaps in chronic osteomyelitis of the lower extremity. Plast Reconstr Surg 115: 142-7.*

**Hsieh CH, Liang CC, Kueh NS, Tsai HH, Jeng SF (2005):** *Distally based sural island flap for the reconstruction of a large soft tissue defect in an open tibial fracture with occluded anterior and posterior tibial arteries-a case report. Br. J. Plast. Surg. 58: 112–115.*

**Hueman MT, Thornton K, Herman JM, Ahuja N (2008):** *Management of Extremity Soft Tissue Sarcomas Surg Clin N Am 88: 539–557.*

**Hwang JH, Kim ES, Kim KS, Kim DY, Lee SY (2007):** *Latissimus Dorsi Muscle and Its Short Perforator-Based Skin Compound Free Flap.*

## **References**

---

*Annals of Plastic Surgery* 58(4): 381-7.

**Hyodo I, Nakayama B, Takahashi M, Toriyama K, Kamei Y, Torii S (2004):** *The gastrocnemius with soleus bi-muscle flap. The British journal of Plastic Surgery* 57: 77–82.

**Isenberg JS (2001):** *Additional follow-up with microvascular transfer in the treatment of chronic venous stasis ulcers. J Reconstr Microsurg;* 17(8): 603–605.

**Izadi K and Ganchi P (2005):** *Chronic Wounds. Clin. Plastic. Surg.* 32: 209–222.

**Jiga LP, Barac S, Taranu G, et al. (2010):** *The versatility of propeller flaps for lower limb reconstruction in patients with peripheral arterial obstructive disease: initial experience. Ann Plast Surg* 64:193-7.

**John Craven (2004):** *Nerves of the leg and foot. Anaesthesia & Intensive Care Medicine. Vol.5. Issue4: 112,113.*

**Kamath BJ, Joshua TV, Pramod S (2006):** *Perforator based flap coverage from the anterior and lateral compartment of the leg for medium sized traumatic pretibial soft tissue defects: a simple solution for a complex problem. J Plast Reconstr Aesthet Surg* 59: 515 -20.

**Kaplan M, Daly D, Stemkowski S (2009):** *Early intervention of negative pressure wound therapy using vacuum assisted closure in trauma patients: impact on hospital length of stay and cost. Adv Skin Wound Care* 22: 128-32.

**Kasabian AK and Karp NS (2007):** *Lower-Extremity Reconstruction. In: "Grabb And Smith's Plastic Surgery".6th edn., Pp676-688. Edited by Thorne CH., Lippincott Williams & Wilkins, Philadelphia.*

**Kawamura K, Yajima H, Kobata Y, et al (2005):** *Anatomy of Y-shaped configurations in the subscapular arterial system and clinical application to harvesting flow-through flaps. Plast Reconstr Surg.* 116: 1082–1089.

**Kim D, Orron DE, Skillman JJ (1989):** *Surgical significance of popliteal arterial variants. A unified angiographic classification. Ann. Surg.* 210(6): 776-81.

## **References**

---

**Kim HH, Jeong JH, Seul JH, et al. (2006):** New design and identification of the medial sural perforator flap. An anatomic study and clinical applications. *Plast Reconstr Surg* 117: 1609.

**Kim NG, Lee KS, Choi TH, Kim JS, Choi JH, Jang PY, Han KH, Son DG, Kim JH (2008):** Aesthetic reconstruction of lower leg defects using a new anterolateral lower leg perforator flap. *J. Plast. Reconstr. Aesthet. Surg.* 61(8): 934-8.

**Knight SL (2000):** Open tibial fractures: principles of soft-tissue cover. *Current Orthopaedics* 13: 92-98.

**Kobbe P, Lichte P and Pape FC (2009):** Complex extremity fractures following high energy injuries: The limited value of existing classifications and a proposal for a treatment guide. *Injury Int. J.* 40 Suppl. 4: S69-74.

**Koshima I, Fujitsu M, Ushio S, et al. (2005):** Flow-through anterior thigh flaps with a short pedicle for reconstruction of lower leg and foot defects. *Plast Reconstr Surg* 115 :155-62.

**Koshima I, Urushibara K, Inagawa K, et al. (2001):** Free medial plantar perforator flaps for the resurfacing of finger and foot defects. *Plast Reconstr Surg* 107: 1753-8.

**Koski EA, Kuokkanen HO, Tukiainen EJ (2005):** Distally-based peroneus brevis muscle flap: a successful way of reconstructing lateral soft tissue defects of the ankle. *Scand J Plast Reconstr Surg Hand Surg* 39: 299-301.

**Kovach SJ , Levin LS (2007):** Treatment of recalcitrant venous ulcers with free tissue transfer for limb salvage. In: "Venous Ulcers". Pp261-274. Edited by Bergan JJ, Shortell C, Elsevier. London..

**Kumins NH, Weinzweig N, Schuler JJ ( 2000):** Free tissue transfer provides durable treatment for large nonhealing venous ulcers. *J Vasc Surg.* 32: 848–854.

**Kuo YR, Jeng SF, Kuo MH, et al. (2001):** Free anterolateral thigh flap for extremity reconstruction: Clinical experience and functional assessment of donor site. *Plast Reconstr Surg* 107 :1766-71.



## **References**

---

**Lee CK and Hansen SL (2009):** Management of Acute wounds. *Surg. Clin. N. Am.* 89: 659–676.

**Lee HJ, Kim JW, Oh CW, et al. (2009):** Negative pressure wound therapy for soft tissue injuries around the foot and ankle. *J Orthop Surg Res* 9(4):14.

**Lin CH and Yazar S (2004):** Revisiting the serratus anterior rib flap for composite tibial defects. *Plast. Reconstr. Surg.* 114(7): 1871-1877.

**Lin CH, Wei FC (2000):** Widely split latissimus dorsi muscle flaps for reconstruction of long soft-tissue defects in lower extremities. *Plast Reconstr Surg* . 105: 706–709.

**Llanos S, Danilla S, Barraza C, et al. (2006):** Effectiveness of negative pressure closure in the integration of split thickness skin grafts: a randomized, double-masked, controlled trial. *Ann Surg* 244: 700-5.

**Lo LJ, Chen YR, Weng CJ, Noordhoff MS (1993):** Use of split anterior tibial muscle flap in treating avulsion injury of leg associated with tibia exposure. *Ann. Plast. Surg.* 31(2): 112-6.

**Lorenzetti F, Lazzeri D, Bonini L, Giannotti G, Piolanti N, Lisanti M, Pantaloni M (2010):** Distally based peroneus brevis muscle flap in reconstructive surgery of the lower leg: Postoperative ankle function and stability evaluation. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 63 1523-1533.

**Lu TC, Lin CH, Lin CH, Lin YT, Chen RF, Wei FC (2011):** Versatility of the pedicled peroneal artery perforator flaps for soft tissue coverage of the lower leg and foot defects. *Journal of Plastic, Reconstructive & Aesthetic Surgery.* 64: 386-393.

**Lu TC, Lin CH, Lin CH, Lin YT, Chen RF, Wei FC (2011):** Versatility of the pedicled peroneal artery perforator flaps for soft-tissue coverage of the lower leg and foot defects. *J Plast Reconstr Aesthet Surg.* 64(3): 386-93.

**Lutz BS, Wei FC, Machens HG, et al. (2000):** Indications and limitations of angiography before free-flap transplantation to the distal lower leg after trauma: prospective study in 36 patients. *J Reconstr Microsurg* 16: 187-91.

## **References**

---

**Marston W (2007):** *Evaluation and Treatment of Leg Ulcers Associated with Chronic Venous Insufficiency. Clin Plastic Surg 34: 717–730.*

**Masquelet AC and Sassu P (2009):** *Gastrocnemius flap In: "Flaps and reconstructive surgery". 1st edn., 409-421. Edited by Wei F and Mardini S., Elsevier, China.*

**Masquelet AC, Romana MC, Wolf G (1992):** *Skin island flaps supplied by the vascular axis of the sensitive superficial nerves: anatomic study and clinical experience in the leg. Plast. Reconstr. Surg. 89:1115–1121.*

**Mast BA (2001):** *Comparison of magnetic resonance angiography and digital subtraction angiography for visualization of lower extremity arteries. Ann Plast Surg 46:261- 4.*

**Mathes SJ, Nahai F (1997):** *Tibialis anterior flap. In: " Reconstructive Surgery. Principles, Anatomy, & Techniques". 1<sup>st</sup> edn. Pp1501–1512. Edited by Mathes SJ and Nahai F, Churchill Livingstone, New York.*

**Mehrotra S (2007):** *Perforator-plus flaps: a new concept in traditional flap design. Plast Reconstr Surg 119 :590-8.*

**Molnar JA, DeFranzo AJ, Hadaegh A, Morykwas MJ, Shen P, Argenta LC(2004):** *Acceleration of integrin corporation in complex tissue defects with subatmospheric pressure. Plast. Reconstr. Surg. 113: 1339–46.*

**Moore and Agur (2007):** *Lower Limb.In:"Essential Clinical Anatomy".3rd edn., P.319. Edited by Moore and Agur., Lippincott Williams & Wilkin, Philadelphia.*

**Moore KL and Dalley AF (2006):** *Lower limb.In:" Clinically oriented anatomy". 5th edn., P566. Edited by Moore KL and Dalley AF., Lippincott Williams & Wilkin, Philadelphia.*

**Morgan K, Brantigan CO, Field CJ, Paden M (2006):** *Reverse Sural Artery Flap for the Reconstruction of Chronic Lower Extremity Wounds in High-Risk Patients The Journal of Foot & Ankle Surgery 45(6): 417–423.*

**Mustoe TA, O'Shaughnessy K, Kloeters O (2006):** *Chronic wound pathogenesis and current treatment strategies: a unifying hypothesis. Plast Reconstr Surg; 117: 35S–41S.*

## **References**

---

**Nasir S, Aydin MA, Sonmez E, Baykal B (2010):** *Flow-through Free Latissimus Dorsi Flap for Reconstruction of Injured Limbs Evaluation of Hemodynamic Effects on Extremity Circulation* *Ann Plast Surg* 65: 164–169.

**Niranjan NS (2009):** *Posterior tibial artery perforator flap*. In: "Flaps and reconstructive surgery". 1st edn., Pp600-616. Edited by Wei F and Mardini S., Elsevier, China.

**Omar AA, Mavor AID, Jones AM, Homer-Vanniasinkam S (2004):** *Treatment of venous leg ulcers with Dermagraft®*. *Eur J Endovasc Surg*; 27: 666-72.

**Orgill DP, Manders EK, Sumpio BE, Lee RC, Attinger CE, Gurtner GC, Paul Ehrlich HP (2009):** *Clinical review: The mechanisms of action of vacuum assisted closure: More to learn*. *Surgery* 146:40-51.

**Ozkan O, Coskunfirat OK, Ozgentas HE (2004):** *An ideal and versatile material for soft-tissue coverage: experiences with most modifications of the anterolateral thigh flap*. *J Reconstr Microsurg*. 20: 377–383.

**Ozkan O, Ozgentas HE, Dikici MB (2005):** *Flow-through, functioning, free muscu-locutaneous flap transfer for restoration of a mangled extremity*. *J Reconstr Microsurg*. 21:167–172.

**Panagiotopoulos K, Soucacos PN, Korres DS, Petrocheilou G, Kalogeropoulos A, Panagiotopoulos E, Zoubos AB (2009):** *Anatomical study and colour Doppler assessment of the skin perforators of the anterior tibial artery and possible clinical applications*. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 62: 1524 :1529.

**Parrett BM, Matros E, Pribaz JJ, et al (2001):** *The parascapular free flap extended to the inframammary fold*. *Zhonghua Zheng. Xing Wai Ke Za Zhi* 17:339-40.

**Parrett BM, Matros E, Pribaz JJ, et al. (2006):** *Lower extremity trauma: trends in the management of soft-tissue reconstruction of open tibia-fibula fractures*. *Plast Reconstr Surg* 117: 1315-22.

**Patel NP, Labropoulos N, Pappas PJ (2006):** *Current management of venous ulceration*. *Plast Reconstr Surg*; 117 Suppl 7: 254S–260S.

## References

---

**Pinsolle V, Reau AF, Pelissier P, Martin D, Baudet J (2006):** Soft-tissue reconstruction of the distal lower leg and foot: are free flaps the only choice? Review of 215 cases. *Journal of Plastic, Reconstructive & Aesthetic Surgery*. 59: 912-917.

**Prantl L, Babilas P, Roll C, Jung M, Nerlich M, Kinner B, Schreml S, Fuechtmeier B (2009):** The use of 'fasciocutaneous' and 'osteofasciocutaneous' parascapular flaps for lower limb reconstruction: a retrospective study of 20 patients. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 62: 973-980.

**Pu L (2006):** Soft-Tissue Coverage of an Open Tibial Wound in the Junction of the Middle and Distal Thirds of the Leg With the Medial Hemisoleus Muscle Flap. *Ann Plast Surg* 56: 639–643.

**Pu L (2006):** The Reversed Medial Hemisoleus Muscle Flap and Its Role in Reconstruction of an Open Tibial Wound in the Lower Third of the Leg. *Ann Plast Surg* 56: 59–64.

**Pu LL (2005):** Successful soft-tissue coverage of a tibial wound in the distal third of the leg with a medial hemisoleus muscle flap. *Plast Reconstr Surg* . 115: 245–251.

**Rad AN, Christy MR, Rodriguez ED, Brazio P, Rosson GD (2011):** The Anterior Tibialis Artery Perforator (ATAP) Flap for Traumatic Knee and Patella Defects. *Ann Plast Surg* 64: 210–216.

**Raveendran SS, Perera D, Happuharachchi T, Yoganathan V (2004):** Superficial sural artery flap—a study in 40 cases. *Br J Plast Surg* 57(3): 266–269.

**Redett RJ, Robertson BC, Chang B, et al (2000):** Limb salvage of lower-extremity wounds using free gracilis muscle reconstruction. *Plast. Reconstr. Surg*. 106(7): 1507-1513.

**Rios-Luna A, Fahandezh-Saddi H, Villanueva-Martínez M, López AG (2008):** Pearls and tips in coverage of the tibia after a high energy trauma. *Indian J Orthop*. 42(4): 387-94.

## References

---

**Ríos-Luna A, Villanueva-Martínez M, Fahandezh-Saddi H, Villanueva-Lopez F, del Cerro-Gutiérrez M (2007):** Versatility of the sural fasciocutaneous flap in coverage defects of the lower limb injury. *Int. J.* 38: 824—831.

**Robotti E, Carminati M, Bonferraro PP, Bocchiotti MA, Ortelli L, Devalle L, and Righi B (2010):** “On Demand” Posterior Tibial Artery Perforator Flaps: A Versatile Surgical Procedure for Reconstruction of Soft Tissue Defects of the Leg After Tumor Excision. *Ann Plast Surg* 64: 202–209.

**Robson MC, Cooper DM, Aslam A, Gould LJ, Harding K G., Margolis DJ, Ochs DE, Serena TE, Snyder RJ, Steed DL, Thomas DR, Wiersma-Bryant (2006):** Guidelines for the treatment of venous ulcers. *Wound Rep Reg* 14: 649–662.

**Roccha JF, Gilbert A, Masquelet A, et al. (1987):** The anterior tibial artery flap: anatomic study and clinical application. *Plast. Reconstr. Surg.* 79(3): 396-406.

**Rogers LC, Bevilacqua NJ, Armstrong DG (2008):** The use of marrow-derived stem cells to accelerate healing in chronic wounds. *Int Wound J.* 5: 20–25.

**Sakai S, Shibata M (2006):** Free adductor perforator flap in lower leg reconstruction *Journal of Plastic, Reconstructive & Aesthetic Surgery* 59: 990-993.

**Sananpanich K, Tu YK, Kraissarin J, Chalidapong P (2008):** Flow-through anterolateral thigh flap for simultaneous soft tissue and long vascular gap reconstruction in extremity injuries: Anatomical study and case report *Injury, Int. J.* 39 S4: S47–S54.

**Shen JT, Falanga V. (2003):** Innovative therapies in wound healing. *J Cutan Med Surg.* 7(3): 217-24.

**Shi B, Chen SZ, Zhang P, Li JQ (2003):** Effects of vacuum-assisted closure (VAC) on the expressions of MMP-1, 2, 13 in human granulation wound [in Chinese]. *Zhonghua Zheng Xing Wai Ke Za Zhi;* 19:279-81.

## **References**

---

**Shim JS, Kim HH (2006):** A novel reconstruction technique for the knee. *Journal of Plastic, Reconstructive & Aesthetic Surgery* 59: 919-926.

**Sood R, Coleman JJ, Daising M, et al. (2000):** Preoperative evaluation of the lower extremity using color Doppler imaging versus angiography. Presented at American Society of Reconstructive Microsurgery, Miami.

**Stadler F, Brenner E, Todorroff B, et al. (1999):** Anatomical study of the perforating vessels of the lower leg. *Anat. Rec.* 255(4): 374-9.

**Tanaka M, Nakakita N, Kuroyanagi Y (1999):** Allogeneic cultured dermal substitute composed of spongy collagen containing fibroblasts: evaluation in animal test. *J Biomater Sci Polym Ed;* 10(4): 433–453.

**Taylor G (2007):** The blood supply of the skin. In: "Grabb And Smith's Plastic Surgery".6th edn., Pp33-41.. Edited by Thorne CH., Lippincott Williams & Wilkins, Philadelphia.

**Taylor GI, McCarten G, Doyle M (1990):** The use of the Doppler probe for planning flaps: anatomical study and clinical applications. *Br J Plast Surg.* 43: 1-16.

**Taylor GI, Palmer JH (1987):** The vascular territories (angiosomes) of the body:experimental study and clinical applications. *Br Plast Surg.* 40: 113-41.

**Thami GP, Singal A, Bhalla M (2004):** Full-thickness punch grafting in chronic nonhealing ulcers. *J Am Acad Dermatol;* 50: 99-100.

**Thione A, Valdatta L, Buoro M, et al. ( 2004):** The medial sural artery perforators anatomic basis for a surgical plan.. *Ann Plast Surg* 53:250-5.

**Top H, Benlier E, Aygit AC, Kiyak M (2005):** Distally-based sural flap in treatment of chronic venous ulcers. *Ann Plast Surg.* 55: 160 –165.

**Tos P, Innocenti M, Artiaco S, Antonini A, Delcroix L, Geuna S, Battiston B (2001):** Perforator-based propeller flaps treating loss of substance in the lower limb *J Orthopaed Traumatol* 12: 93–99

## **References**

---

**Tu YK, Yen CY, Ma CH, Yu SW, Chou YC, Lee MS, Ueng SW (2008):** *Soft-tissue injury management and flap reconstruction for mangled lower extremities Injury, Int. J. 39: 75- 95.*

**Tu YK, Yen YC (2007):** *The role of vascularized bone graft in lower extremity osteomyelitis. Orthop. Clinic. Of North. Am.38: 37-49.*

**Valencia IC, Falabella A, Kirsner RS, Eaglstein WH (2001):** *Chronic venous insufficiency and venous leg ulceration. J Am Acad Dermatol; 44(3): 401-421.*

**Villarreal PM, Monje F, Gañán Y, Junquera LM, Morillo AJ (2004):** *Vascularization of the peroneal muscles Critical evaluation in fibular free flap harvesting. Int. J. Oral Maxillofac. Surg. 33: 792-797.*

**Vuerstaek J, Vainas T, Neumann M (2007):** *Topical negative pressure techniques in chronic leg ulcers. In:"Venous ulcers".22edn. Pp307-326. Edited by Bergan JJ, Shortell CK, Elsevier Inc, London.*

**Weinzweig N, Babbitt R, Dunn RM (2010):** *Leg Ulcers. In:"Plastic surgery secrets". 2nd edn. Pp 616-625. Edited by Weinzweig J, Mosby, Philadelphia*

**Weissman IL (2000):** *Stem cells: units of development, units of regeneration, and units in evolution. Cell. 100:157 - 68.*

**Werner S, Grose R (2003):** *Regulation of wound healing by growth factors and cytokines. Physiol Rev. 83: 835-70.*

**Whetzel TP, Barnard MA, Stokesl RB (1997):** *Arterial fasciocutaneous vascular territories of the lower leg. Plast. Reconstr. Surg. 100(5): 1172-83.*

**Wong CH and Tan BK (2007):** *Intermittent short saphenous vein phlebotomy. An effective technique of relieving venous congestion in the distally based sural artery flap. Ann Plast Surg. 58: 303–307.*

**Wright JK, Watkins RP (1981):** *Use of the soleus muscle flap to cover part of the distal tibia. Plast. Reconstr. Surg. 68(6): 957-8.*

**Yang D and Morris SF (2002):** *Reversed sural island flap supplied by the lower septocutaneous perforator of the peroneal artery. Ann. Plast. Surg. 49: 375–378.*

## **References**

---

**Yang YL, Lin TM, Lee SS, et al. (2005):** *The distally pedicled peroneus brevis muscle flap anatomic studies and clinical applications. J Foot Ankle Surg 44: 259-63.*

**Yousif NJ (1996):** *The anterior tibial artery flap. In:" Atlas of microsurgical composite tissue transplantation".1<sup>st</sup> edn.,P459-72. Edited by Serafin D. WB Saunders, Philadelphia.*

**Zaulyanov L, Kirsner RS (2007):** *A review of a bi-layered living cell treatment (Apligraf®) in the treatment of venous leg ulcers and diabetic foot ulcers. Clinical Interventions in Aging; 2(1): 93–98.*

**Zielinski M, Pukacki F, Oszkinis G, Gabriel M, Waliszewski K, Majewski W (2008):** *The usefulness of topical negative pressure (TNP) as a supporting method for the surgical treatment of venous ulcers with the use of middle-split thickness skin graft. Polish Journal of Surgery. 80(3): 124-131.*

**Zweifel-Schlatter M, Haug M, Schaefer DJ, et al (2006):** *Free fasciocutaneous flaps in the treatment of chronic osteomyelitis of the tibia: a retrospective study. J Reconstr Microsurg 22: 41-7.*