

REFERENCES

- Abbott, W.W. (1925).
A method of computing the effectiveness of an insecticide.
J. Econ. Ent. 18:265-267.
- Ahmed, M. (1976).
Effect of phosphine fumigation on the germination of edible legume seeds.
J. Stored Prod. Res. 12:211-212.
- Aliniabee, M.T. (1971).
The effect of carbon dioxide gas alone or in combinations on the mortality of T. castaneum (Herbst) and T. confusum (Duval) Coleoptera, Tenebrionidae).
J. Stored Prod. Res. 7:243-252.
- Anon (1974).
Fumigant-Mode of action use and residue analysis.
J. Stored Prod. Res. 17:43-52.
- Bailey, S.W. (1955).
Air-tight storage of grain: its effects on insect pests. 1. Calandra granaria.
Aust. J. Agr. Res. 6:33-51.
- Bailey, S.W. (1956).
Air-tight storage of grain, its effect on insect pests. II. Calandra oryzae (small strain).
Aust. J. Agric. Res. 7:7-19.

Bailey, S.W. (1957).

Air-tight storage of grain, its effect on insect pests. III. Calandra oryzae (large strain).
Aust. J. Agric. Res. 8:595-603.

Banks, H.J. and Sharp, A.K. (1979).

Insect control with CO₂ in a small stock of bagged grain in a plastic film enclosure.
Aust. J. Exp. Agric. Anim. Husb. 19:102-107.

Barbara, D.H.; Bell, C.H.; Mills, K.A. and Goodship, G. (1976).

The toxicity of phosphine to all developmental stages of 13 species of stored product beetles.
J. Stored Prod. Res. 12:235-244.

Bell, C.H. (1984).

Effect of oxygen on the toxicity of carbon dioxide to stored insects.
Developments in Agric. Engineerings. Controlled atmosphere atmosphere and fumigation in grain storage, edited by Rippetal, B.E., 67-74.

Bell, C.H. (1976).

The tolerance of developmental stages of four stored product Moths to phosphine.
J. Stored Prod. Res. 12:77-86.

Bell, C.H. (1979).

The efficiency of phosphine against diapausing larvae of E. elutella, over a wide range of concentrations and exposure times.

J. Stored Prod. Res. 15:53-58.

Bell, C.H. and Glanville, V. (1973).

The effect of concentration and exposure in tests with methyl bromide and with phosphine on diapausing larvae of Ephestia elutella (Hubner) (Lepidoptera, Pyralidae).

J. Stored Prod. Res. 9:165-170.

Bell, C.H.; Hole, B.D. and Evans, P.H. (1977).

The occurrence of resistance to phosphine in adult and egg stages of R. dominica.

J. Stored Prod. Res. 13:91-94.

Beratliet, C. and Alexandrescu, S. (1964).

Effect of phosphine fumigation on the germination of wheat and maize seed.

Probleme Agric. 16:45-51.

Berck, B. (1968).

Sorption of phosphine by cereal products.

J. Agric. Fd. Chem. 16:419-425.

Bond, E.J. (1973).

Increased tolerance to ethylene dibromide in a field population of Tribolium castaneum (Herbst)

J. Stored Prod. Res. 9:61-63.

Bond, E.J. (1980).

Sorption of tritiated phosphine by various stages of T. castaneum.

J. Stord Prod. Res. 16:27-31.

Bond, E.J. and Buckland, C.T. (1978).

Control of insects with fumigants at low temperatures: toxicity of fumigants in atmosphere of CO₂.
J. Econ. Ent. 71:307-309.

Bond, E.J. and Buckland, C.T. (1979).

Development of resistance of carbon dioxide in the German weevil.
J. Econ. Ent. 72:770-771.

Bond, E.J. and Dumas, J. (1967).

Loss of warning odour from phosphine.
J. Stored Prod. Res. 3:389-392.

Bond, E.J. and Monro, H.A.U. (1961).

The toxicity of various fumigants to the cadelle Tenebroides mauritanicus.
J. Econ. Ent. 54:451-454.

Bond, E.J.; Monro, H.A.U. and Buckland, C.T. (1967).

The influence of oxygen on the toxicity of fumigants to Sitophilus granarius.
J. Stored. Prod. Res. 3:289-294.

Bond, E.J.; Robinson, J.R. and Buckland, C.T. (1969).

The toxic action of phosphine-absorption and symptoms of poisoning in insects.
J. Stored Prod. Res. 5:289-298.

Bruce, R.B.; Robbins, A.J. and Tuft, T.O. (1962).

Phosphine residues from phostoxin treated grain.
J. Agric. Fd. Chem. 10:18-21.

- Calderon, M. and Navarro, S. (1979).
Increased toxicity of low oxygen atmospheres supplemented with CO₂ on T. castaneum adults.
Ent. Exp. Appl. 25:39-44.
- Calderon, M. and Navarro, S. (1980).
Synergistic effect of CO₂ and O₂ mixtures on two stored grain insect pests.
Developments in Agric. Engineering I controlled atmosphere storage of grains, edited by Shejbal, J. pp. 79-84, Italy.
- Champ, B.R. and Dyte, C.E. (1976).
Report of the FAO global survey of pesticide susceptibility of stored grain pests.
FAO Plant Prod. Prot. Serv. No. 5.
- Childs, D.P. and Overby, J.E. (1983).
Mortality of the cigarette beetle in High-Carbon dioxide atmospheres.
J. Econ. Ent. 76:544-546.
- Corques, M.A. and Adeyemi, A.O. (1969).
A comparison of phosphine and ethylene dibrom for fumigation of cowpeas in polythene lined hessian sacks.
Bulletin of the Entomological Society of Nigeria, 2(1):45-50.
- Desmarchelier, J. M. (1984).
Effect of carbon dioxide on the efficacy of phosphine against different stored product insects
Mitteilung aus der Biologischen Bundesanstalt für land-und Forstwirtschaft-Berlin-Dahlem Heft 220 Juli, 1984.
- Dieterich, W.H.; Mayr, G. and Hild, K. (1967).
Hydrogen phosphide as a fumigant for foods, feeds and processed food products.
Residue Rev., 19:135-149.

El-Deeb, H.A.H. (1981).

Studies on some factors affecting the efficiency of atmospheric fumigation with methyl bromide and phosphine against the rice weevil, Sitophilus oryzae (L.) and the granary weevil, S. granarius (L.).

Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.

El-Lakwah, F.A. (1978).

The efficacy of combined fumigation using methyl bromide and phosphine in combatting khapra beetles (Trogoderma granarium (Everts)).

Federal Biological Establishment for Agriculture and Forestry, Institute of stored product protection, Berlin-Dahlem Translation No. 3-12 a.

El-Lakwah, F.A. (1990).

Fumigation experiments with phosphine in traditional mud silos in Egypt to control stored product insects.

Egypt. J. Appl. Sci., 5(3):54-65.

El-Lakwah, F.A.; Abdel-Gawaad, A.; Meuser, F.; Wohlgenuth, R. and Darwish, A. (1991).

Efficiency of phosphine alone and in mixtures with carbon dioxide against the adults of Tribolium castaneum and Sitophilus oryzae.

Egypt. J. Appl. Sci., 4(2):527-545.

El-Lakwah, F.A. and Khattab, M.M. (1989).

Effect of phosphine fumigation on germination of seeds.

J. Agric. Sci. Mansoura Univ. 14(2):794-796.

El-Lakwah, F.A.; Khattab, M.M.; Ahmed, S.M. and Adel-Latif, A.M. (1991).

Selection of the Red flour Beetle. (Tribolium castaneum (Herbst) for resistance to phosphine in the laboratory and Biological observation on the resistant strain.

Egypt J. Appl. Sci., 6(11):121-141.

- El-Lakwah, F.A.; khattab, M.M.; Ahmed, S.M. and Abdel-Latief, A.M. (1991).
Susceptibility of some field strains of Tribolium castaneum (Herbst) and Sitophilus oryzae (L.) collected from different Governorates in Egypt to phosphine.
Egypt. J. Appl. Sci., 6(11):142-159.
- El-Lakwah, F.A.; Kndil, M.M.; Khattab, M.M. and Halawa, A.Z. (1991).
Efficacy of phosphine to four species of mites, infesting stored products in Egypt.
Egypt. J. Appl. Sci., 6(10):51-68.
- El-Lakwah, F.A.; Kndil, M.M.; Khattab, M.M. and Halawa, A.Z. (1991).
Effect of the sublethal dose of phosphine on the biological characteristics of the survivors of the flour mite Acarus siro (L.).
Egypt. J. Appl. Sci., 6(10):285-291.
- El-Lakwah, F.M.; Meuser, F.; Abd-El-Gwaad, A.; Wohlgemuth, R. and Darwish, A. (1991).
Efficiency of phosphine alone and in mixtures with carbon dioxide against Angoumois Grain Moth Sitotroga cerealella (Olivier); (Gelechiidae, Lepidoptera).
J. of Plant Diseases and Protection 98(1):92-102.
- El-Lakwah, F.A.; Reichmuth, C.; Khattab, M.M. and Darwish, A.A. (1989).
Sorption of phosphine by larvae and pupae of Sitotroga cerealella (Olivier); Lepidoptera;
Egypt. J. Appl. Sci., 4(2):1-21.

- El-Lakwah, F.A.; Wohlgemuth, R. and Khattab, M.M. (1989).
Efficiency of phosphine and combination of phosphine with carbon dioxide against Khapra Beetle larvae Trogoderma granarium (Everts).
Anz.Schädlingsskde-pflanzenschutz, Umweltschutz. 62:85-88.
- El-Nahal, A.K.M.; Barakat, A.A.; El-Halafawy, M.A. and Hassan, H. (1986).
Susceptibility of natural strains of Sitophilus oryzae (L.), S. granarius (L.), Rhizopertha dominica (F.) and Tribolium castaneum (Herbst) collecting from the different governorates of Egypt to Methyl bromide and phosphine.
Bull. Entomol. Soc. Egypt. Econ. Ser. 5(14): 141-154.
- FAO/World Health Organization (1967).
Evaluation of some pesticide residues in food. Rome. FAO PL: CP/15, Geneva, WHO/Food Add. 67. 32. (C.F. Monro, 1969).
- Finney, D.J. (1971).
Probit Analysis, 3rd ed.
Cambridge University Press.
- Flury, F. and Zernik, F. (1931).
Schädliche Gase. Manual of fumigation for insect control.
Canada Department of Agric. London, Ontario.
- Girish, G.K. (1972).
Analysis of some important factors affecting the susceptibility of stored product insect pests to fumigants.
Bull. grain Tech. 10:197-210.
-

- Hole, B.D.; Bell, C.H. and Bowley, C.R. (1985).
The toxicity of methyl chloroform to stored product insects.
J. Stored prod. Res. 21(2):95-100.
- Hole, B.D.; Bell, C.H.; Mill, K.A. and Goodsh, G. (1976).
The toxicity of phosphine to all developmental stages of thirteen species of stored product beetles.
J. Stored Prod. res. 21:235-244.
- Howe, R.W. (1965).
Losses caused by insects and mites in stored foods and feeding stuffs.
Nutr. Abstr. Rev. 35:285-293.
- Howe, R.W. (1973).
The susceptipility of the immature and adult stages of S. granarius to phosphine.
J. Stored Prod. Res. 8:241-262.
- Irwin, D.G.; Smith, L.W. and Pratt, J.J. (1972).
Effects of CO₂ and N₂ on the secretion of para-benzoquinones by T. castaneum.
J. Stored Prod. Res. 8:213-219.
- Jalil, M.; Ross, I.J. and Rodriguez, J.G. (1970).
Methyl bromide and phosphine as fumigants for some Acarid mites.
J. Stored Prod. Res. 6:33-37.
- Jay, E.G. (1980).
Methods of applying carbon dioxide for insect control in stored grain.
United States Department of Agric. (AAT-5-13):
7 pp.

- Jay, E.G. and Pearman, G.C. (1973).
CO₂ for control of an insect infestation in stored corn .
J. Stored Prod. Res. 9:25-29.
- Jay, E.G.; Redlinger, L.M. and Laudani, H. (1970).
The application and distribution of CO₂ in a peanut silo for insect control.
J. Stored Prod. Res. 6:247-254.
- Kashi, K.P. (1982).
Dose mortality responses of 5 species of stored products insects to phosphine.
Int. Pest Control 24(2):46-49.
- Kashi, K.P. and Bond, E.J. (1975).
The toxic action of phosphine: Role of CO₂ on the toxicity of phosphine to S. granarius and T. confusum.
J. Stored Prod. Res. 11:9-15.
- Kranz, J.; Schmutterer, H. and Koch, W. (1977).
Diseases, pests and weeds in Tropical crops.
Verlag Paul Parey. Berlin and Hamburg 666 pages.
- Krishnamurthy, T.S.; Spratt, E.C. and Bell, C.H. (1986).
The toxicity of carbon dioxide to adult beetles in low oxygen atmospheres.
J. Stored Prod. Res. 22:145-151.
- Kumar, A.; Srivastava, J.L.; Pandey, G.P. and Varma, B.K. (1981).
Phosphine residue in multi fumigated rice stocks stored in commercial warehouses.
Bull. Grain Tech., 19:81-85.

- Lindgren, D.L. and Vincent, L.E. (1959).
Biology and control of Trogoderma granarium
Everts.
J. Econ. Ent., 52(2):312-319.
- Lindgren, D.L. and Vincent, L.E. (1966).
Relative toxicity of hydrogen phosphide to various
stored-product insects.
J. Stored Prod. Res., 2:141-146.
- Lindgren, D.L. and Vincent, L.E. (1970).
Effect of atmospheric gases alone or in combinat-
ion on the mortality of granary and rice weevils.
J. Econ. Ent. 63:1926-1929.
- Lindgren, D.L.; Vincent, L.E. and Strong, R.G. (1958).
Studies on hydrogen phosphide as a fumigant.
J. Econ. Ent. 51:900-903.
- Lindgren, D.L. and Vincent, L.E. (1960).
The relation of moisture content and temperature
of stored grain to the effectiveness of grain
fumigants under forced circulation.
J. Econ. Ent. 53:1071-1077.
- Liscombe, E.A.R. (1963).
Hydrogen phosphide in tablet form as a grain
fumigant.
Research for Farmers, 8:6-7.
- Mayr, G. and Hild, K. (1966).
Unpublished data quoted by Dietrich et al. (1967).
- Monro, H.A.U. (1969).
Manual of fumigation for insect control.
F.A.O. Agricultural studies no. 79:381 pp.

- Monro, H.A.U.; Musgrave, A.J. and Upitis, E. (1961).
Induced tolerance of stored-product beetles to methyl bromide.
Ann. Appl. Biol. 49:373-377.
- Murthy, K.S.R.K. and Srivastava, B.P. (1971).
Effect of food on the susceptibility of pulse beetle, Callosobruchus maculatus (Fab) (Bruchidae: Coleoptera) to different fumigants.
Indian J. of Entomol., 33(2):148-151.
- Nakakita, H. and Winks, R.G. (1981).
Phosphine resistance in immature stages of a laboratory selected strain of T. castaneum.
J. Stored Prod. Res. 17:43-52.
- Nakakita, H.; Saito, T. and Iyatomi, K. (1974).
Effect of phosphine on the respiration of adult S. zeamais.
J. Stored Prod. Res. 10:87-92.
- Navarro, S. and Calderon, M. (1974).
Exposure of Ephestia cautella pupae to CO₂ concentrations at different relative humidities the effect on adult emergence and loss in weight.
J. Stored Prod. Res. 10: 237-241.
- Navarro, S.; Dias, R. and Donahage, E. (1985).
Induced tolerance of S. oryzae adults to carbon dioxide.
J. Stored Prod. Res. 21:207-213.
- Neitzert, K. (1953).
Influence on the baking quality of wheat of phostoxin fumigation.
Published report to Degesch Co., Frankfurt-Am. Main, Germany.

- Noack, S. and Reichmuth, Ch. (1978).
Einrechnerisches Verfahren zur Bestimmung von beliebigen Dosis-Werten eines Wirkstoffes aus empirisch Dosis-Wirkungs-Daten-Mitt.
Biol. Bundesanstalt für Land-u. Forstwirtschaft., Berlin-Dahlem, Heft 185:1-49.
- Noack, S. Wohlgemuth, R. (1985).
PH₃-Rückstände in Haselnüssen, Sojabohnen und Weizen nach Phosphine-Begasungen mit zeitlich nicht konstanter Konzentration.
Z. Lebensm. Unters. Forsch. 180:101-108.
- Noack, S.; Reichmuth, Ch. and Wohlgemuth, R. (1983).
PH₃-Rückstände bei Vorratsschutzbegasungen in Abhängigkeit von der Konzentration, Einwirkungszeit und Lagerdauer nach der Begasung.
Zeitschrift für Lebensmittel-Untersuchung und Forschung. 177:87-93.
- Noack, S.; Reichmuth, Ch. and Wohlgemuth, R. (1984).
Rückstandsverhalten von Phosphine in begasteten Lebensmitteln in Abhängigkeit von der Lagertemperatur und der Belüftung.
Zeitschrift Lebensmittel-Untersuchung und Forschung. 178:31-37.
- Ozar, A.I. (1974).
Investigation on the effect of phosphine on Bruchus lentis Frohl and Callosobruchus maculatus F.
Bitki Karuma Bulteni, 14(1):43-54.

- Palaniswamy, P.T. and Dakshinamurthy, A. (1986).
Biogas to control rice storage pests India.
International Rice Research Newsletter, N. 11(6):
25.
- Pinto de Matos, A.H. (1961).
Influence of insecticide treatment on the germination of groundnut seeds.
Garcia de Orta, 9:471-477. (C.F. Monro, 1969).
- Price, N.R. (1981).
A comparison of the uptake and Metablism of ³²P-Radiolabelled Phosphine in susceptible and resistant strains of the lesser grain borer (Rhizopertha dominica).
Comp. Biochem. Physiol. C; 69(1):129-131.
- Price, N.R. (1985).
Review, the mode action of fumigants.
J. Stored Prod. Res. 21:157-164.
- Price, N.R. and Dance, S.J. (1983).
Some biochemical aspects of phosphine action and resistance in 3 species of stored product beetles.
Comp. Biochem. Physiol. C. Comp. Pharmacol. Toxicol. 76(2):277-282.
- Quereshi, A.H.; Bond, E.J. and Monro, H.A.U. (1965).
Toxicity of hydrogen phosphide to the granary weevil S. granarius and other insects.
J. Econ. Ent. 58:324-331.

Rai, L.; Sarid, J.N. and Pingale, S.V. (1962).

Fumigation of food grains in India with hydrogen phosphide. I. Tests in concrete bins.

Bull. Grain Tech. 1:3-17.

Reichmuth, Ch. (1984).

The effect of the fumigant hydrogen phosphide (phosphine) on stored product pests.

German Biological Research Center/Institute for Stored Products Protection.

Reichmuth, Ch. (1986).

The significance of changing concentrations in toxicity of phosphine. "GASGA seminar on fumigation technology in developing countries".

Tropical development and research institute, storage department, Slough, 18-21 March 1986, 88-98.

Reynolds, E.M.; robinson, J.M. and Howells, C. (1967).

The effect on Sitophilus granarius (L.) (Coleoptera, Curculionidae) of exposure to low concentrations of phosphine.

J. Stored Prod. Res. 2:177-186.

Robinson, J.R. and Bond, E.J. (1970).

The toxic action of phosphine Studies with $^{32}\text{PH}_3$, terminal residues in biological materials.

J. Stored Prod. Res. 6:133-146.

- Singh, K.N. and Srivastava, B.P. (1980).
Studies of the efficacy and extent of residues
of phosphine in stored pulses.
Pesticides 14(2):32.
- Singh, K.N. and Srivastava, B.P. (1983).
Relative toxicity of various fumigants to the
adults of two species of *Callosbruchus* and the
susceptibility of different sexes to these fumig-
ants.
Indian J. Entomol., 45(2):130-138.
- Singh, K.N.; Srivastava, B.P. and Nath, G. (1983).
Phosphine residues and effect on the germination
of stored pulses.
Indian J. Entomol. 45:71-80.
- Sinha, R.N. and Watters, F.L. (1985).
Insect pests of flour mills, grain elevators,
and feed mills and their control.
Research Branch Agriculture Canada, Publication
1776 E, Cat. No. A 43-1776/1985 E.
- Sinha, R.N.; Berck, B. and Wallace, A.H. (1966).
Effect of phosphine on mites, insect, and micro-
organisms.
Contribution no. 221 Canada Dept. of Agric. Rec.
Station, Winnipeg Accepted for Publication August
29, 1966.
- Sinha, R.N.; Berck, B. and Wallace, A.H. (1967).
Effect of phosphine on mites, insects, and micro-
organisms.
J. Econ. Ent. 60:125-132.

Sowunmi, E.O. (1982).

Effect of storage length and insecticide treatment on cowpea (Vigna unguiculata L. Walp).

Tropical Grain Legume Bulletin 25:21-27.

Spratt, E.C. (1975).

Some effects of the CO₂ absorbency of humidity controlling solutions on the results of life history experiments with stored products insects. J. Stored prod. Res. 11:127-134.

Spratt, E.C. (1979a).

Some effects of a mixture of O₂, CO₂ and N₂ in the ratio 1 : 1 : 8 on the oviposition and development of S. zeamais.

J. Stored Prod. Res. 15:73-80.

Spratt, E.C. (1979b).

The effects of a mixture of O₂, CO₂ and N₂ in the ratio 1 : 1 : 8 on the longevity and the rate of increase of populations of S. zeamais.

J. Stored Prod. Res. 15:81-85.

Srivastava, J.L. (1985).

Tamil NADU Agric. Univ., Coimbatore (India), pp. 205-207.

Storey, C.L. (1978).

Mortality of cowpea weevil in a low-oxygen atmosphere.

J. Econ. Ent. 71(5):833-834.

Storey, C.L.; Sauer, D.B. and Wlker, D. (1984).

Present use of pest management practices in wheat corn and date stored on the farm.

J. Econ. Ent. 77(3):784-788.

- Stoyanova, S. and Shikrenov, D. (1976).
Storage of cereals in an atmosphere with a high CO₂ concentration. Effect of 20 and 40% CO₂ on insect pests.
Development in Agric. Engineering. I Controlled atmosphere storage of grains, edited by Shejbal, J. pp. 101-118 Italy.
- Strong, R.G. and Lindgren, D.L. (1960).
Germination of small legume seeds after fumigation with hydrogen phosphide.
J. Econ. Ent. 53:1-4.
- Tunc, I. (1983).
Mortality of T. confusum adults in various atmospheric gas compositions.
Zeitschrift für angewandte Entomologie. Sonderdruck aus Bd. 95, H. 3, S. 263-267.
- Tyler, P.S.; Taylor, R.W. and Rees, D.P. (1983).
Insect resistance to phosphine fumigation in food warehouses in Bangladesh.
Int. Pest control 25:10-13.
- Upitis, E.; Monro, H.A.U. and Bond, E.J. (1973).
Some aspects of inheritance of tolerance to methyl bromide by S. granarius.
J. Stored Prod. Res. 9:13-17.
- Van den Bruel, W.E. and Bollaerts, D. (1956).
The hydrogen phosphide fumigation, a new method with a west range of application.
- Verma, A.K. and Wahdi, S.R. (1978).
Susceptibility of walnut pests to CO₂ and N₂ and effect of gas storage on keeping quality of walnut kernels.
Indian J. Ent. 40:290-298.

- Vincent, L.E. and Lindgren, D.L. (1971).
Comparison of the sorption of hydrogen phosphide, methyl bromide, ethylene dibromide and hydrocyanic acid by wheat and corn of different moisture contents and load factors.
J. Econ. Ent. 64:122-123.
- Vincent, L.E. and Lindgren, D.L. (1972).
Toxicity of phosphine to the life stages of four species of dermestidae.
J. Econ. Ent. 65:1429-1431.
- Vincent, L.E. and Lindgren, D.L. (1975).
Toxicity of phosphine and methyl-bromide at various temperatures and exposure periods to the four metamorphic stages of Trogoderma variable
J. Econ. Ent., 68(1):53-56.
- Vincent, L.E. and Lindgren, D.L. (1977).
Toxicity of methyl bromide and phosphine at various temperatures and exposure periods to the metamorphic stages of Lasioderma serricorne.
J. Econ. Ent., 70(4):497-500.
- Wigglesworth, V.B. (1965).
The principles of insect physiology.
Menthuen, London (C.F. Bond and Monro, 1967).
- Williams, P. Minett, W.; Buchana, S.A.; Wilson, S.D.; Savage, P.; Rosenbaum, H. and Guiffre, V. (1985).
Assessment of the commercial application of a methyl bromide and carbon dioxide mixture for the control of grain insects.
Plant Prot. q. 1(2):51-56.

Winks, R.G. (1969).

Resistance to the fumigant phosphine in a strain of T. castaneum.
Insect Tox. Inf. Serv. 12:178-180.

Winks, R.G. (1985).

The toxicity of phosphine to adults of T. castaneum phosphide-induced narcosis.
J. Stored Prod. Res. 21:25-29.

Winks, R.G. (1986).

Strategies for effective use of phosphine as a grain fumigant and the implications of resistance.
Proc. 4th Int-Work. conf. Stored- Product TelAviv, Israel Sept. 1986 [Eds. E. Donahaye and S. Navaro]. pp. 335-344.

Yadav, T.D. and Singh, S. (1977):

Respiratory behaviour of developing stages of Callosbruchus maculatus (Fab) and C. chinensis (Linn) under airtightness.
Bulletin of Grain Technology 15(1):67-70.