

REFERENCES

- Alexander, M.** (1977). Introduction to soil microbiology , 2nd ed. John Wiley & Sons, Inc. , New York
- Alexander , M.** (1994). Biodegradation and Bioremediation. Academic Press , Inc. New York
- Anderson, T. A.** (1992). Comparative plant uptake and microbial degradation of Trichloroethylene in the rhizosphere of five plant species, ORNL/ TM- 12017, Oak Ridge National Laboratory, Oak Ridge, T N
- Anderson, T. A. , E.A. Guthrie, and B. T. Walton** (1993). Bioremediation in the rhizosphere. Environ. Sci. Technol. 27, 2630-2636
- Anderson, T. A., E.L. Kruger, and J.R. Coats** (1994). Biological degradation of pesticide wastes in the root zone of soils collected at an agrochemical dealership. In: Bioremediation through rhizosphere technology ; Anderson, T. and J.R.Coats. (eds), American Chemical Society, Washington. DC 1994
- A.O.A.C.** (1970). Offical Methods of Analysis . Assoc. Offical Analytical Chemists, Washington, D.C.

Audus, L.J. (1964). In the Physiology and Biochemistry of Herbicides. (Audus, L.J. ed.). pp 163-206. Academic Press. London

Baird, D.D., R.P. Upchurch, W.B. Homesly, and J.E. Franz (1971). Introduction of a new broad spectrum postemergence herbicide class with utility for herbaceous perennial weed control. Proceedings of the North Central Weed Control Conference, 26, 64-68

Bell, R. M. (1992). Behaviour of pollutants in the plant-soil environment. In: Higher plant accumulation of organic pollutants from soils. Report No. EPA/ 600 / R- 92 / 138 , 1992

Berenson, M. L., D. M. Levine, and D. Rindskopf (1988). Applied statistics. A first course, Prentice Hall, Englewood Cliffs, New Jersey 07632

Bollag, J.M., G.G. Briggs, J.E. Dawson, and M. Alexander (1968). 2,4-D Metabolism. Enzymatic degradation of chlorocatechols. J. Agric. Food Chem. 16, 829-833

Boulter, D., J.J. Jeremy, and M. Wilding (1966). Amino acids liberated into the culture medium by pea seedling roots. Plant and Soil 24, 121-127

Cheshire, M.V. and C. Mundie (1990). Organic matter contributed to soil by plant roots during the growth and decomposition of maize. *Plant and Soil*, 121, 107-114

Chiou, C.T., V.H. Freed, D.W. Schmedding, and R.L. Kohnert (1977). Partition Coefficient and bioaccumulation of selected organic chemicals. *Environ. Sci. and Technol.* 11(5), 475-478

Colab, T., W.A. Althaus, and H.L. Wooten (1979) Fate of ^{14}C Trifluralin in soil. *J. Agric. Food Chem.* 27, 163-179

Cole, D.J. (1985). Mode of action of glyphosate- A literature analysis. In: Grossbard, E. and D. Atkinson (eds). *The herbicide glyphosate*. Butterworths, London

Crafts, A.S. (1964) Herbicide behaviour in the plant. In: *The physiology and biochemistry of herbicides*, L.J. Audus (ed.), Academic Press, London & New York, pp 75-110

Cullimore, D.R. (1981). The enumeration of 2,4-D degraders in Saskatchewan soils. *Weed Sci.* 29, 440-443

Cunningham, S. and W. Berti (1993). Remediation of contaminated soils with green plants: An overview . In Vitro Cell. Dev. Biol. 29, 207-212

Curl, E.A., and B. Truelove (1986). The Rhizosphere. Spring-Verlag. Berlin, 288 p.(1986)

Devine, M., S.O. Duke, and C. Fedtke (1993). Physiology of herbicide action. PTR, Prentice Hall, Englewood Cliffs, New Jersey 07632 (1993)

Doxtader, K.G. and Saleh-Rastine, N. (1979). Population dynamics of microorganisms capable of degrading 2,4-D. In Agronomy Abstract, Proceedings of ASA, CSSA, and SSSA Annual Meetings, p 127

Foth, H. D. (1990) Fundamentals of Soil Science, 8th ed. John & Wiley sons Inc., New York.

Fournier, J.C., P. Codaccioni, and G. Soulas (1981). Soil adaptation to 2,4-D degradation in relation to the application rates and the metabolic behaviour of the degrading microflora. Chemosphere 10, 977-984

Franz, Z. J. E. (1985). Discovery, development, and chemistry of glyphosate. In: Grossbard, E., and D.

Atkinson (eds). The Herbicide Glyphosate. Butterworths, London

Freed, V.H., and M.L. Montgomery (1963). The metabolism of herbicides by plants and soils. Residue Rev. 3, 1-17

Grover, R. (1973). The adsorptive behaviour of acid and ester forms of 2,4-D on soils. Weed. Res. 13, 51-58

Hance, R. J. (1976). Adsorption of glyphosate by soils. Pesticide Science, 7, 363-366

Hanne, L.F., L.L. Kirk, S.M. Apple, A.D. Narayan, and K.K. Bains (1993). Degradation and induction specificity in actinomycetes that degrade p-nitrophenol. Appl. Environ. Microbiol. 59, 3505-3508

Harms, H. and C. Langebartels (1986). Standardized cell suspension test systems for an ecotoxicologic evaluation of the metabolic fate of xenobiotics. Plant Sci. 45, 1-9

Hartter, D.R. (1985). The use and importance of nitroaromatic chemicals in the chemical industry. In: Richtert, D.E. (ed). Toxicity of nitro aromatic compounds. Chemical Industry Institute of Toxicology Series. Hemisphere, Washington.D.C. pp 1-14

Hogland, D.R., and D.I. Arnon (1950) The water culture method for growing plants without soil, p. 32. Circular 347, California Agr. Expt. Station, Univ. of California, Berkeley

Horvath, R.S. (1972). Microbial co-metabolism and the degradation of organic compounds in nature. Bacteriol. Reviews. 36, 146-155

Hsu, T. S. and R. Bartha (1979). Accelerated mineralization of two organophosphorous insecticides. Appl. Environ. Microbiol. 37, 36-41

Husain, S.S. and W.E. McKeen (1963). Interaction between strawberry roots and Rhizoctonia gragariae. Phytopathology, 53, 541-545

Jvanov, V.P., G.A.Y. Jacobsen, and B.S. Fomenko (1964). Effects of soil moisture on metabolism of root excretions. Fiziol. Rast. 11, 630-637

Katznelson, H., J.W. Rouatt, and T.M.B. Payne (1955). The liberation of amino acids and reducing compounds by plant roots. Plant and Soil 7, 35-48

Keith, L.H., and W.A. Telliard (1979). Priority pollutants. A perspective view. Environ. Sci. Technol. 13, 416-423

Kerr, A. (1964). The influence of soil moisture on infection of peas by Pythium ultimum. Aust. J. Biol. Sci. 17, 676-685

Kinouchi.T., and Y. Ohnishi (1983). Purification and characterization of 1-nitropyrene nitroreductases from *Bacteriodes fragilis*. *Appl. Environ. Microbiol.* 45, 1234-1241

Knaebel,D.B., and J.R. Vestal (1994). Intact rhizosphere microbial communities used to study microbial biodegradation in agricultural and natural soils. In: *Bioremediation through rhizosphere technology*, Anderson, T.A. and J.R. Coats (eds); American Chemical Society, Washington, D.C.

Kunc,F., and J. Rybarova (1983a). Mineralization of carbon atoms of ^{14}C - 2,4-D side chain and degradation ability of bacteria in soil. *Soil Biol Biochem.* 15, 141-144

Laanio,T.L., P.C. Kearney, and D.D. Kaufman (1973). Microbial metabolism of dinitramine. *Pestic. Biochem. Physiol.* 3, 271-277

Lichtenstein,E.P.,and K.R. Schulz (1964). The effect of moisture and microorganisms on the persistence and metabolism of some organophosphorus insecticides in soils, with special emphasis on parathion. *J. Environ. Entomol.* 57, 618-627

Liu, C. M., P.A. Mclean, C.C. Sookdeo, and F.C. Cannon (1991). Degradation of the herbicide glyphosate by members of the family Rhizobiacea. Appl. Environ. Microbiol. 57, 1799

Loos, M.A. (1975a). Phenoxyalkanoic acids. In: Kearney, P.C. and D.D. Kaufman (eds), Herbicides: chemistry, degradation, and mode of action. vol. 1. Marcel Dekker, New York, pp 1- 128

Loos, M.A., I.F. Schlosser, and W.R. Mapham (1979). Phenoxy herbicides degradation in soils. Quantative studies of 2,4-D and MCPA- degrading microbial populations. Soil Biol. biochen. 11,377-385

Lyayi, C.B. (1986) The metabolism of p-nitrophenol by Penicillium Chrysogenum. Microbios 45, 153-159

Markle, R.A., A.F. Fentiman, J.R. Steadman, and R.A. Mayer (1980). Potentially toxic and hazardous substances in the industrial organic chemicals and organic dyes and pigment industries. EPA 600/2-80-056. National Technical Information Service, Springfield, Va.

Matsumoto, H., K.Okada, and E. Takahasi (1979). Excretion products of maize roots from seedling to seed development stage. Plant and Soil 53, 17-26

Masse, R., M. Badr, C. Ayotte, and M. Sylvestre (1985).
Gas chromatographic- Mass spectrometric
characterization of bacterial metabolites of 4-
nitrophenol formed in gram negative strain B206.
Toxicol. Environ. Chem. 10, 225-246

McCall, P. J., S. A. Vrona, and S. S. Kelly (1981). Fate of
uniformly carbon-14-ring labeled 2,4-5-
trichlorophenoxyacetic acid and 2,4-
dichlorophenoxyacetic acid. J. Agric. Food Chem. 29, 100-
107

McCarty, P. L., B. E. Rittmann, and E. J. Bouwer (1984).
Microbiological processes affecting chemical
transformation in groundwater, In: Groundwater
Pollution Microbiology, Bitton. B. and Gebra, P. G.,
(eds.), Wiley, New York, 1984, 89

Merckx, R., A. Dijkstra, A. D. Hartog, and J. A. Van Veen
(1987). Production of root-derived material and
associated microbial growth in soil at different
nutrient levels. Biology and Fertility of Soils. 5, 126-
132

Moore, F. K., H. D. Braymer, and A. D. Larson (1983).
Isolation of a *Pseudomonas* sp. which utilizes the

phosphonate herbicide glyphosate. Appl. Environ. Microbiol. 46, 316-320

Moreale, A., and R. Van Bladel (1980), Behaviour of 2,4-D in Belgian soils. J. Environ. Qual. 9, 627- 633

Moshier, L. J., and D. Penner (1978). Factors influencing microbial degradation of ^{14}C glyphosate to ^{14}C CO_2 in soil. Weed Sci. 26, 686-691

Munnecke, D.M., and D.P.H. Hsieh (1974). Microbial decontamination of parathion and p-nitrophenol in aqueous media. Appl. Microbiol. 28, 212-217

Munnecke, D.M. (1976). Enzymatic hydrolysis of organophosphatic insecticides, a possible pesticide disposal method. Appl. Environ. Microbiol. 32, 7-13

Nishino, S.F. and J.C. Spain (1993). Cell density-dependent adaptation of *Pseudomonas putida* to biodegradation of p-nitrophenol. Environ. Sci. Technol. 27, 489-494

Nomura, N.S. and H.W. Hilton (1977), The degradation of glyphosate in Hawaiian sugar cane soils. Weed Research, 17, 113-121

Ogram, A.V., R.E. Jessup, L.T. Ou, and P.S.C. Rao (1985). Effects of 2,4-dichlorophenoxyacetic acid in soils. Appl. Environ. Microbiol. 49, 582-587

Ou, L.T., D.F. Rothwell, W.B. Wheller, and J.M. Davidson (1978b). The effects of high 2,4-D concentrations on degradation and carbon dioxide evolution in soils. J. ..Environ. Qual. 7, 241-246.

Ou, L.T. (1984). 2,4-D degradation and 2,4-D degrading microorganisms in soils. Soil Sci. 137, 100-107

Parker, L.W. (1979). The kinetics of the microbial decomposition of 2,4-D in soil. Ph.D thesis. Colorado State University, Fort Collins

Parker, L.W. , K.G.D. Doxtader (1982). Kinetics of microbial decomposition of 2,4-D in soil : Effect of herbicide concentration. J. Environ.Qual. 11, 679-684

Parker, L.W. and K.G. Doxtader (1983). Kinetics of the microbial degradation of 2,4-D in soil: Effects of temperature and moisture. J.Environ. Qual. 12, 553-558

Quilty, S.P. and M.J. Geoghegan (1976). The degradation of glyphosate in peat. Proceedings of the society for general microbiology. 3, 129

Raymond, D. G. M., and M. Alexander (1971). Microbial metabolism and cometabolism of nitrophenols. Pest. Biochem. Physiol. 1, 123-130

Reddy, B. R. and N. Sethunathan (1983a), Mineralization of parathion in the rice rhizosphere. Appl. Environ. Microbiol. 45, 826-829

Reddy, B.R. and N. Sethunathan (1983b). Mineralization of parathion in the rhizosphere of rice and pearl millet. J. Agric. Food Chem., 31, 1379-1381

Reddy, B.R. and N. Sethunathan (1994). Mineralization of p-nitrophenol in the rhizosphere of rice. Agriculture, Ecosystems and Environment. 47,313-317

Rochkind, M.L., J.W. Blackburn, and G.S. Sayler (1986). Microbial decomposition of chlorinated aromatic hydrocarbons. EPA /600/2-86/090

Rovira,A.D. (1956). Plant root excretions in relation to the rhizosphere effect. 1- The nature of root exudates from oats and peas. Plant and Soil, 7, 178-194

Rovira,A.D. (1959). Plant root excretion in relation to the rhizosphere effect.

IV- Influence of plant species, age of plant, light, temperature and calcium nutrition on exudation. Plant and Soil 11, 53-64

Rovira, A.D. (1969). Plant root exudates. The Botanical Review. 35, 35-37

Rovira, A.D., and C.B. Davy (1971). Biology of the Rhizosphere. In: The plant root and its environment, Carson, E.W. (ed.) 153-203

Rubin, H.E., and M. Alexander (1983). Effect of nutrients on the rates of mineralization of trace concentrations of phenol and p-nitrophenol. Environ. Sci. Technol. 17, 104

Rueppel, M., B. Brightwell, J. Schaefer, and J. Marvel (1977). Metabolism and degradation of glyphosate in soil. J. Agric. Food Chem. 25, 517-528

Sandmann, E. R.I.C., and M.A. Loos (1984). Enumeration of 2,4-D degrading microorganisms in soils and crop plant rhizosphere using indicator media; high populations associated with sugar cane (*Saccharum Officinarum*). Chemosphere 13, 1073-1084

Sandmann, E.R.I.C., M.A. Loos , and L.P. Van Dyk (1988).
The microbial degradation of 2,4-dichlorophenoxyacetic acid in soil. Review of Environmental Contamination and Toxicology. 101,1-53

Sethunathan, N., and T. Yoshida (1972). Conversion of parathion to p-nitrophenol by diazinon-degrading *Flavobacterium* sp. Proc. Inst. Environ. Sci. N.Y. Meet. 18 , 255-357

Shann, J.R., and J.J. Boyle (1994). Influence of plant species on In Situ rhizosphere degradation. In: Bioremediation through rhizosphere technology, Anderson, T.A., and J.R. Coats (eds), American Chemical Society, Washington, DC 1994

Sharmila, M., K. Ramanand, and N. Sethunathan (1989). Effect of yeast extract on the degradation of organophosphorus insecticides by soil enrichment and bacterial cultures. Can. J. Microbiol. 35, 1105-1110

Shimp, J.F., J.C. Tracy, L.C. Davis, E. Lee, W. Huang, and L.E. Erickson (1993). Beneficial effects of plants in the remediation of soil and groundwater contaminated with organic materials. Crit. Rev. in Environmental Science and Technology 23 (1) 41-77

Simpson, J.R. and W.C. Evans (1953). The metabolism of nitrophenols by certain bacteria, *Biochem. J.* 55, XXIV

Smith, W. H. (1969). Release of organic materials from the roots of tree seedlings. *Forest Sci.* , 15, 138-143

Smith, W. H. (1970). Root exudates of seedlings and mature sugar maple. *Phytopathology*, 60, 701-703

Spain, J.C., O. Wyss, and D.T. Gibson (1979), Enzymatic oxidation of p-nitrophenol. *Biochem. Biophys. Res. Commun.* 88, 634-641

Sprankle, P., W. Meggitt, and D. Penner (1957b). Adsorption, mobility, and microbial degradation of glyphosate in soil. *Weed Science*, 23, 229-234

Stark, J. (1982). Persistence of herbicides in forest soils. Dissertation. Swedish Univ. of Agricultural Sciences, Uppsala Dept. of Microbiology, Report 15

Stott, D. E., J. P. Martin, D.D. Focht, and K. Haider (1983). Biodegradation, stabilization in humus, and incorporation into soil biomass of 2,4-D and chlorocatechol carbons. *Soil Sci. Soc. Am. Proc.* 47, 66-70

Sudhakar-Barik, Siddaramappa, R., and N. Sethunathan (1976). Metabolism of nitrophenols by bacteria isolated from parathion amended soils. *Antonie Van Leeuwenhoek* 42, 461- 470

Sylvester, M., R. Masse, F. Messier, J. Fauteux, J-G. Bisailon, and R. Beaudet (1982). Bacterial nitration of 4-chlorobiphenyl. *Appl. Environ. Microbiol.* 44, 871-877

Thompson, D.G., G.R. Stephenson, K.R. Solomon, and A.V. Skepasts (1984). Persistence of 2,4-dichlorophenoxyacetic acid and 2-(2,4-dichlorophenoxy) propionic acid in agricultural and forest soils of northern and southern Ontario. *J.Agric. Food Chem.* 32, 578-581

Torstensson, N.T.L., and A.Aamisepp (1977). Detoxification of glyphosate in soil. *Weed Research*, 17, 209-212

Torstensson, L., and J. Stark (1979). Persistence of glyphosate in forest soils. *Weed and Weed Control*. 20th Swedish Weed Conference, pp 145-149

Torstensson, N.T.L. , and J. Stark (1981). Decomposition of ^{14}C labelled glyphosate in swedish forest soils. *Proceedings of the EWRS symposium on Theory and*

Practice of the use of soil applied herbicides, pp, 72-79

Torstensson, L. (1985). Behaviour of glyphosate in soils and its degradation. In: Grossband, E. and D. Atkinson (eds). The Herbicide Glyphosate. Butterworths, London

Vancure, V. (1964). Root exudates of plants.1- Analysis of root exudates of barley and wheat in their initial phases of growth. Plant and Soil 21, 231-248

Vancure, v., and J.L. Garcia (1969). Root exudates of reversibly wilted millet plants, " *Panicum miliaceum* ". Oecol. Plant. (Gautheir Villars) . 4, 93-98

Whipps, J.M. (1990). Carbon economy. In: Lynch, J. M. (ed.). The Rhizosphere. Wiley, Chichester, pp. 59-97

Zoeteman, B. C. J. , K. Harmren, J.B.H.J. Linders, C.F.H. Morra, and W. Slooff (1980). Persistent organic pollutants in river water and groundwater of the Netherlands. Chemosphere 9 , 231-249