

REFERENCES

- Abbott, D.C., Holmes and J.O.G. Tatton(1969)
Pesticides residues in the total diet in Eng-
land and Wales, 1966-1967; Organochlorine
pesticide residues in the total diet.
- Abdallah, El.A.M. and Morsy, F.A.R., 1980: "Side effects of
certain dithio carbamate fungicides". Egyptian
Society of Microbiology. Proc. IV Conf. Microbi-
ology, Cairo 1980. Vol. II. Plant ppathology
211-219.
- _____, Mokhtor, H.K. and Morsy, F.A., 1981. Relation
between chemical structure and insecticidal
activity. Part V-Biological activity of 2-Methyl-
6-styryl pyridiene dicarboxylic acid and its
derivatives. Proc. 4 th Arab Pesticide Conf. Tan-
ta. Univ. Vol IIIA, 235-245.
- Abdel-Gawaad, A.A. (1985): Survey of pesticides used in
Egypt. 2nd Int. Cong. for Soil Poll. and Prot.
from Pesticide Residues. Cairo, Vol II, 33 - 86.
- _____, and El-Gayar, F.H. (1972): Effect of some
soil insecticides on plants: III - Effect on ger-
mination, dry weight, and nodulation in Trifoli-
um Alexandrinum.
Indian J. of Agric. Sci. 42 (11) : 969.
- _____, (1973): Effect of some
soil insecticides on plants: II - Effect on germi-
nation of cotton seeds, growth, dry weight and
cotton yield. Cotton growing Review. 50: 152 -
160.

_____, F.H., Soliman, A. Rehab, F.I., and Ali, N.M. (1972 a) : Effect of some soil insecticides on plants : 1- On cotton, clover, corn and bean.
Phytopath. 2., 73 : 189 -200.

_____, El-Minshawy, A.M. and Zeid, M. (1972 b) : Studies on soil isecticides : VIII - Effect of some soil insecticides on broad beans and Egyptian clover nodule forming bacteria.
Z. Bakter. Parasit., 127 (2) : 172 - 177

_____, Khishen,, M.K. and Zakaria Seleim (1981): Soil pollution by pesticide residues: 18-Effect on sprouted onion roots. 1st Intern. Cong. soil. poll. PartII, 349-361.

Abdel-Kader M.H.K. (1982); Malathion and fenitrothion in stored wheat, low temperature degradation, distribution and quantification of metabolites.
Dissertation Abstracts International B: 42(8) 3173.

Abdel-Kader, M.H.K. and Webster G.R.S. (1982): Analytical methods developed and quantitative determination of malathion and fenitrothion and metabolites in stored wheat.
Dept. of soil scienc. Univ. of Manitoba-Canada

_____, (1983): Analytical methods developed and quantitative determination of malathion and fenitrothion and metabolites.

Sixteenth annual workshop on pesticide residues analysis (Western Canada) pp. 93 - 97

_____ and .RB. Webster (1980): Low temperature degradation and translocation of malathion

and fenitrothion in stored wheat.

Canadian Plains Proc. 9:143-153.

Abu-Elamayem M.M., A. Abdel-Rii, G.A. Tantawy, A.S.M. Marei (1979): Fate of leptophos in milk and wheat during the processing steps. Alexandria J. of Agric. Res. 27(3): 659-663

Adachi, K., N. Dhokuni and T. Metsuhashi (1984): Simple analytical method for organophosphorus pesticide determination in unpolished rice, using removal of fats by zinc acetate. J. Assoc. off. Anal. Chem. 67,4,798-800

Adler, I.L., C.K. Hofmann and S.S. Stawinski (1984). High-performance liquid chromatographic Analysis of Fenridazon potassium in wheat grain and straw. J. Agric. Food. Chem. 32 : 1352-1361

Al-Adil, K.M., White, E.R., Winterlin, W.L. and Kilgore, W.W. (1973): Uptake and translocation of Guthion by beans and barley. J. Agric. Food Chem. 21: (3), 376 - 379.

Alnaji, L.K. and A.M. Kadoum. (1979): Residues of Methyl phoxim in wheat and Milling fractions. J. Agric. Food. Chem. 27,3:583-584

(1981): Note on the effects of baking on methyl Phoxim and malathion residues in bread. Cereal chemistry 58 (1): 74 - 75.

Amer, S.M. (1965): Cytological effects of pesticides: I - mitotic effects of N-methyl-1-naphthyl carbamate "sevin". Cytologia, 30 (1): 175 - 181.

Rmer, S.M., E.M. Aji (1974) : Cytological effects of pesticides: V - Effect of some herbicides on *Vicia faba*. *Cytologia*, 39 (4) : 633 - 643.

_____ and Farah, O.R. (1974): Cytological effects of pesticides: VI - Effect of the insecticide "Roger" on the mitosis of *Vicia faba* and *Gossypium barbadense*.
Cytologia, 39 (4) : 507 - 514.

_____, Hammouda, M.R. and Farah O.R. (1971): Cytological and morphological effects of the insecticide N-methyl-1-naphthyl carbamate "sevin".
Flora, 160: 433 - 439.

Anderegg, B.N, Lichtenstein and Kemp, J.D. (1977): Effect of lindane on DNA, RNA and protein synthesis in corn roots.
J. Agric. Food Chem. 25(4): 923 - 928.

_____ and Madison L.J.(1983): Effect of insecticide distribution and storage time on the degradation of [^{14}C] Malathion in stored wheat.
J. Econ. Ent. 76(5), 1009-1013.

Anderson, R.F. (1955): Internal Medication of plants for the control of insect.
J.Econ.Ent. 48(2):187-190

Appaiah, K.M. ,R. Ramakrishna, K.R. Subba Rao, and O. Kapur. (1982) : Spectrophotometric determination of Carbaryl in grains.
J. Assoc. Off. Anal. Chem. 65: 32-34

_____, Nagaraja, K.U. and Kapur, O.P., (1982): Spectorphotometric method of determination of carbofuran residues.

J. Food Science and Tech., India, 19(5): 211-212.

Ashour, M.B.A.M. (1979): Side effects of phosphorus insecticides on plants.

D.Ph. Thesis, Faculty of Agriculture, Zagazig University.

Rult, J. A., Schofield, M., Johnson, L.D. and Waltz R.H. (1979):
Automated Gel Permeation Chromatographic preparation of vegetables, fruits and crops for organophosphate residue determination utilizing flame photometric detection.
J. Agric. Food Chem. vol. 27 : 825-828

Bai Qing-Yun and Chang-Wu Liu (1987) :
Dissipation of Propiconazole residues in and on wheat Triticum aestivum L.
Pestic. Sci. 19, 229-234

Balayannis P.G. (1983) : Effect of soil applications of Aldi carb on the growth, yield and chemical composition of the tobacco plants.
J. Agric. Food Chem. 31 : 1351-1355

Balyaeva, M.G. and Zh. L. Lukpanov (1975): Physiologic Characteristics of the effect of pesticides on cucumbers. Tr. Kaz. Nauchno-Issled.Inst. Zashch Rest.,13:116-121.

Barman, M.C., C.L. Molder and L.G. Rushing (1978): Trace analysis of etrimfos and two degradation products in corn and alfalfa.
J. of Agric. Food Chem. 26 (1): 35-42

Barraws, H.L., Caro, J.H., Armiger, W.H. and Edwards, W.M. (1969): Contribution of aerial contamination to

- the accumulation of dieldrin by mature corn plants.
Environmental Sc. Technol. 3: 261 -263.
- Barrentine B.F. and J.D.Cain 1969 Residues of endrin and DDT in soybeans grown on soil treated with these compounds .
Pesticides Monitoring J.III(2) ,77-79.
- Bates, R.N. and Rowlands, D.G. (1964): Interference by extractives in the determination of malathion residues in rice bran.
Analyst 89, 286 -287.
- Beitz, H., Seefeld, F. Hartisch, J. and Heinisch, E. (1971): Investigation on the uptake of DDT from the soil by plants with lipophilic constituents.
Rev. Appl. Ent. 59 (11) 1777 (3433).
- Bengston M., R.A.H. Davies, J.M. Desmarchlier, R. Menning and W. Marray (1983): Organophosphorothioates and synergised synthetic pyrethroids as grain protectants on bulk wheat.
Pestic. Sci. 14(4): 373-384
- Bennet, S.H. (1957). The behaviour of systemic insecticides applied to plants.
Ann. Rev. Ent. 2: 279 -295.
- Berck B. (1974): Fumigant residues of carbon tetrachloride, ethylene dichloride, and ethylene dibromide in wheat, flour, bran, middlings and bread.
J. Agric. Food Chem. 22:6:977-984
- Bhan, A.K. and Kaul, B.L. (1975): Cytogenetic activity of dichlorvos in barley.
Indian J. Exp. Biol. 13(4): 403 -405.

Bunnemaison, L. and Jourdheuil, P. (1951): Influence of various insecticides on the germination and growth of several cultivated plants.

Parasitica C. Gembloux 7 : 81 -108.

Bowman, M.C., C.L. Holder and L.G. Rushing (1978).

Trace analysis of etrimfos and two degradation products in corn and alfalfa.

J. Agric. Food Chem. 26(1) 35-42.

Bradburg, F.R. (1963): The systemic action of benzene hexachloride seed dressing.

Ann. App. Biol., 52: 361-361-370.

Bradburg, F.R. and Whitaker, W.O. (1956): The systemic action of benzene hexachloride in plants: Quantitative

measurements.

J. Sci. Food. Chem. 7(2): 248 -253.

Brass C.L. and Ware G. (1960): "BHC translocation from treated soil and the effect on growth of red clover.

J. Econ. Ent. 53(1): 110-113.

Brindley, T.A., Schopp, R. and Hinnan, F.G. (1950): Effect of initial high dosages of DDT on the yields of peas and wheat. J.Econ. Ent. 43: 565 - 567.

Bruce, R.B., A.J. Robbins and T.O. Tuft (1962): Phosphine residues from phostoxin-treated grain.

J. Agric. Food Chem. 10, 18-21

Caro, J.M. (1969): Accumulation by plants of organochlorine insecticides from the soil.

Phytopathology 59: 1191 - 1197.

_____ and Taylor, A.W. (1971): Pathways of loss of dieldrin from soils under field conditions.

J. Agric. Food Chem., 19(2): 379 -384.

- Carpentier, S. and Fromageot, C. (1950):** Activite mitotique des isomeres et de l'hexachlorocyclohexane, avec des observations sur l'influence du mesoinositol et du mesoinositophosphate de sodium. *Biochem. Biophys. Acta*, 5: 290 -296.
- Chapman, R.K. and Allen, T.C. (1948):** Stimulation and suppression of some vegetable plants by DDT. *J. Econ. Ent.* 41(4): 616 -623.
- Chapman, R.R. and J.R. Robinson (1977):** Simplified method for the determination of residues of carbofuran and its metabolites in crops using gas-liquid chromatography-mass fragmentography. *J. of Chromatography* 140 (3):209-218.
- Charnetski, W.R., Lichtenstein, E.P. and Evert, R.F. (1973):** Effect of lindane on cell structure of pea roots. *Can. J. Bot.* 51 (11): 211 -2117.
- Choi, S.Y. and Lee, H.R. (1977):** Control of some insects on soybeans with granular systemic insecticides applied in seedling-pits. *Hanguk Sikmul Poho Hakhoe Chi.* 16(1): 41 - 45.
- Chopra, S.L. and Nandra, K.S. (1969):** Effect of thiometon on the germination of Sarson. *J. Agric. Food Chem.* 17(4): 805 - 806.
- Clower, M. Jr., James P. McCarthy and D.M. Rains (1985):** Effect of cooking on levels of ethylene dibromide residues in rice. *J. Assoc. Off. Anal. Chem.* 68: 4: 710-711
- Clower, M. Jr., James P. McCarthy and L.J. Carson (1986):** Comparison of methodology for determination of

ethylene dibromide in grains and grain-based foods.

J. Assoc. Off. Anal. Chem. 69, 1: 87-93

Cordon, C. (1986): Gas chromatographic determination of flucythrinate synthetic pyrethroid residues in a range of crops.

J. Agric. Food Chem. 34(6): 953-955

Costet-Corio, M.F. and P Benveniste (1988): Sterol metabolism in wheat treated by N-substituted morpholines.

Pestic. Sci. 22: 343-357

Cullinan, E.P. (1949): Some new insecticides. Their effect on plants and soil.

J. Econ. Ent. 42(2), 387-391.

Das, N.M. and Chandrika, S. (1972): Effect of seed treatment with systemic insecticides on the germination of rice and the growth of seedlings.

Agr. Res. J. Kerala, 10(2): 152 -156.

David, W.R.L. (1951): Insecticidal action studies with bisdimethylaminophosphorus anhydride containing phosphorous 32. Ann. Appl. Biol. 38: 508-524.

_____ and Aldridge, W.N. (1957): The insecticidal action in leaves of plants growing in soil treated with parathion.

Ann. Appl. Biol. 45: 332-346.

_____ and B.O.C. Gardiner (1952): Insecticidal action studies with bisdimethyl aminofluorophosphine oxide containing 32 phosphorous.

Ann. Appl. Biol. 39(2): 203-210.

David, W.R.L. and Gardiner, B.O.C., (1955): The aphicidal action of some systemic insecticides applied to seeds. Ann. Appl. Biol 43 (4): 594-614.

Dennis, E.W. and Edwards, C. R. (1964): Phytotoxicity of insecticides and acaricides. III. Soil applications. *Plant Path. B.* 173-177.

Desmarchelier, J.M. (1976): Analysis of pyrethroids on grains.
J. Stored Prod. Res. 12, 245-252.

Desmarchelier, J.M. (1978): Mathematical examination of availability to insects of aged insecticide deposits on wheat.
Journal of stored products research 14(4): 213-222.

_____ : Loss of fenitrothion on grains in storage.
Pesticide Science 9 (1), 33-38.

Bewidar, R.E. and Rabie S.H. (1978): Effect of trifluralin on the growth regulators of Amaranthus chlorostachys. seedling
J.Faculty Ed. Ain Shams Univ. 17-26.

Devries, J.W., J.M. Broge, J.P. Schroeder, R.H. Bowers, P.A. Larson, and N.M. Burns (1985): Headspace gas chromatographic method for determination of methyl bromide in food ingredients.
J. Assoc. Off. Anal. Chem. 62,6: 1112-1116

Ding, H.D and I.S. Krull (1984): Trace analysis for organothio-phosphate agricultural chemicals by high-performance liquid chromatography-photolysis-electrochemical detection.
J. Agric. Food Chem. 32 C3: 622-628

Dumas T. (1980): Phosphine sorption and desorption by stored wheat and corn.

J. Agric. Food Chem. 28: 337-339

Edwards, C.A. (1965): Some side effects resulting from the use of persistent insecticides.

Biol. 55(22):329-331

Edwards, C.A. (1974): Environmental Pollution by pesticides.

"Environmental science research series". Pesticides residues in soil and water. PP 409-458. Blackwell. Oxford.

Eno-Charles, F. and Everett, P.H. (1958): Effect of soil application of 10 chlorinated hydrocarbon insecticides on soil microorganisms and the growth of stringless Black Valentine beans.

Soil Sci. Amer. Proc. 22(3): 235-238.

Foster, R.C. (1951): Some plants response to certain insecticides in soil.

USA Agric. Circ.: 862

Getzin, L.W. and Chapman, R.K. (1959): Effect of soil upon the uptake of systemic insecticides by plants.

J. Econ. Ent. 52(6): 1160-1165

Gimenez-Martin, G. (1962): Accion a-Motorica del parathion.

Phyton 18: 23-26 (cf Ashour, M.B., 1979)

Gomaa, E.A. (1977): Effect of organophosphorus insecticides on dry beans.

Actes Congr. Int. Compos Phosphores 1st 1977 (Pub. 1978), 671-676.

Gough, H.C. and Wood, A. (1954): Seed dressings for the control of wheat bulb fly.

Nature, 174(4442): 1151-1153.

Gowen, J.A. Wiersma, G.B. Tai, H. and Mitchell, W.G. (1976):
Pesticide levels in hay and soils from nine
states, 1971.
Pesticide Monitoring J. 10 (3), 114-116.

Hacskaýlo, J. (1957): Growth and fruiting properties and
carbohydrate, nitrogen and phosphorus levels
of cotton plants as influenced by Thimet.
J. Econ. Ent. 50(3): 280-284.

Lindquist, D.A. and Davich, T.B. (1961): Dime-
thoate adsorption and its translocation and
distribution in the cotton plants.
J. Econ. Ent. 54:1206-1209.

Hagley, E.A.C. (1965): Effect of insecticides on the growth of
vegetable seedling.
J. Econ. Ent. 58 : 777-778.

Haines R.G. (1956): Evidence of lindane translocation in corn
plants.
J. Econ. Ent. 49 (4): 563-564.

Handa S.K. (1979): Colorimetric determination of dillapiole in
wheat.
J. of the Ass. Off. Anal. Chem. 62(1):203-204

Harris, G.E. and Sans W.W. (1969): Vertical distribution of
residues of organo-chlorine insecticides in soil
collected from six farms in Southwestern On-
tario.
Proc. Ent. Soc. 100: 156-164.

Hassan, M. A. (1979): Cytological and physiological studies
of the effect of herbicide Dinitramine on roots of
Allium cepa and Vicia faba. (M.Sci. Botany), Ain

Shams University.

- _____, Saad, R.S. and Mansour, M.H. (1975): Effect of certain insecticides on some cotton pests and cotton plants. *Bull. Entomol. Soc. Egypt. Econ. Ser.* 8: 221-226.
- Heikes D.L. (1985): Purge and trap method for determination of ethylene dibromide in whole grains, milled grain products, intermediate grain-based foods and animal feeds. *J. Assoc. Off. Chem.* 68, 6: 1108-1111
- _____, and M.L. Hopper (1986): Purge and trap method for determination of fumigants in whole grains, milled grain products, and intermediate grain-based foods. *J. Assoc. Off. Anal. Chem.* 69: 990-998
- Hopkins, H.T. (1952): Inhibition of growth by benzene hexachloride isomers and protective effect of glucose as measured by cell counting technique. *Plant physiol.* 27: 526-540.
- Isshiki K., S. Tsumara, and T. Watanabe (1978): Residual piperonyl butoxide in agricultural products. *Bull. Environ. Cont. and Tox.* 19(5): 518-523
- Ishii, S. and Hirano, C.,(1962): *Japanese J. Appl. Zool.* 6:28.
- James, M.B., M. Desmarchelier, B. Hayward, Ronald Henni, J.H. Moulden, R.M. Noble, G. Smith, J.T. Snelson, R. Sticka, D. Thomas, B.E. Wallbank and D.J. Webley (1987) : Synergised Cyfluthrin and Cypermethrin as grain protectants on bulk wheat. *Pestic. Sci.* 0031-613 H 187/503 50: 23-37

Joia B.S., L.P. Sarna and G.R.S. Webster (1985): Gas chromatographic determination of cypermethrin and fenvalerate residues in wheat and milled fractions.

Inter. J. of Environ. Analytical Chem. 21(3):179-184

Joia, B.S., G.R.B. Webster and S.R. Loschiavo (1985): Cypermethrin and fenvalerate residues in stored wheat and milled fractions.

J. Agric. Food Chem. 33, 618-622

Kadoun R.M. and L. Alnaji (1978):

Effect of grain moisture content on the degradation rate of methyl phoxim in corn, sorghum and wheat.

J. Agric. Food Chem. 26(2) 507-509

Kathpal T.S. and R.S. Dewan (1976):

Endosulfan residues in/or sorghum under different agroclimatic conditions of India.

Indian J. of Agric. Sci. 46(8) 354-358

Kearney, P.C., Oliver, J.E., Kantson, R., Fiddler, W. and Pensabene, J.W. (1980): Plant uptake of dinitroaniline herbicide related nitrosamines. J. Agric. Food Chem. 28-633-635.

Keul, M., Dintila R. and Qchesanu (1976): Dynamics of carbohydrates and respiration in corn, following treatment with Duplitor, aldrin and heptachlor.

Stnd. Cercet. Biol. Ser. Biol. Veg. 28(2): 127-133.

_____ and Fabian R. (1977): Modification of the carbohydrate content of corn after lindane, hep-

tachlor and aldrin treatment.

Stud. Univ. Babeş. Bolyai, Ser. Biol. 22(1): 28-33.

Khalifa, M.A.S., El-Din, A.T., Ahmed A.H. and Halawa, A. (1981):
Phytotoxicity of certain carbamate insecticides.
Proc. 4th Arab Pesticide Conf. Tanta Univ. 1, 393-404.

Kobayashi, M., K. Yagi, M. Mori, K. Endo, R.K. Tanaka and
N. Umetsu (1987):
A method for residue analysis of benfuracarb and
its metabolites, carbofuran, 3-hydroxy-
carbofuran and
3-keto-carbofuran in crops by gas chromatogra-
phy.
J. of Pestic. Sci. 12(4): 689-697

Kostoff, D. (1949): Induction of cytogenetic changes and a
typical growth by hexachloro cyclohexane.
Science 109: 467-468.

Kozlova, E.N. (1950): Penetration of organic insecticides into
tissues. Doklady Uzesoyuz.
Ordenz Lenina Akad-Sel. SKD-Khoz. Nauk ison.
U.I.

Ku C.C., I.P. Kapoor and J.D. Rosen (1978):
Metabolism of Cytrolane (mephosfolan) systemic
insecticide (diethoxyphosphinyl) dithioimido
carbonic acid, cyclic propylene ester in a simu-
lated rice paddy.
J. of Agric. Food Chem. 26(6): 1352-1357

Kumar, A., R.B. Daharey, G.P. Pandey and B.K. Verma. (1983)
Determination of residues of pirimiphos ethyl in
stored food grains by thin - layer chromatogra-
phy.
Indian J. of Agric. Sciences 53,6: 460-462.

Lahue, D.W.,(1977): Pirimiphos methyl: effect on popultions of Tribolium confusum and T. castaneum in wheat.
J. Econ. Ent. 70(1): 135-137.

_____ and Dicke, E.B. (1977): Evaluation of selected insecticides applied to high moisture grain sorghum to prevent stored grain insect attack.
Marketing research Report USDA No. 1063, 10 pp.

Lawrence, J.F. and Leduc, R. (1977): Direct analysis of carbofuran and two nonconjugated metabolites in crops by high-pressure liquid chromatography with UV absorption detection.
J. Agric. Food Chem. 25(6): 1362-1365.

_____ (1978): High-pressure liquid chromatographic analysis of carbofuran and two non-conjugated methabolites in crops as fluorescent dansyl derivatives.
Journal of chromatography. 152(2): 507-513.

Lawrence J.F., L.G. Panopio and H.A. McLeod (1981):
Analysis of difenzoquat herbicide in wheat products by reversed phase liquid chromatography.
J. Agric. Food Chem. 29, 887-889.

Lazer-Keul, G., Soran, U., Dintila, and Keul, M. (1979): Effect of Lindane and Methylchlor on cell nucleus DNA content of wheat (Triticum vulgare) and broad bean (Vicia faba) root meristem. Rev. Roum. Biol. Ser. Biol. Veg. 24(1): 69-75.

Leahy, J.P. and E.A. Curl (1982):
The degradation of pirimiphos-method on stored grains.
Pestic. Sci. 13(5): 467-474

- _____ and P.K. Carpenter (1980):
The uptake of metabolites of permethrin by plants
grown in soil treated with permethrin.
Pestic. Sci. 11(2): 279-289
- Lee, S.R., S.Y. Kang and T.H. Kim (1980):
Occurrence of organochlorine insecticides in ve-
getable oils produced in Korea.
Korean J. of Food Sci. and Technology 12(3):
216-218
- Lee, Y.W. and Neil D. Westcott (1981):
Gas Chromatographic quantitative analysis and
persistence of dinmethoate and dimethoxon resi-
ues on and in wheat plants.
J. Agric. Food Chem. 29, 860-862
- Lichtenstein, E.P. (1959): Absorption of some chlorinated hy-
drocarbon insecticides from soils into various
crops.
J. Agric. Food Chem. 7(6): 430-433.
- _____, T.T. Liang and M.K. Koeppe (1985):
Penetration of volatile carbon-14 residues into
the aerial parts of plants grown adjacent to or in
[¹⁴C] phonofos or [¹⁴C] carbofuran-treated soils.
- _____, Myrdel, G.R. and Schulz, K.R. (1965): Adsorption
of insecticidal residues from contaminated soils
into five carrot varieties.
J. Agric. Food Chem. 13(1): 57-63.
- _____ and Schulz K.R. (1959): Breakdown of lindane
and aldrin in soils.
J.Econ. Ent. 52(1): 118-124.
- _____ and Millington, W.F. and Cowley G.T. (1962): Ef-

fect of various insecticides on growth and respiration of plants.

J. Agric. Food Chem. 10(3): 251-256.

_____ and G.T. Cowley (1963): Inhibition of the conversion of aldrin to dieldrin in soils with methylenedioxy phenyl synergists. J.Econ. Ent. 56(4): 485-489.

Lindquist, R.K., Krugger, H.R., Spadafora, R.R. and Mason, J.E. (1972): Application of aldicarb to green house tomatoes, plant growth , fruit yields, greenhouse whitefly control and residues in fruits. J.Econ. Ent. 65(3): 862-864.

Lockwood, L.M., S.K. Majumder and D.R. Lineback (1974): Degradation of Organophosphate Pesticides in cereal Grains during milling and cooking in India. Cereal Science Today. 19,8, 330-333

Lofroth, G. (1970): Alkylation of DNA by dichlorvos. Naturwissenschaften 57: 393-394.

Maclean, D.C., D.C. Mc Cune and R.E. Schneider (1984): Growth and yield of wheat and sorghum after sequential exposures to hydrogen fluoride. Environmental Pollution Series A, 36: 351-365

Magallona, E.Dand Celino, L.P. (1977): Organophosphate insecticide residues in grains. II. Effect of cooking on malathion and pyrimiphos methyl residues in rice. Philippine Entomologist. 3 (5/6) 295-299.

Mc Kay, G. (1985): Gas chromatographic determination of ethylene dibromide in flour products. J. Assoc. Off. Anal. Chem. 68, 2: 203-204

Mensah G.W.K. (1979): Persistence and translocation of organophosphorus insecticides on structural surfaces and in stored grain.
Abstracts International B, 40 (4): 1510

_____ (1987): Accumulation of malathion residues in grain stored on Malathion-treated wood surfaces.
Pestic. Sci. 20: 161 - 166.

_____, Watters, F.L. and Webster, G.R.S.(1979): Translocation of malathion, bromophos and iodofenphos into stored grain from treated structural surfaces.
J. Econ. Ent. 72(30): 325-39)

_____: Insecticide residues in milled fractions of dry or tough wheat treated with malathion, bromophos, iodofenphos and pirimiphosmethyl.
J. Econ. Ent. 72(5): 728-731

_____ (1979): Uptake of bromophos into bulk stored wheat from treated granary surfaces.
J. Econ. Ent 72(2): 275-276.

Metcalf, R.L. (1955): Advances in pest control.
Vol. (3). Interscience, Pub. N.York, London.

_____, Fukuto, T.R. and Ralph B. March (1957): Plant metabolism of dithiosyston and thimet. **J. Econ. Ent. 50(3): 438-445.**

_____ and Ralph B. March (1952): Behaviour of octamethyl pyrophosphoroamide in citrus plants.

J. Econ. Ent. 45(6): 988-977.

Mitchell, L.W., Smale, B.C. and Metcalf, R.L. (1960): **Advances in pest control. vol.(3).**
Interscience, Pub. Inc. N.York.

Mishra, G.M. and Sinha, S.P. (1979): **Effect of malathion on mitotically dividing onion (*Allium cepa*) root-tip.**
Cell. India J. Exp. Biol. 17(7): 716-717.

Morrison, H.E., Crowell, H.H., Grumb, S.E. and Lauderdale, R.W., Jr. (1948): **The effect of certain new soil insecticides on plants.**
J. Econ. Ent. 41(3): 374-378.

Mole, U.N. (1978): **Effect of systemic insecticides on the germination and subsequent growth of pea (*Pisum sativum* L.) seed.**
Seed Res. 6(1): 62-66.

Nethawy, A., Wilson, G.B. and Hoopingarner, R. (1965): **Cytological and genetic studies on the effect of Rue-lene.** J. Econ. Ent. 58(3): 511-513.

Noble, R.M. and D.J. Hamilton (1985): **Stability of cypermethrin and cyfluthrin on wheat in storage.**
Pestic. Sci. 16(2): 179-185

_____ and W.J.Osborne (1982): **Stability of pyrethroids on wheat in storage.**
Pestic. Sci. 13(3): 246-252

Nybohm, N. and B. Knutsson (1947): **Investigations on C-mitosis in *Allium cepa*. 1. The cytological effect of hexachlorocyclohexane.**
Hereditas 33: 220-234.

Parouchais, C. (1973): **Determination of diuron in wheat.**
J. of the RORC 56, 4: 831-835

- Pulizu, R. and Floru, S. (1972): Analela Institutului de Cercetari pertru Protectia Plantelor 8, 299-302.
- Pool, R.R.F. (1977): A rapid bioassay for pesticide phytotoxicity. J. Agric. Food Chem. 25(5):1216-1318.
- Rains, D.M. and J.W. Holder (1981): Ethylene dibromide residues in biscuits and commercial flour. J. Assoc. Off. Anal. Chem. 64, 5:1252-1254
- Rajanna, B. and Dela Cruz, R.R. (1977): Stands establishment and early growth of field crops as influenced by seed vigor and pesticide residues. Seed Sci. Technol. 5(1): 71-85.
- Rajkannu, K. and K.K. Krishnamoorthy (1979): Uptake and persistence of phorate and carbofuran in I.R. 20 rice. Riso 28 (4): 339-343
- _____, M.S. Venugopal, P. Vasudevan and M. Balasubramanian (1979): Residues of quinalphos, phosalone and malathion in/on sorghum. Current Sci. 48(2):71
- Rangaswamy, J.R. (1985): Phosphine residues and its desorption from cereals. J. Agric. Food Chem. 33: 1102-1106
- _____ and M. Muthu (1985): Spectrophotometric determination of phosphine residues in rice. J. Assoc. Off. Anal. Chem. 68, 2: 205-208

- Reddy, M.U. and Rao, B.U. (1969): Cytological effects of insecticides (Dimecron-100 and Roger-40) on Vicia faba. Cytologia 34: 408
- Reynolds, R.L. (1958): Research advances in seed and soil treatment with systemic and non-systemic insecticides. Advances in pest control Research Vol.II. Ed. Metcalf, R.L. Interscience Publisher Inc., N.York.
- Ridgeway, R.L., Jones, S.L., Coppedge, T.R. and Lindquist, B.A. (1968): Systemic activity of 2-methyl-2-(methyl thio)propionaldehyde O-(methyl carbonyl) oxime (U.C. 21149) in the cotton plant with special reference to the both weevil. J. Econ. Ent. 61(6): 1705-1712.
- Ripper, W.E. (1957): Advances in Pest Control Research, Vol (I), Interscience publisher, N.York, London.
- Rodrigues, J.G., Maynard, D.E. and Smith W.T. Jr. (1960): Effect of soil insecticides and absorbents on plants sugars and resulting effect on mite nutrition. J. Econ. Ent. 53(4): 491-495.
- Rowlands, D.G. (1964): Degradation of malathion on stored maize and wheat grains. J. Sci. Food Agric., 15, 824-829.
- Rowlands, D.G. and J.E. Clements (1965): The degradation of malathion in rice brans. J. Stored Prod. Res. 1, 101-103
- _____ and J.S. Bromhall (1977): The uptake and translocation of malthion by the stored wheat grain. J.Store Prod. Res. 13, 13-22.

- Rus, B. Funduc, I. Crainiceanu, R. and Trestianu, S. (1977):
Gas chromatographic column for separation of
organochlorine insecticide residues.
Analytical Chem. 49(13): 2123-2124.
- Saivaraj, K., Habeebullah, B., Balasubramanian, G. And Bala-
subramanian, M. (1980): Insecticide residues in
ragi. *Madras Agric. J.* 67(3): 192-193.
- Sastry C.S.P., D.Ujaya and R.K. Ekambareswara (1986):
Sensitive spectrophotometric determinations of
carbaryl and propoxur in formulations, water,
grains and pulses.
Food Chem. 20(2): 157-162
- Sax, K. and Sax, H. (1968): Possible mutagenic hazards of
some food additives, beverages and insecticides.
Japan. J. Genetics 43: 89-94.
- Scheel, D., W. Schafer, and H. Sandermann (1984): Metabo-
lism of pentachlorophenol in cell suspension
cultures of soybean (*Glycine max* L.) and wheat
(*Triticum aestivum* L.). General results and isola-
tion of lignin metabolites.
J. Agric. Food Chem. 32: 1237-1241
- Scheidt, I., W. Pfannhauser and H. Woidich (1982): Determi-
nation of pyrethrins in flour.
Recent developments in food analysis. pp. 461-
463.
- Schloess, M.E. (1953): The effect of hecachlorcyclohexane
in roots of the onion (*Allium cepa*) and straw-
berry (*Fragaria vesca*).
J.Hortic. Sci. 28: 49-67.
- Schloess, M.E. (1955): The effect of aldrin, dieldrin, isodrin,

endrin and DDT on mitosis of roots of onion (Allium cepa).

J. Hortic. Sci. 30: 181-187.

Scopes, N.E.A.(1969): Some quantitative effects of soil applied organophosphorus insecticides on crop growth. Plant Path. 18(1): 10-15.

Scott, P.M., P. Lan and S.R. Kanhere (1981): Gas chromatography with electron capture and mass spectrometric detection of deoxynivalenol in wheat and other grains.

J. Assoc. Off. Anal. Chem. 44, 6: 1364-1371

_____ (1985): Effectiveness of cold solvent extraction system for the determination of fumigant residues in cereal grains.

Pestic. Sci.18: 33-53

Scudamore K.A. (1987): Fumigant residues in wheat and other cereal grains resulting from the use of 1,1,1-trichloroethane and bromoethane as a liquid fumigant mixture.

Pestic. Sci. 20: 1-10

_____ and Goodship G. (1986): Determination of phosphine residues in fumigated cereals and other foodstuffs.

Pestic. Sci. 17, 185-195

Shanimov, Kh.I. and Isin, M.M. (1977): Effect of insecticides on apple tree development. Khim. Selsk. Khoz. 15(5); 61-63.

Sherjugjit Singh and R.P. Chawla (1980): Comparative persistence of residues of pirimiphos-methyl on store wheat, maize and paddy.

Bull. of grain tech.18(3):181-187

- Shimabukuro, R.H., W.C. Walsh and R.A. Hoerauf (1979):
Metabolism and selectivity of diclofop-methyl in
wild oat and wheat.
J. Agric. Food Chem. 27; 3 : 615-623
- Sidney, D. (1975): Diazinon and carbaryl effects on the veg-
etation of soybean ecosystem.
Bull. N. York Acad. Sci. 20(1): 1-4.
- Simkover, M.G. and Shenefelt, R.D. (1952): Phytotoxicity of
some insecticides to coniferous seedlings with
particular reference to benzenhexachloride.
J. Econ. Ent. 45(1): 11-15.
- Simon, M. and Guinochat, M. (1939): Anomalies morpholo-
giques et caryologiques provoquées sur les
jeunes plantules, par les dérivés halogènes des
carbures cycliques.
C.R. Liebd. Seances Mean. Soc. Biol. 131: 22-229.
- Sinclair, W.B., D.L. Lindgren and R. Forbes (1962): Recovery
of ethylene dibromide residues from fumiwhole
kernel and milled wheat fractions.
J. Econ. Ent. 55, 6: 836-842
- Singh, D.P., Mathon, S.m. and Gangrads, G.A. (1974): Granular
insecticides on soybean stem fly, *Melanogromy-
za phaseoli*.
Pesticides 8(1): 31-32.
- Singh, B.D., Singh, R.B., Singh, R.M. and Singh, J. (1979): Ef-
fect of insecticides on germination, early
growth and cytogenetic behavior of barley (Hor-
deum vulgare). Environ. Exp. Bot. 19(3): 127-
132.
- , Singh, Y., Singh, R.B., Singh, R.M., Singh, N.D. Shar-
ma, R.K. and Singh, J. (1977): Cytogenetic aber-
rations and morphological changes induced by

insecticide treatments of barley seeds.
Indian J. Exp. Biol. 15(8): 688-691.

Smith D.C.(1971): Pesticide residues in the total diet in
 Canada.
Pestic. Sci. 2, 92-96

Smith R.E. (1984): Gas Chromatographic method for Analysis
 of 2,4-D in wheat: Interlaboratory Study.
J. Assoc. Off. Chem. 67, 4: 794-797

Smith, G.N., Watson, B.S. and Fischer, F.S. (1967): Investiga-
 tion on dursban insecticide. Uptake and translo-
 cation of (³⁶Cl) 0,0-diethyl 0-3,5,6-trichloro-2-
 Pyridyl phosphorothioate and (¹⁴C)0,0-diethyl-
 3,5,6-trichloro-2-pyridyl phosphorothioate and
 (¹⁴C)0,0-diethyl-3,5,6-trichloro-2-pyridyl phos-
 phorothioate by beans and corn.
J.Agric.Food Chem. 15(1): 127-131.

Starnes, E. (1950): adsorption -translocation of insecticides
 through the root system of plant.
J. Econ. Ent. 43(3): 338-342.

Strek, H.J. and Weber, J.B. (1980): Absorption and transloca-
 tion of polychlorinated biphenyls (PCBs) by
 weeds. *Proc. South Weed Sci. Soc.* 33: 226-232.

_____, Shea, P.J., Mrozek, E. Jr. and Over-
 cash, M.R. (1981): Reduction of polychlorinated
 biphenyl toxicity and uptake of carbon-14 activ-
 ity by plants through the use of activated car-
 bon.
J. Agr. Food Chem. 29, 288-293.

Strelkova, E.A. (1975): Effectiveness of treating barley
 seeds with insecto-fungicides under the condi-
 tions of the northeastern part of Bielorussia.

Tr. Bielorus. S. Kh. Akad. 140: 10-14.

Sullivan, J.H. (1980): Pesticide residues in imported spices. A survey for chlorinated hydrocarbons.
J. Agric. Food Chem. 28, 1031-1034.

Suzuki, T. Ishikawa, K., Sato, N., Sakai, K.I. (1979): Determination of chlorinated pesticide residues in foods. III. Simultaneous analysis of chlorinated pesticide and phthalate ester residues by using AgNO₃-coated florasil column chromatography for cleanup of various samples.
J.AOAC. 62(3): 689-694.

Szeto, S.Y., Wilkinson, R.T.S. and Brown, M.J. (1984): A gas chromatographic method for the determination of bendiocarb in soil and corn: Application to the analysis of residues in corn.
J. Agric. Food Chem. , 32, 78-80

Taleker, N.S., Lee, E.M. and Sun, L.T. (1977): Absorption and translocation of soil and foliar applied ¹⁴C-Carbofuran and ¹⁴C-phorate in soybean and mung bean seeds.
J. Econ. Ent. 7(6): 685-688.

_____, Chen, J.S. and Sun, L.T. (1978): Insecticide residues in subtropical soil : Their degradation, persistence and absorption into root crops.
Biotrop. Spec. Publ. 7, 105-115.

_____ and Lichtenstein, S.B. (1971): Influence of plant nutrition on lindane penetration and its translocation with pea plants.
J.Agr. Food Chem. 19(5): 846-850.

Tanski, H.H., C. Rosenberg, H. Siltanen, S. Kilpi and P. Simojoki (1985):
The effect of the annual use of pesticides on

soil microorganisms, pesticide residues in the soil and barley yields.

Pestic. Sci. 16, 141-148

Taylor, I.S. and F.P. Keenan (1970): Studies on the analysis of hexachlorobenzene residues in foodstuffs. J. of the RORC 53, 6:1293-1295

Thielert, W., W. Steffens, F.Fuhr, H.Kuck and H. Scheinpflug (1988): Uptake of triadimenol through wheat Caryopses after application by seed treatment. Pestic. Sci. 22: 93-105

Tietz, H. (1954): Hofchen-Brief, 7:1-55. (C.F. Talekar and Lichtenstein, 1971).

Tranto, C.M., Fraga, C.G. and Naves, O.S. (1960): Effect of application of insecticides to the soil on growth of cotton.

Soil and fertilizers, 24 (306), 2, 148.

Turner, B.C. and Caro, J. (1973): Uptake and distribution of carbofuran and its metabolites in field grown corn plants.

J. Environmental Quality, 2,2: 245-247.

Vaarama, A. (1947): The influence of DDT pesticide upon plant mitosis.

Hereditas 33: 191-219.

Van Middelen, C.H. (1969): Residues in food and feed. Pesticide

Monitoring Journal 3,2, 70-71.

Diljoen, J.H., Coatzer J.J., Vermaak, C.J., Pope A.R., Dixon, J.B.P. and Nagy, J. (1984): Dichlorvos from high pressure cylinders and aerosol cans for the

- control of storage pests in mills.
Phytophylactica, 16(4): 267-274.
- Dolpi, M. (1951): Cytological and lethal action of some phosphoric esters used as insecticides.
Ann. Sper. Agrar. 5: 1511-1525.
- Wagner, K. and Frehse, H. (1976): Method for the determination of residues of Folimat insecticide/ acaricide.
Pflanzenschutz-Nachrichten Bayer 29(1):54-66.
- Wallace, C.R (1947): Small-scale tests with DDT and Benzene Hexachloride incorporated in the soil.
J. Australian Inst. Agric. Sci. 13:132-137.
- Way, M.J. (1959): Experiments on the mode of action of insecticidal seed dressings especially against *Leptohy lemia Coarctata* F. muscidae , the wheat bulb fly.
Ann. App. Biol. 47(4): 783-801.
- Weaver, J.B., All, J.N., Weaver, D.B. and Hornyak, E.P. (1979): Influence of various insecticides on yield parameters of two cotton genotypes.
J.Econ.Ent. 77(1) 119-122.
- Webley, D.J. and Kilminster, K.M. (1980): The persistence of insecticide spray deposits on woven polypropylene and jute sacking.
Pesticide Science: 11:(6): 667- 673.
- Wedzisz, A.T., Stoczynska and J. Szwejda (1977): Effect of the preparation gardona 24 E.C. on some components of vegetables.
Bromatol. Chem. Toksykol 10(1); 29-34.
- Westcott, N.D. (1985): Gamma-HCH in rape seedlings grown

from treated seeds.

Pestic. Sci. 16, 416-421

Wheeler, W.B., Maye, H.A., Van Middelen, C.H., Thompson, N.P. and Tappan, W.B. (1969): Residues of endrin and DDT in turnips grown in soil containing these compounds. Pesticides monitoring J. 3(2): 72-76.

White, N.D.G. (1986): Uptake of malathion and pirimiphos-methyl by rye, wheat or triticale stored on treated surfaces.
J. Econ. Entomol. 78(6): 1315-1319.

Winteringham, F.P.W., A. Harrison, R.G. Bridges and P.M. Bridges (1955): The fate of labelled insecticide residues in food products: 11- The nature of methyl promide residues in fumigated wheat.
J. Sci. Food Agric. 6, 251-261

Wiresma, G.B., Mitchell, W. and Stanford, C.L. (1972):
Pesticide residues in onions and soil.
Pestic. Monit. J., 5: 354-347.

Wuu, K.D. and Grant, W.F. (1967): Chromosomal aberrations induced by pesticides in mitotic cells of barley.
Cytologia 32: 31-41.

Zaki, Mohib and Reynolds H.T. (1961)
Effect of various soil types and methods of application upon uptake of three systemic insecticides by cotton plants in the green house
J. Econ. Ent. 54(3): 568-573.