

REFERENCES

REFERENCES

- Abdallah, A.R. (1972).** Studies on the effect of moisture content of the soil on the rhizosphere microflora of different plants. M.Sc. Thesis, Fac. of Agric., Univ. of Assiut.
- Abdallah, A.R.; El-Dahtory, T.H., and Abdel-Moneim, A.A. (1986).** Effect of salinity on nodulation of some leguminous plants. Minia J. Agric. Res. & Dev. Vol.8(2): 504-511.
- Abdel-Aziz, R.A. (1987).** Ecological studies on *Rhizobium trifolii*. Ph. D. Thesis, Faculty of Agric. Ain Shams Univ., Cairo, Egypt.
- Abdel-Ghaffar, A.S. (1987).** Factors affecting nitrogen fixation and yield of faba bean under Egyptian field conditions. Arid Soil Research and Rehabilitation, 1: 65-75.
- Abdel-Wahab, S.M.; Hassan, M.E.; Saber, H.A.; Saleh, S.A., and El-Waraky, M.K. (1995).** Symbiotic performance of some naturally saline tolerant strains of *R. leguminosarum* bv. *viceae* under soil salinity stress. Minia J. Agric. Res. & Development. 17: 35-47.
- Abdel-Wahab, S.M.; Mehrashan T. El-Mokadem; Fatma A. Helemish, and Mona M. Abou El-Nour (1993).** The symbiotic performance of *Bradyrhizobium japonicum* under stress of salinized irrigation water. In " cuurent development in soybean - *Rhizobium* symbiotic nitrogen fixation (Ed. Dou

Xintian), Heilingjiang Science and Technology Publishing House, Harbin, China, pp. 442-454.

- Abu-Shakra, and Bassir, A. (1972).** Effect of inoculation and nitrogen fertilization nodulation, seed, yield and quality of soybean. *J. Agric. Sci.* 78: 179-182.
- Afza, R., Hardarson, G., Zepata, F., Fried, M., and Blume, W.E.H. (1983).** Effect of starter fat soil and foliar fertilizer application of N-15 labelled urea on yield and N-fixation of soybean. *Agron. Abst.* p. 163.
- Al-Rashidi, R.K. Loyna Chan, T.E., and Fredrick, L.R. (1982).** Desiccation tolerance of four strains of *Rhizobium japonicum*. *Soil Biol. and Biochem.* 14: 489-493.
- Al-Rashidi, R.K., and Aziz N.Y. (1990).** Efficiency and persistance of alfalfa rhizobia in soils as affected by salinity and desiccation. *Z. Mikrobiol.* 145: 195-202.
- Alexander, M. (1984).** Ecological constraints on nitrogen fixation in agriculture ecosystems. *Adv. Microbiol. Ecol.* 8: 163-183.
- Allos, H.F., and Bartholomew, W.V. (1959).** Replacement of symbiotic fixation by available nitrogen. *Soil Sci.* 87: 61-66.
- Allen, O.N., and Allen, E.K. (1950).** Biochemical and symbiotic properties of the rhizobia. *Bacterial. Rev.* 14: 273-330.

- Asanuma, S., and Ayanaba, A. (1990).** Variation in acid-Al tolerance of *Bradyrhizobium japonicum* strains from African soils. *Soil Sci. Plant Nutr.* 36: 309-317.
- Ayanaba, A., Asanuma, S., and Munns, D.N. (1983).** An agar plate method for rapid screening of *Rhizobium* for tolerance to aluminium stress. *Soil Sci. Soc. Amer. J.* 47: 256-258.
- Bala, N.; Sharma, P.K., and Lakshminarayana, K. (1990).** Nodulation and nitrogen fixation by salinity-tolerant in symbiosis with tree legumes. *Agric. Ecosys. Environ.* 33: 33-46.
- Baldani, J. Ivo, and Weaver R.W. (1992).** Survival of clover *Rhizobia* and their plasmid cured derivatives in soil under heat and drought stress. *Soil Biol. Biochem.* vol 24, No. 8, pp. 737-742.
- Bekki, A., Trinchant J.C., and Rigand I. (1987).** Nitrogen Fixation (C_2H_2 reduction) by *Medicago* nodules and bacteroids under sodium chloride stress. *Plant. Physiol.* 71: 61-67.
- Bhardwaj, K.K.R. (1975).** Survival and symbiotic characteristics of *Rhizobium* in saline-alkali soils. *Plant & Soil.* 43: 377-385.
- Bolshnoi, U.R. and Pancholy, D.K. (1980).** Comparative salt tolerance in triticale, wheat and rye during germination. *Plant and Soil.* 55: 491-493.
- Booth, I.R. (1985).** Regulation of cytoplasmic pH in bacteria. *Microbiol. Rev.* 49: 359-378.

- Bordeleau, L.M., and Prevost, D. (1994).** Nodulation and nitrogen fixation in extreme environments. *Plant and Soil* 161: 115-125.
- Bordeleau, L.M., Antoun, H., and Lachance, R.A. (1977).** Effects des souches de *Rhizobium meliloti* et des coupes successives de la luzerne (*Medicago sativa*) sur la fixation symbiotique d'azote. *Can. J. Plant Sci.* 57: 433-439.
- Bottomley, p.J. (1992).** Ecology of *Bradyrhizobium* and *Rhizobium*. In: *Biological nitrogen fixation*. pp. 293-347. (Eds. Stacy G.; Burris, R.H. and Evans, H.J.). Chapman and Hall, New York and London.
- Brockwell, J., and Whalley, R.D.B. (1970).** Studies on seed pelleting as an aid to legume seed inoculation. Z. Survival of *Rhizobium meliloti* applied to medic seed sown into clay soil. *Aust. J. Exp. Agr. Anim. Husb.* 10: 455-459.
- Brockwell, J., Pilka, A., and Holliday, R.A. (1991).** Soil pH as a major determinant of the numbers of naturally-occurring *Rhizobium meliloti* in non-cultivated soils of New South Wales. *Aust. J. Exp. Agric.* 31: 211-219.
- Bushby, H.V.A. (1990).** The role of bacterial surface charge in the ecology of root-nodule bacteria: a hypothesis. *Soil Biol. Biochem.* 22: 1-9.

Bushby H.V.A., and Marshall K.C. (1977 a). Some factors affecting the survival of root-nodule bacteria on desiccation. *Soil Biol. Biochem.* 9: 143-147.

Bushby, H.V.A. and Marshall, K.C. (1977 b). Water Status of rhizobia in relation to their susceptibility to desiccation and to their protection by montmorillonite. *J. of Gen. Microbiol.* 99, 19-27.

Busse, M.D., and Bottomley, P.J. (1989). Growth and nodulation response of *Rhizobium meliloti* to water stress induced by permeating and non permeating solutes. *Apl. Environ. Microbiol.* 55: 2431-2436.

Buttery, B.R., Park, S.J., and Dhanvantari, B.N. (1990). Effects of combined nitrogen *Rhizobium* strain and substrate on a supernodulating mutant of *phaseolus vulgaris* L. *Can. J. Plant Sci.* 70: 955-963.

Buttery B.R., Park S.I., and Findly, W.I. (1987). Growth and Yield of white bean (*Phaseolus vulgaris* L.) in response to nitrogen, phosphorus and potassium fertilizers and to inoculation with *Rhizobium*. *Can. J. Plant Sci.* 67, 425-432.

Caetano-Anolles, G., Lagares, A., and Favelukes, G. (1989). Adsorption of *Rhizobium meliloti* to alfalfa roots: dependence on divalent cations and pH. *Plant and Soil.* 117: 67-74.

Carroll, B.J., McNeil, D.L., and Gresshoff, P.M. (1985). Super-nodulation and nitrate tolerant

symbiotic (nts) soybean mutant. Plant Physiol. 78: 34-40.

Casino-Colon, A., and Marquis, R.E. (1988). Role of the arginine leiminase system in protection oral bacteria and an enzymatic basis for acid tolerance. Appl. Environ. Microbiol. 54: 1318-1324.

Caudry-Rezinck, S., Prevost, D., and Schulman, H.M. (1986). Some properties of Archtic rhizobia. Arch. Microbiol. 146: 12-18.

Chao, W. and Alexander M. (1982). Influence of soil characteristics on the survival of *Rhizobium* in soils undergoing drying. Soil Sci. Soc. Amer. J. 46: 949-952.

Chatel, D.L., and Parker, C.A. (1973 b). Survival of field grown rhizobia over the dry summer period in Western Australia. Soil Biol. Biochem. 5: 415-423.

Chen, H., Richardson, A.E., and Gartner, E. (1991). Construction of an acid tolerant *Rhizobium leguminosarum* biovar *trifolii* strain with enhanced capacity for nitrogen fixation. Appl. Environ. Microniol. 57: 2005-2011.

Cooper, J.E. (1982). Acid production, acid tolerance and growth rate of lotus rhizobia in laboratory media. Soil Biol. Biochem. 14: 127-131.

Cunningham, S.D., and Munns, D.N. (1984). The correlation between extracellular polysaccharide

production and acid tolerance in *Rhizobium*. Soil Sci. Soc. Amer. J. 48: 1273-1276.

Dart, P.J., Day, J., Islam, R., and Dobereiner, J. (1976). Symbiosis in tropical grain legumes : Some effects of temperature and the composition of the rooting medium. In: Symbiotic Nitrogen Fixation in Plants (P.S. Nutman, Ed.). Cambridge Press. Cambridge. UK. pp. 361-384.

Date, R.A., and Halliday, J. (1979). Selecting *Rhizobium* for acid, infertile soils of the tropics. Nature (London). 277: 62-64.

Date, R.A., and Roughley, R.J. (1977). Preparation of legume seed inoculants. In: A treatise on dinitrogen fixation. Agronomy and Ecology. Edited by R.W.F. Hardy and A.H. Gibson. John Wiley and Sons, New York. pp. 243-275.

Davis, L.C., and Imsande, J. (1988). Direct test for altered gas exchange rates in water-stressed soybean nodules. Ann. of Bot. 61: 169-177.

Day, D.A., B.J. Carroll, A.C. Delves, and Gresshoff, P.M. (1989). Relationship between autoregulation and nitrate inhibition of nodulation in soybeans. Physiol. Plant. 75: 37-42.

Day, D.A., Price, G.D., Schuller, K.A., and Gresshoff, P.M. (1987). Nodule physiology of a supermodulating soybean (*Glycine max*) mutant. Aust. J. Plant Physiol., 14: 527-538.

Dazzo, F.B., Hollingsworth, R.T., Sherwood, J.E. Abe. M., and Hrabak, E.M. (1985). Recognition and

infection of clover root hairs by *Rhizobium trifolii*. Appl. Environ. Microbiol. 48: 1140-1150.

Diatolff, A. (1970). Relationship of soil moisture, temperature and alkalinity to soybean nodulation failure. Q. J. Agric. Anim. Sci. 27: 279-293.

Diatolff, A. (1967). Effect of soil moisture fluctuations on legume nodulation and nitrogen fixation in a black earth soil. Queensland J. Agric. and Animal Sciences. 24: 315-321.

Drevon, J.J., Heckmann, M.O., Soussana, J.F, and Satsac, L. (1988 a). Inhibition of nitrogen fixation by nitrate assimilation in legume. *Rhizobium* symbiosis. Plant Physiol. Biochem. 26: 197-203.

Eaglesham, A.R.J., and Ayanaba, A. (1984). Tropical stress ecology of rhizobia, root nodulation and legume fixation. In: current developments in biological nitrogen fixation. Edited by N.S. Subba Rao. Edward Arnold Publishers, London, U.K. pp. 1-35.

Eaglesham, A.R.J.; Stowers, M.D.; Maina, M.I.; Goldman, B.J; and Ayanaba, A. (1987). Physiological of *Bradyrhizobium* sp. (Vigna) from three West African soils. Soil Biol. Biochem. 19: 575-581.

Ebady, D.M.N. (1986). Effect of NPK and foliar fertilizers on nodulation of nitrogen, growth and seed yield of soybean under field condition. Ph.D. Thesis, Fac. Agric. Tanta Univ.

- EK-Jander, J., and Fahraeus, G. (1971).** Adaptation of *Rhizobium* to subactic environment in Scandinavia. Plant and Soil Spec. p. 1971: 129-137.
- El-Essawi, T.M., Balba, A.A., and El-Said, M. (1982).** Effect of soil salinity and nitrogen dressing on symbiotic nitrogen fixation in two soybean varieties. J. Agric. Res., Tanta Univ. 8: 10-15.
- El-Shakweer, M.H.A., and Barakat, M.A. (1984).** Nitrogen fixation and phosphorus, potassium and water uptake by faba bean as affected by salinity stresses. 2nd Conf. of the Agric. Res. Center, Giza, Egypt. 9-11.
- El-Sheikh, E.A.E., and Wood, M. (1989 a).** Response of chickpea and soybean rhizobia to salt: influence of carbon source, temperature and pH. Soil Biol. Biochem. 21: 883-887.
- El-Sheikh, E.A.E. and Wood, M. (1989 b).** Response of chickpea and soybean rhizobia to salt: Osmatic and specific ion effects of salt. Soil Biol. Biochem. 21: 889-895.
- El-Sherbeeny, M.H., Nassib, M., and Rizk, M.A. (1982).** Effect of NPK fertilizers on nodulation seed yield and yield components of soybean growth on sandy calcareous soil. The First Inter. Africian Conf. on Biofertilizers, Cairo, Egypt. 22-26 March.
- Elsaeid, N.B.; Wothum, A.G., and Musse Lwhite, M.S. (1990).** Growth temperature tolerance and survival

of peanut Rhizobia in peat inoculants Turrlaiba
Vol. 40, No.3, pp. 287-291.

Foulds, W. (1971). Effect of drought on three species
of *Rhizobium*. Plant & Soil. 35: 665-667.

Frier, J.R. (1984). Important limiting factors in soil
for the *Rhizobium* - legume symbiosis. In: BNF
Ecology, Technology and physiology. Ed. M.
Alexander. pp. 51-74. Pleneem Press, New York,
London.

Fuhrmann J., Davery C.B., and Wollum, A.G. II (1986).
Desiccation tolerance of clover rhizobia in
sterile soils. Soil Sci. Soc. Amer. J. 50: 639-
644.

Gewaily, E.M., and Khedr, A.K. (1985). Effect of
nitrogen fertilization on symbiotic fixation and
growth of broad bean. 6th Inter. Symp. on N.
Fixat., Aug. 4-10. Oregon, U.S.A.

Gibson, A.H. (1967). Physical environment and symbiotic
nitrogen fixation. IV. Factors affecting the
early stages of nodulation. Aust. J. Biol. Sci.
20: 1087-1094.

Gibson, A.H. (1977). The influence of the environment
and symbiotic nitrogen fixation. VI. Nitrogen
relation within the nodules of *Trifolium*
subterranean L. Aust. J. Biol. Sci. 22: 829-838.

Gibson, A.H., and Harper, J.E. (1985). Nitrate effect on
nodulation of soybean by *Bradyrhizobium*
japonicum. Crop Sci. 25: 497-501.

- Gibson, A.H., and Jordan, D.C. (1983).** Ecophysiology of nitrogen fixing systems In: Encyclopedia of plant physiology New Series. Vol. 12C: 301-390.
- Gibson, A.H., and Nutman, P.S. (1960).** Studies on the physiology of nodule formation. VII. A reappraisal of the effect of preplanting. Ann. Bot. (London) 24: 420-433.
- Graham, P.H. (1979).** Influence of temperature on growth and nitrogen fixation in cultivars of *Phaseolus vulgaris* L. inoculated with *Rhizobium*. J. Agric. Sci. Camb., 93: 365-370.
- Graham, P.H. (1992).** Stress tolerance in *Rhizobium* and *Bradyrhizobium*, and nodulation under adverse conditions. Can. J. Microbiol. 38(6): 475-484.
- Graham, P.H. and Rosas, J.C. (1978).** Nodule development and nitrogen fixation in cultivars of *Phaseolus vulgaris* L. as influenced by planting density. J. Agric. Sci. 90: 19-29.
- Graham, P.H., and C.A. Parker (1964).** Diagnostic features, in the characterisation of the root nodule bacteria of legumes. Plant and Soil. 20, 383-396.
- Graham, P.H., Draeger, K.J., and Ferrey, M.L. (1992).** Occurrence and characterization of acid-pH tolerance in strains of *Rhizobium* and *Bradyrhizobium* submitted Can. J. Microbiol.
- Graham, P.H., Parker, G.A., Oakley, A.E., Tange, R.T., and Sanderson, J.J.V. (1963).** Spore formation and

- heat resistance in *Rhizobium*. J. Bacteriol. 86: 1353-1354.
- Graham, P.H.; Viteri, S.E.; Mackie, F.; Vargas, A.T., and Palacias, A. (1982). Variation in acid soil tolerance among strains of *Rhizobium phaseoli*. Field Crops Res. 5: 121-128.
- Habish, H.A., and Khairi, S.L.M. (1970). Nodulation of legumes in the Sudan. II. *Rhizobium* strains and cross inoculations of *Acacia* spp. Expt. Agric. 6: 171-176.
- Hafeez, F.Y.; Asad, S., and Malik, K.A. (1991). The effect of high temperature on *Vigna radiata* nodulation and growth with different bradyrhizobial strains. Environ. Experim. Biol. 31: 285-294.
- Halliday, J, and Graham, P.H. (1978). Coal compared to peat as a carrier of rhizobia. Turrialba 28(4): 348-349.
- Ham, G.E.; Liener, I.E.; Evans S.D.; Frazier, R.D. and Nelson, W.W. (1975). Yield and composition of soybean seed as affected by N and S fertilization. Agron. J. 67: 293-297.
- Hamdi, Y.A. (1970). Soil water tension and the movement of rhizobia. Soil Biol. and Biochem. 3: 121-126.
- Hamdi, Y.A., and Amira AI-Tai, M. (1981). Salt tolerance of strains of *R. meliloti* and *R. trifolii* to chlorides of sodium, calcium and magnesium. Egypt. J. Microbiol. 16(1-2): 1-7.
- Hardarson G., Zapata F., and Danso S.K.A. (1984 b). Effect of plant genotype and nitrogen fertilizer on symbiotic nitrogen fixation by soybean cultivars. Plant and Soil. 82: 397-405.

- Hardarson, G., and, Gareth Jones, D. (1979). Effect of temperature on competition among strains of *Rhizobium trifolii* for nodulation of tow white clover varieties. *Ann. Appl. Biol.* 92: 229-236.
- Harper, J.E., and Cooper, R.L. (1971). Nodulation response of soybeans (*Glycine max* L. Merrill.) to application rate and placement of combined nitrogen. *Crop. Sci.* 11: 438-440.
- Harper, J.E., and Gibson, A.H. (1984). Differential nodulation tolerance to nitrate among legume species. *Crop Sci.* 24: 797-801.
- Hartel, P.G., and Alexander, M. (1984). Temperature and desiccation tolerance of cowpea rhizobia. *Can. J. Microbiol.* 30: 820-836.
- Hassan, M.Z., Abdalla, S., and Nassib, A.M. (1986). Soybean nodulation and yield as affected by *Rhizobium* inoculation, sowing methods and irrigation regime during the early growth stage. *Annals Agric. Sci. Fac. Agric., Ain Shams Univ.* 31(1): 325-336.
- Heckmann, M.O., J.J. Drevon, P. Saglio, and L. Salsac (1989). Effect of oxygen and malate on NO_3^- inhibition of nitrogenase in soybean nodules. *Plant Physiol.* 90: 224-229.
- Helemish, F.A. (1986). Response of rhizobia to saline, alkaline and acid conditions in laboratory media. *J. Bacteriol.* 171: 4543-4548.

Helemish, F.A., and El-Gammal, S.M.A. (1987). Salt and pH tolerance of *Rhizobium leguminosarum* TAL 271. Zentrablatt. Mikrobiol. 142: 211-214.

Hernandez-Armenta, R.; Wien, H.C., and Eaglesham, A.R.J. (1989). Maximum temperature for nitrogen fixation in common bean. Crop. Sci., 29: 1260-1265.

Howieson, J.G., and Ewing, M.A. (1989). Annual species of *Medicago* differ greatly in their ability to nodulate on acid soils. Aust. J. Agric. Res. 40: 843-850.

Howieson, J.G. Ewing, M.A., and D'Antuono, M.F. (1988). Selection for acid tolerance in *Rhizobium meliloti*. Plant and Soil. 105: 179-188.

Hug, S.M.I, and Larther, F. (1983). Effects of NaCl salinity on the growth and nitrogen status of nodulated cowpea (*Vigna sinensis* L.) and mung bean (*Phaseolus aureus* L.). Zeitschrift fur Pflanzen Physiologie. 112(1): 79-87.

Ibrahim, A.N., and Khadr, M.S. (1974). The effect of soil improvement on the nodule formation and plant growth of some legumes crops in saline areas of the Nile delta. Agrochimica es Talajtan 23: 111-120.

Ibrahim, A.N.; Kamel, M., and Khadr, S. (1970). Nodule formation and growth of legumes as influenced by certain sodium salts. Agrochimica es Talajtan, 19(1-2) : 164-172.

- Islam, R., and Afandi, F. (1980).** Responses of lentil cultivars to *Rhizobium* inoculation and nitrogen fertilization. *Lens Newsletter*. 7: 50-51.
- Jackson, M.L. (1962).** Soil chemical Analysis. Perntic Hall, Inc. Engleweeod Cliffs, N.J..
- Jacobsen, E., and Feenstra, W.J. (1984).** A new pea mutant with efficient nodulation in the presence of nitrate. *Plant Sci. Letters* 33: 337-344.
- Kamel, M., Ibrahim, A.N., Hamissa, M.R., and Khadr, M. (1971).** Effect of reclamation of saline soil nodule formation and yield of broad bean. *Agric. Res. Rev.* 49(6): 145-155.
- Karanja, N.K., and Wood, M. (1988 a).** Selecting *Rhizobium phaseoli* strains for use with beans (*Phaseolus vulgaris* L.) in Kenya: Infectiveness and tolerance of acidity and aluminium. *Plant & Soil*. 112: 7-13.
- Karanja, N.K., and Wood, M. (1988 b).** Selecting *Rhizobium phaseoli* strains for use with beans (*Phaseolus vulgaris* L.) in Kenya. Tolerance of high temperature and antibiotic resistance. *Plant and Soil*. 112: 15-22.
- Katoch, K.K., Aggarwal, G.C., and Garg, F.C. (1983).** Effect of nitrogen, soil compaction and moisture stress on nodulation and yield of soybean. I. *The Indian Soc. Soil Sci.* 31(2): 215-219.
- Keyser, H.H., and Munns, D.N. (1979 a).** Effects of Ca, Mn, and Al on growth of rhizobia in acid media. *Soil Sci. Soc. Amer. J.* 43: 500-503.

- Keyser, H.H., and Munns, D.N. (1979 b).** Tolerance of rhizobia to acidity, aluminum and phosphate. Soil Sci. Soc. Amer. J. 43: 51.
- Keyser, H.H., Munns, D.N., and Hohenberg, J.S. (1979).** Acid tolerance of rhizobia in culture and in symbiosis with cowpea. Soil Sci. Soc. Amer. J. 43: 719-722.
- Kishinevsky, B.D., and Weaver, R.W. (1992).** Effect of high root temperature on *Bradyrhizobium*-peanut symbiosis. Plant and Soil 143: 275-282.
- Koermendy, A., and Eaglesham, A.R.J. (1984).** The effect of starter N on soybean nodulation p: 539. In: C veeger and W.E. Newton (ed.) Advances in Nitrogen Fixation Research. Nijhoff, Junk, The Hague.
- Kremer, R., and Harold, L. Peterson (1983).** Effect of carrier and temperature on survival of *Rhizobium* spp. in legume inocula : Development of an improved type of inoculant. Applied and Environmental Microbiology, June, 1983: 1790-1794.
- Kumar, V.; Vimal, K.; Bala, R. and Nafees, N. (1990).** Effect of salt salinity and acidic conditions on growth of cowpea *Rhizobium*. Acta Bacterica Indica.18: 290-292.
- Kuwahara, M., Hoshii, S., and Yashidia, T. (1986).** Soil management and N-fertilization for increasing the yield of soybean in Japan. Technical Bulletin ASPAC Food and Fertilizer Technology Center. No. 98, p. 9.

- La Faver, A.K., and Eaglesham, A.R.J. (1986).** The effect of high temperature on soybean nodulation and growth with different strains of bradyrhizobia. *Can. J. Microbiol.* 32: 22-27.
- La Favre, A.K., and Eaglesham, A.R.J. (1987).** Effects of high soil temperatures and starter nitrogen on the growth and nodulation of soybean. *Crop. Sci.* 27: 742-745.
- Ladha, J.K.; Garcia, M.; Pareek, R.P., and Parivoson, G. (1992).** Relative contributions to nitrogenase (acetylene reducing) activity of stem and root nodules in *sesbania rostrata*. *Can. J. Microbiol.*
- Lakshmi-Kumari, C.S.; Singh, C.S., and Subbarao, N.S. (1974).** Root hair infection and nodulation in lucerne (*Medicago sativa*) as influenced by salinity and alkalinity. *Plant & Soil.* 40: 261-268.
- Lawn, R.J., and Brun, W.A. (1974).** Symbiotic N-fixation in soybeans. III. Effect of supplemental nitrogen and intervarietal grafting. *Crop Sci.* 14: 22-25.
- Lauter, D.J.; Munns, D.N. and Clarkin, K.L. (1981).** Salt response of chickpea as influenced by N supply. *Agron. J.* 73: 961-966.
- Lie, T.A. (1969).** Effects of low pH on different phases of nodule formation in pea plants. *Plant & Soil.* 31: 391-406.
- Lie, T.A. (1971).** Temperature dependent root-nodule formation in pea cv. Iran. *Plant and Soil.* 34: 751-752.

- Lie, T.A. (1974).** Environmental effects on nodulation and symbiotic nitrogen fixation. In: *The Biology of Nitrogen Fixation* (Qisspel A. Ed.). Elsevier. New York, USA, pp. 555-582.
- Lie, T.A. (1981).** Gene Centres, a source for genitic variants in symbiotic nitrogen fixation host-induced ineffectivity in *Pisum sativum* ecotype Fulvum. *Plant and Soil*. 61: 124-134.
- Lie, T.A. (1984).** Host gene in *Pisum sativum* L. conferring resistance to European *Rhizobium leguminosarum* strains. *Plant and Soil*. 82: 415-425.
- Lie, T.A.; D. Goktan and M. Engin (1988).** *Rhizobium* strains from wild and primitive legumes : A nuisance or a valuable gene pool. In : *Nitrogen fixation by legumes in Mediterranean Agriculture* (D.P. Beck and L.A. Materon (editors)) ICARDA, ISBN pp. 121-129.
- Lie, T.A., D. Hille Ris Lambers and A. Houwers (1976).** Symbiotic specialisation in pea plants : Some environmental effects on nodulation and nitrogen fixation pp. 319-333 . In: *Symbiotic Nitrogen Fixation in Plants* (Nutman, P.S. ed.). University Press, Cambridge, England.
- Lindstrom, K.; Sarsa, M.L.; Polkunen, J., and Kansanen, P. (1985).** Symbiotic nitrogen fixation of *Rhizobium* (Galega) in acid soils and its survival in soil under acid and cold stress.
- Lowendorf H.S., and Alexander, M. (1983).** Selecting *Rhizobium meliloti* for inoculation of alfalfa

planted in acid soils. Soil Sci. Soc. Amer. J. 47: 935-938.

Lowendorf, H.S., Bays, A.M., and Alexander, M. (1981). Survival of *Rhizobium* in acid soil. Appl. Environ. Microbiol. 42: 951-957.

Mahler, R.L., and Wollum, A.G. (1981). The influence of soil water potential and soil texture on the survival of *Rhizobium japonicum* and *Rhizobium leguminosarum* isolates in the soil. Soil Sci. of Amer. J. 45: 761-766.

Maier, R.J., and Brill, W.J. (1978). Mutant strains of *Rhizobium japonicum* with increased ability to fix nitrogen for soybean. Science 201: 448-450.

Manhart, J.R., and Wong, P.P. (1980). Nitrate effect on nitrogen fixation (acetylene reduction). Plant Physiol. 65: 502-505.

Marquis, R.E.; Bender, G.R.; Murray, D.R., and Wong, A. (1987). Arginine deiminase system and bacterial adaptation to acid environments. Appl. Environ. Microbiol. 53: 198-200.

Marshall, K.C. (1964). Survival of root-nodule bacteria in dry soils exposed to high temperatures. Aust. J. Agric. Res. 15: 273-281.

Martensson, A.M.; Bautli, L., and Ljunggren, T. (1989). Competition between strains of *Bradyrhizobium japonicum* for nodulation of soybeans at different nitrogen fertilizer levels. Plant & Soil. 117: 219-225.

- Martinez-Romero, E., Segovia, L., and Mercante, F.M. (1991).** Anovel *Rhizobium* species nodulating *Phaseolus vulgaris* L. beans and *Leucaena* sp. trees. *Int. J. Syst. Bacteriol.* 41: 417-426.
- Mary, P.; Ochin, D., and Tailliez, R. (1986).** Growth status of rhizobia in relation to their tolerance to low water activities and desiccation stresses. *Soil Biol. Biochem.* 18: 179-184.
- McNeil, D.L. (1982).** Variation on ability of *Rhizobium japonicum* strains to nodulate soybeans and maintain fixation in the presence of nitrate. *Appl. Environ. Microbiol.* 44: 647-652.
- Minchin, F.R., Minguez, M.I., Sheehy, J.E., Werry, F.J., and Skot, L. (1986 a).** Relationships between nitrate and oxygen supply in symbiotic nitrogen fixation by white clover. *J. Exp. Bot.* 37: 1103-1113.
- Minchin, F.R., Summerfield, R.J., Eaglesham, A.R.J., and Stewart, K.A. (1976).** Effects of short-term water logging on growth and yield of cowpea (*Vigna unguiculata*). *J. Agric. Sci. (Camb.)*. 90: 355-366.
- Mohamed, R.M. Campbell W.F., and Rumbaugh, M.D. (1989 b).** Acetylene reduction activity in salt-tolerant alfalfa *Rhizobium* under drought stress. *Arid Soil Res. Rehab.* 3: 431-438.
- Munever, F., and Wollum, A.G. (1982 b).** Response of soybean plants at high root temperatures as

affected by plant cultivar and *Rhizobium* strain.
Agron. J. 74: 138-142.

Munevar, F., and Wollum, A.G. (1981 a). Growth of *Rhizobium japonicum* strains at temperature above 27 °C. Appl. and Environ. Microbiol. 42: 272-276.

Munevar, F., and Wollum, A.G. (1981 b). Effect of high root temperature and *Rhizobium* strain on nodulation, nitrogen fixation, and growth of soybeans. Soil Sci. Soc. Amer. J. 45 : 1113-1120.

Munns, D.N. (1987). Soil acidity and related factors. p. 211-236. In: J.M. Vincent (ed.). Exploiting the legume rhizobia.

Munns, D.N. (1986). Acid soil tolerance in legumes and rhizobia. Adv. Plant Nutr. 2: 63-91.

Munns, D.N. (1978). Soil acidity and nodulation. p. 247-264. In: C.S. Anrew and E.J. Kamprath (ed.). Mineral nutrition of legumes in soils of the tropics and subtropics. C.S.I.R.O., Melbourne.

Munns, D.N. (1977). Mineral nutrition and the legume symbiosis, In : " A Treatise on dinitrogen fixation. IV. Agronomy and Ecology" R.W.F. Hardy and A. H. Gibson, eds., Wiley Interscience, New York.

Munns, D.N. (1968 b). Nodulation of *Medicago sativa* in solution culture. I. Acid-sensitive steps. Plant and Soil, 28: 129-146.

- Munns, D.N., Fogle, V.M., and Hallock, B.G. (1977). Alfalfa root nodule distribution and inhibition of nitrogen fixation by heat. *Agron. J.* 69: 377-380.
- Nasef, M.A. (1995). Selection of efficient strains of *Rhizobium leguminosarum* with special traits and studying their serological relationship. Ph. D. Thesis, Faculty of Agric., Ain Shams Univ., Cairo, Egypt.
- Nelson, L.M. (1987). Response of *Rhizobium leguminosarum* isolates to different forms of inorganic nitrogen during nodule development in peat (*Pisum sativum* L.). *Soil Biol. Biochem.* 19: 759-763.
- Nelson, L.M. (1983). Variation in ability of *Rhizobium leguminosarum* isolates to fix dinitrogen symbiotically in the presence of ammonium nitrate. *Can. J. Microbiol.* 29: 1626-1633.
- Norris, D.O., and Date, R.A. (1976). Legume bacteriology In Shaw, N.H., and W. Payan (ed.), *Tropical pasture research* pp. 134-174. Commonwealth Agric. Bureaux Bulletin 51, U.K.
- Nortcliff, S. (1988). Soil formation and characteristics of soil profiles. In: Wild, A. (ed.), *Russell's Soil Conditions and Plant Growth*. IIth edition, Longman, London, pp. 168-212.
- Nutman, P.S. (1962). The influence of the legume in root nodule symbiosis. *Roy. Soc. Lond., B.*, 156, 122.

- O'Hara, G.W., Goss, T.J., Dilowrth, M.J., and Glenn, A.R. (1989). Maintenance of interacellular pH and acid tolerance in *Rhizobium meliloti*. Appl. Environ. Microbiol. 55: 1870-1876.
- Osa-Afiana, L.O., and Alexander, M. (1982). Differences among cowpea rhizobia in tolerance to high temperature and desiccation in soil. Appl. Environ. Microbiol. 43: 435-439.
- Ozawa, T., Shima, T., and Yamaguchi, M. (1988). Soil aggregate as a favourable habitat for *Bradyrhizobium japonicum*. Soil Sci. Plant. Nutr. 34: 605-608.
- Pal, U.R. and Saxena, M.C. (1975). Response of soybean to symbiosis and N-fertilization under humid subtropical conditions. Exp. Agric. 11: 221-226.
- Pate J.S., and Dart, P.J. (1961). Nodulation studies in legumes. IV. The influence of inoculum strain on symbiotic response. Plant and Soil 15: 329-346.
- Pate, J.S., and Alkins, C.A. (1963). Nitrogen uptake, transport and utilization. In: Nitrogen Fixation Vol. 3. Legumes. Ed. W.J. Broughton. pp. 245-298. Clarendon Press, Oxford.
- Pena-Cabriaes J., and Alexander M. (1979). Survival of *Rhizobium* in soils undergoing drying. Soil Sci. Soc. of Amer. J. 43: 962-966.
- Philpotts, S.H.C. (1967). The effect of soil temperature on nodulation of cowpea (*Vigna sinensis*). Aust. J. of Exp. Agric. Hus. 7: 372-376.

- Piha, M.I., and Munns, D.N. (1987 a). Nitrogen fixation potential of bean (*Phaseolus vulgaris* L.) compared with other grain legumes under controlled conditions. *Plant and Soil*. 98: 169-182.
- Piha, M.I., and Munns, D.N. (1987 b). Sensitivity of the common bean (*Phaseolus vulgaris* L.) symbiosis to high soil temperature. *Plant and Soil*. 98: 183-194.
- Pijnenborg, J.W.M., Lie, T.A., and Zehnder, A.J.B. (1991). Nodulation of Lucerne (*Medicago sativa* L.) in an acid soil : effects of inoculum size and lime pelleting. *Plant and Soil*. 131: 1-10.
- Rai, R. (1983). The Salt tolerance of *Rhizobium* strains and lentil genotypes and the effect of salinity on aspects of symbiotic N_2 -fixation. *J. Agric. Sci., (Cambridge)*. 100: 81-86.
- Rai, R., and Prasad, V. (1983). Salinity tolerance of *Rhizobium* mutants: Growth and relative efficiency of symbiotic nitrogen fixation. *Soil Biochem*. 15: 217-219.
- Rao, A.V. and Venkateswarlu, B. (1987). Nitrogen fixation as influenced by water stress in selected crop legumes of the Indian arid zone. *Arid Soil. Research and Rehabilitation*. 1: 89-96.
- Rennie, R.J., and Kemp, G.A. (1981). Dinitrogen fixation in *Phaseolus vulgaris* at low temperatures: interaction of temperature, growth

- stage and time of inoculation. Can. J. Bot. 60: 1423-1427.
- Rice, W.A. (1982). Performance of *Rhizobium meliloti* strains selected for low pH tolerance. Can. J. Plant Sci. 62: 941-948.
- Rice, W.A., and Olsen, P.E. (1988). Root - temperature effects on competition for nodule occupancy between two *Rhizobium meliloti* strains. Biol. Fertil. Soils. 6: 137-140.
- Rice, W.A., and Olsen, P.E. (1983). Inoculation of alfalfa seed for increased yield on moderately acid soil Can. J. Soil Sci. 63: 541-545.
- Rice, W.A., Penney, D.C., and Nyborg, M. (1977). Effects of soil acidity on rhizobial numbers, nodulation and nitrogen fixation by alfalfa and red clover. Can. J. Soil Sci. 57: 197-203.
- Richardson, A.E., and Simpson, R.J. (1989). Acid tolerance and symbiotic effectiveness of *Rhizobium trifolii* associated with a *Trifolium subterraneum* L. based pasture growing in acid soil. Soil Biol. Biochem. 21: 87-95.
- Richardson, A.E., and Simpson, R.J. (1988). Enumeration and distribution of *Rhizobium trifolii* under a subterranean clover based pasture growing in an acid soil. Soil Biol. Biochem. 20: 431-438.
- Richardson, A.E., Simpson, R.J., Djordjevic, M.A., and Rolfe, B.J. (1988 b). Expression of nodulation genes in *Rhizobium leguminosarum* bv. *trifolii* is affected by low pH and by Ca and Al ions. Appl. Environ. Microbiol. 54: 2541-2548.

- Rigaud, J. (1981).** Comparison of the efficiency of nitrate and nitrogen fixation in crop yield. In: Nitrogen and carbon metabolism pp. 17-48. Ed. J.D. Bewley. Martinus Mijboff, Dordrecht, The Netherlands.
- Robert, H.B., and Evans, H.J. (1992).** In : Biological Nitrogen Fixation. New York, pp. 475-476.
- Robert, M.Z.; Dennis, D.E. and Glen, H.C. (1981).** Nodulation and N-fixation of field grown California cowpea as influenced by well irrigated and droughted conditions. Agron. J. 73: 9-12.
- Ross, M., Rainbird, Craig, Athkins, and John S. Pate (1983).** Effect of temperature on nitrogenase functioning in cowpea nodules. Plant Physiol. 73: 392-394.
- Rossi, A., and Arst H.N. (1990).** Mutant of aspergillus nidulans able to grow at extremely acid pH acidity the medium less than wild type strains when grown at more moderate pH. FEMS. Microbiol. Letts. 60: 51-54.
- Roughley, R.J., and Dart, P.J. (1970).** Root temperature and root-hair infection of *Trifolium subterraneum* L. cv. cranmore. Plant and Soil. 32: 518-520.
- Roughley, R.J. (1982).** The storage, quality control and use of legume seed inoculants. In: Biological Nitrogen Fixation Technology for Tropical Agriculture. Edited by P.H. Graham and S.C. Harris. CIAT, Cali, Columbia. pp. 115-126.
- Roughley, R.J. and Dart, P.J. (1970).** Root temperature and root-hair infection of *Trifolium subterraneum* L. cv. Cranmore. Plant and Soil. 32: 518-520.

- Roughley, R.J. and Dart, P.J. (1969).** Reduction of acetylene by nodules of *Trifolium subterraneum* as affected by root temperature, *Rhizobium* strain and host cultivar. *Arch. Microbiol.* 69: 171-179.
- Russell, J.S. (1976).** Comparative salt tolerance of some tropical grasses. *Aust. J. Exp. Agric. Anim. Husb.* 16: 103-109.
- Ruschel, A.P., Vose, P.B., Matsui, E., Victoria, R.L., and Tsa Saito, S.M. (1979).** Field evaluation of N_2 -fixation and N utilization by phaseolus bean varieties determined by ^{15}N isotope dilution. *Plant & Soil.* 65: 397-407.
- Salem, S.H.; Gewaily, E.M., and El-Zamik, F.I. (1981).** Effect of soil salinity on the symbiosis of root nodule bacteria and broad bean plants. *Agrok. Talajan.* 30(1-2).
- Salema, M.P.; Parker, C.A.; Kidby, D.K.; Chatel, D. and Armitage, T.M. (1982).** Rupture of nodule bacteria on drying and rehydration. *Soil Biol. Biochem.* 14: 15-22.
- Sauvage, D., Hamelin, J., and Larher, F. (1983).** Glycine betain and other structurally related compounds improve the salt tolerance of *Rhizobium meliloti*. *Plant Sci. Lett.* 31: 291-302.
- Segovia, L.; Pinero, D.; Palacios, R. and Martinez-Romero, E. (1991).** Genetic structure of a soil population of nonsymbiotic *Rhizobium leguminosarum*. *Appl. Environ. Microbiol.* 57: 426-433.

- Semu, E., Hume, D.J. and Cokke, C.T. (1979). Influence of soybean inoculation and nitrogen levels on populations and zero groups of *Rhizobium japonicum* in Ontario. Can. J. Microbiol. 25: 739-745.
- Senaratine, R.; Amornpimol, C. and Hardarson, G. (1987). Effect of combined nitrogen on nitrogen fixation of soybean (*Glycine max* L. Merr.) as affected by cultivar and rhizobial strain. Plant & Soil. 103: 45-50.
- Serraj, R.; Dervon, J.J.; Obaton, M. and Vidal, A. (1992). Variation in nitrate tolerance of nitrogen fixation in soybean (*Glycin max*)-*Bradyrhizobium* symbiosis. Plant Physiol. 140: 366-371.
- Sidiqui, S. and Kumar, S. (1987). Note on nodule metabolism under salinized and desalinized conditions in pea c.v. Banneville. Current Agriculture 11(1-2) 69-72.
- Simpson, J.R. and Gibson, A.H. (1970). Effect of combined nitrogen levels on nodulation of legumee. Soil Biol. Biochem. 2: 295.
- Singelton, P.W. and Bohlool, B.B. (1984). Effect of salinity on nodule formation by soybean. Plant Physiol. 74(1): 72-76.
- Singelton, P.W., (1983). A split-root growth system for evaluating the effect of salinity on the components of the soybean *Rhizobium japonicum* symbiosis. Crop. Sci. 23: 259-262.

Singleton, P.W., and Bohlool, B.B. (1983). Effect of salinity, on the functional components of the soybean - *Rhizobium japonicum* symbiosis. Crop Sci. 23: 815-818.

Singleton, P.W.; El-Swaily, S.A. and Bohlool, B.B. (1982). Effect of salinity on *Rhizobium* growth and survival. Appl. and Environ. Microbiol. 44(4): 884-890.

Sirry, A.R.; S.H. Salem; E.M. El-Gewaily and F.J. El-Zamik (1980). Effect of soil reclamation on the symbiotic relationship between *Rhizobium* and some leguminous plants. In: Proc. of the Intern. Symp. on salt-affected soils, Kernak, India, pp. 461-467.

Smith, D.L. and Hume D.J. (1985). Effect of irrigation and fertilizer N on $N_2(C_2H_2)$ fixation and Yield of white bean and soybean. Can. J. Plant Sci. 65: 307-316.

Smith, R.S., and Del Rio Esturra, G.A. (1982). Soybean inoculant types and rates evaluated under dry and irrigated field conditions. J. Agric. Univ. P.R. 66: 241-249.

Smith, R.S., Ellis, M.A., and Smith, R.E. (1981). Effect of *Rhizobium japonicum* inoculant rates on soybean nodulation in a tropical soil. Agron. J. 73: 505-508.

Snedecor, G.W. and Cochran, W.G. (1980). Statistical Methods. Iowa State Univ. Press, U.S.A.

Somasegran, P., Reyes, V.G., and Holen, H.J. (1984).

The influence of high temperatures on the growth and survival of *Rhizobium* spp. in peat inoculants during preparation, stage, and distribution. *Can. J. Microbiol.* 30: 23-30.

Soos, T., Balint, H. and Konya, K. (1984). Effect of Rhizonit "S" inoculum irradiated with ^{60}Co isotope and nitrogen fertilizers on N-fixation by soybean plants. *Soil Biol. and Conservation on the Biosphere* 2: 495-501. (ed. by Szegi, J. Budapest, Hungary).

Sprent J.I. (1984). Effect of drought and salinity on heterotrophic nitrogen fixing bacteria and on infection of legumes by rhizobia. In: *Advances in Nitrogen Fixation Research*, Pp. 295-302. EdC Veeger and WE Newton. Martinus Nijhoff, The Hague.

Sprent, J.I (1975). Nitrogen fixation by legumes subjected to water and light stresses. In *symbiotic Nitrogen Fixation in plants* "(P.S. Nutman ed.). Cambridge Univ. Press, London and New York, 405-420 .

Sprent, J.I. (1972 a). The effects of water stress on nitrogen fixing root nodules. IV- Effects on whole plants of *Vicia faba* and *Glycine max.* *New Phytol.* 71: 603-611.

Sprent J.I. (1971 b). Effect of water stress on nitrogen fixation in root nodules. *Plant and Soil, Spec.* 225-228.

- Sprent, J.I. (1969).** Prolonged reduction of acetylene by detached soybean nodules. *Planta*, Berlin, 88: 372-375.
- Sprent J.I. and Zahran H.H. (1988).** Infection, development and functioning of nodules under drought and salinity. In: *Nitrogen Fixation by Legumes in Mediterranean Agriculture*, pp. 145-151, Eds DP Beck and LA Materon. Martinus Nijhoff/Dr. W. Junk, Dordrecht.
- Stephens, B.D. and C.A. Neyna (1983).** Nitrate and nitrite reduction in relation to nitrogenase activity in soybean nodules and *Rhizobium japonicum* bacteroids. *Plant Physiol.* 71: 731-735.
- Streeter, J.G. (1988).** Inhibition of legume nodule formation and N_2 fixation by nitrate *CRC Crit. Rev. Plant Sci.* 7: 1-23.
- Subba Rao, N.S. (1977).** Soil Micro organisms and plant growth. Oxford and IBH Publishing Company, New Delhi, India.
- Subba Rao, G.W.; Johnsen, C.; Kumar Rao, J.V.D.K.; and Jana M.K. (1990).** Response of the pigeonpea - *Rhizobium* symbiosis to salinity stress: Variation among *Rhizobium* strain symbiotic ability. *Fertil Soils.* 9: 49-53.
- Swelim, D.M. (1996).** Environmental and Nutritional studies on *Rhizobium* spp. (*Leucaena*). Ph.D. Thesis. Fac. Agric. Cairo Univ.
- Sylvester-Bradley, R.; Mosquera, D. and Mendez, I.E. (1988).** Selection of rhizobia for inoculation of

forage legumes in savanna and rainforest soils of tropical America. In: Nitrogen Fixation by Legumes in Mediterranean Agriculture. Edited by D.P. Beck and L.A. Materon Martinus Nijhoff, Dordrecht, The Netherlands. pp. 225-233.

Tanaka, A., Yamauchi, J., Fujita, K., Koto, H. and Uroyama, M. (1984). Yield of soybeans as influenced by genetic characteristics, climatic conditions, and nitrogen nutrition. Soil Sci. and Plant Nutrition. 30 (4): 533-541.

Tang, C. , and Robson, A. (1993). The pH above 6.0 reduces nodulation in Lupinus species. Plant & Soil. 152: 269-276.

Tanner, J.W. and Anderson, I.C.C. (1963). Effect of nitrogen on nodulation and yield of irrigated soybeans. Can. J. Plant Sci. 43: 542.

Taylor, R.W., Williams, M.L., and Sistrani, K.R. (1991). N₂ fixation by soybean - *Bradyrhizobium* combinations under acidity. low P and high Al Stresses. Plant & Soil. 131: 293-300.

Terraccino, J.S.; Rapaport and Kashket (1988). Stress and growth phase proteins of *Clostridium acetobutylicum*. Appl. Environ. Microbiol., 54 : 1989-1995.

Thornton, H.G. (1936). Atmospheric nitrogen fixing by legumes. Proceedings of the Royal Society (London). 119-474.

Toomsan, B. (1990). Groundnut microbiology research at Khon Kaen University. In: Patanothai, A. (ed),

Groundnut Improvement Project at Khon Kaen University. Report of Work for 1986-1988. Khon Kaen University. Thailand, pp. 89-111.

Toomsan B., Rupela, O.P. and Dart, P.J. (1982). Studies on soil and rhizosphere populations of *Rhizobium* sp. nodulating *Cicer arietinum* In: Proceedings of the National Symposium on Biological *Cicer arietinum* In: Proceedings of the National Symposium on Biological Nitrogen Fixation. IARI, Delhi, pp. 517-531.

Toro, N. (1996). Nodulation competitiveness in the *Rhizobium* - legume symbiosis. World J. Microbiol. Biotechnology. 12: 157-162.

Tu, J.C. (1981). Effect of salinity on *Rhizobium* root-hair interaction, nodulation and growth of soybean. Can. J. of Plant Sci. 61(2): 231-239.

Valera, C.L., and Alexander, M. (1965). Effect of inoculum, nitrogen and phosphorus on root nodulation and yield of soybean varieties. Nature. 206: 326.

Van Rensberg, H.J. and Strijdom, B.W. (1980). Survival of fast and slow-growing *Rhizobium* spp. under conditions of relatively mild desiccation. Soil Biol. Biochem. 12: 353-356.

Van Schreven, D.A. (1972). On the resistance of effectiveness of *Rhizobium trifolii* to a low pH. Plant & Soil. 37: 49-55.

Van Schreven, D.A. (1958). In: Nutrition of the legumes (Ed. E.G. Hallsworth), Proc. Univ. Nottingham 5th

Easter School in Agricultural Science,
Butterworths, London, P. 328.

Vargas, A.A.T and Graham P.H. (1988). *Phaseolus vulgaris* cultivar and *Rhizobium* strain variation in acid-pH tolerance and nodulation under acid conditions. *Field Crops Res.* 19: 91-101.

Vasilas, B.L., Walker, W.M. and Ham, G.E. (1984). Dry matter and primary nutrient accumulation in soybeans as affected by combined nitrogen levels. *J. Plant Nutrition.* 7(12): 1731-1743.

Velgaleti, R.R., and Marsh S. (1989). Influence of host cultivars and *Bradyrhizobium* strains on the growth and symbiotic performance of soybean under salt stress. *Plant & Soil.* 119: 133-138.

Velagaleti, R.R., Marsh, S., and Kramer, D. (1990). Genotypic differences in growth and nitrogen fixation among soybean (*Glycine max* (L.) Merr) cultivars grown under salt stress. *Trop. Agric. (Trinidad),* 67: 169-177.

Vincent, J.M. (1970). A manual for the practical study of root nodule bacteria. International Biological programme handbook No. 15. Blackwell Publications. Ltd., Oxford.

Weaver, R.W., and Holt, E.C. (1990). Short-term survival of rhizobia on arrowleaf clover seed at different depths. *Plant & Soil.* 122: 147-150.

Weber, D.F., and Miller, V.L. (1972). Effect of soil temperature on *Rhizobium japonicum* serogroup

distribution in soybean nodules. Agron. J. 64: 796-798.

Wilkins, J. (1967). The effects of high temperatures on certain rootnodule bacteria. Aust. J. Agric. Res. 18: 299-304.

Wilson, D.O. and Trang, K.M. (1980). Effects of storage temperature and enumeration method on *Rhizobium* spp. numbers in peat inoculants. Trop. Agric. (Trinidad). 57: 233-238.

Wood, M., and Shepherd, G. (1987). Characterization of *Rhizobium trifolii* isolated from soils of different pH. Soil Biol. Biochem. 19: 317-321.

Wood, M.; Cooper, H.E. and Bjourson, A.J. (1988). Response of *Lotus* rhizobia to acidity and aluminum in liquid culture and in soil. Plant and Soil. 107: 227-231.

Worral, V.S., and Roughley, R.J. (1976). The effect of moisture stress on infection of *Trifolium subterranean* L., by *Rhizobium trifolii* Dang. J. Exp. Bot. 27: 1233-1241.

Wych, R.O. and D.W. Rains (1979). Nitrate absorption and acetylene reduction by soybeans during reproductive development. Physiol. Plant. 47: 200-204.

Yadov, N.K. and Vyas, S.R. (1971). Response of root nodule rhizobia to saline, alkaline and acid conditions. Indian J. Agric. Sci. 41: 875-881.

- Yelton, M.M.; Yang, S.S.; Edie, S.A. and limp, S.T. (1983). Characterization of an effective salt-tolerant fast-growing strain of *Rhizobium japonicum*. J. Gen. Microbiol. 129: 1537-1547.
- Zahran, H.H. and Sprent I.T (1986). Effects of sodium chloride and polyethylene glycol on root-hair infection and nodulation of *Vicia faba* L. plants by *Rhizobium leguminosarum*. Planta . 167: 303-309.
- Zhang, X., Harper, F., Karsisto, M., and Lindstrom, K. (1991). Diversity of *Rhizobium* bacteria isolated from the root nodules of leguminous trees. Int. J. Syst. Bacteriol. 41: 104-113.