

REFERENCES

- Abdeen, S.A.O. (1985).
Some new approaches to control cotton bollworms.
Ph.D. Thesis, Fac. of Agric., Al-Azhar University.
- Abd El-Hafez, A. (1982).
Further studies on the ecology and behaviour of the
pink bollworm, Pectinophora gossypiella (Saund.).
Ph.D. Thesis, Fac. of Agric., Ain Shams University.
- , A.G. Metwally and M.R.A. Saleh (1982).
Rearing pink bollworm, Pectinophora gossypiella
(Saunders) on kidney bean diet in Egypt (Lepidoptera :
Gelechiidae).
Res. Bull., Fac. Agric., Zagazig Univ., No. 576, 10 pp.
- Abdel-Hafez, M.M.; M.M. Abd El-Sattar and M.A. El-Malla
(1984).
Changes in esterases, phosphatases and amino acid
transferases of the pink bollworm moth during the
course of insecticides poisoning.
Bull. ent. Soc. Egypt, Econ. Ser., 14, p. 429.
- Adkisson, P.L. (1961).
Fecundity and longevity of the adult female pink boll-
worm reared on natural and synthetic diets.
J. Econ. Entomol., 54 (6) : 1224-1227.
- (1962).
Timing of defoliants and desiccants to reduce popula-
tions of the pink bollworm in diapause.
J. Econ. Entomol., 55 : 949-951.
- (1964).
Action of the photoperiod in controlling insect
diapause.
The American Naturalist, Vol. XCVII, No. 902, Sept.
1964.

- Adkisson, P.L. (1966).
Internal clocks and insect diapause.
Reprinted from Science, Vol. 154, No. 3746 : 234-241.
- and J.C. Gaines (1960).
Pink bollworm control as related to the total cotton
insect control program of Central Texas.
Texas Agric. Expt. Sta. Misc. Publ., 444.
- and S.G. Wellso (1962).
Effect of DDT poisoning on the longevity and fecun-
dity of the pink bollworm.
J. Econ. Entomol., 55 (6) : 842-845.
- Ahmed, S.M., M.K. El-Khishin, M.A. Rashad and S.M. Nagieb
(1987).
Relative susceptibility to different insecticides and
effect of their sublethal doses on the full-grown
larvae of the pink bollworm Pectinophora gossypiella
(Saunders).
Ann. Agric. Sci., Moshtohor, 25 (2) : 1899-1908.
- , F.F. Shalaby and M.M. Assar (1980).
Primary studies on the effect of cotton root bark and
gossypol on egg laying and fertility of eggs of house-
fly Musca domestica L. (Diptera : Muscidae).
Ann. Agric. Sci., Moshtohor, 13 : 115-121.
- Allam, S., K. Gohar, A.S. Kansouah and Z.H. Zidan (1978).
The effect of sublethal doses of cyolane, Dursban
and Lannate on certain biological aspects and
potentiality of Spodoptera littoralis (Boisd.).
4th Conf. Pest Control, NRC, Cairo, 1978.
- Andrewatha, H.G. (1952).
Diapause in relation to the biology of insects.
Biol. Rev., 27 : 50-107.
- and L.C. Birch (1954).
The distribution and abundance of animals.
The University of Chicago Press, Chicago, Illinois.

- Bailey, A.E. (1948).
Cottonseed and cottonseed products. The Chemistry
and chemical technology.
Interscience Publishers, Inc., New York, U.S.A.
- Bariola, L.A., D.L. Kittock, H.F. Arle, P.V. Vail and T.J.
Henneberry (1976).
Controlling pink bollworm : Effect of chemical
termination of cotton fruiting on populations of
diapausing larvae.
J. Econ. Entomol., 69 (5) : 633-636.
- Barry, B.D. and P.L. Adkisson (1966).
Certain aspects of the genetic factors involved in
the control of larval diapause of the pink bollworm.
Ann. Ent. Soc. Amer., 59 (1) : 122-125.
- Belal, A.H., F.A. Adal, M.M. Essaway and M.A. Khalifa (1988).
The role of bound free amino acids and transaminases
(alanine and asparatic) in regulating the resistance
mechanism of Spodoptera littoralis (Boisd.) to
cypermethrin and profenofos.
Proc. 3rd Egypt. Conf. Agron., Kafr El-Sheikh,
5-7 September.
- Bell, C.H. (1983).
The regulation of development during diapause in
Ephestia elutella (Hubner).
J. Insect Physiol., 29 (6) : 485-490.
- Benz, G. (1963).
Physiopathology and histochemistry in insect pathology.
(ed. by Steinhauser), Vol. 1 : 299-338. Academic
Press, New York and London.
- Bottger, G.T., E.T. Sheehan, and Lukefahr.
Relation of gossypol of cotton plants to insect
resistance.
J. Econ. Entomol., 57 : 283-285.

- Brazzel, J.R. and L.D. Newsom (1959).
Diapause in Anthonomus grandis Boh.
J. Econ. Entomol., 52 (4)
- Brazzel, J.R. and D.F. Martin (1957).
Oviposition sites of the pink bollworm on cotton.
J. Econ. Entomol., 50 (2) : 122-124.
- Bull, D.L. and P.L. Adkisson (1962).
Fat content of the larval diet as a factor influencing diapause and growth rate of the pink bollworm.
Ann. Ent. Soc. Amer., 55 (5) : 499-502.
- Chapman, A.J., L.W. Nobel, O.T. Robertson and L.C. Fif (1960).
Survival of the pink bollworm under various cultural and climatic conditions.
USDA Prod. Res. Rep., No. 34, 21 pp.
- Chautbani, A.R. and P.L. Adkisson (1966).
Effect of sublethal doses of certain insecticides on eggs, larvae, and adults of two species of Heliothis.
J. Econ. Entomol., 59 (3) : 1070-1074.
- Chen, P.S. (1971).
Biochemical aspects of insect development. pp. 56-56.
S. Karger, Basel.
- Chippendale, G.M. and S. Kikukawa (1983).
Effect of day length and temperature on the larval diapause of the sunflower moth, Homoeosoma electellum.
J. Insect Physiol., 29 (8) : 643-649.
- El-Sayed, M.A.R. (1970).
The relation between the gossypol content of cotton varieties and its toxicity to insects.
M.Sc. Thesis, Fac. of Agric., Cairo Univ.
- El-Sayed, M.T. and Z.M.F. Rustom (1960).
Factors affecting the initiation of diapause in the pink bollworm, Pectinophora gossypiella (Saunders).
Bull. ent. Soc. Egypté, 44 : 253-264.

Fife, L.C. (1949).

Studies of diapause in the pink bollworm in Puerto Rico. V.S. Dept.
Agric. Tech. Bull., 977 : 26 pp.

----- (1961).

Factors influencing pink bollworm pupation and moth emergence from overwintering larvae in central Texas. J. Econ. Entomol., 54 : 908-913.

Finney, D.J. (1971).

Probit Analysis.
Cambridge Univ. Press, London, 3rd ed., 333 pp.

Firling, C.E. (1977).

Amino acid and protein changes in the haemolymph of developing fourth instar Chironomus tentans. J. Insect Physiol., 23 : 17-22.

Flint, H.M. and R.L. Smith (1977).

Laboratory evaluation of TH 60-40 against the pink bollworm.
J. Econ. Entomol., 70 (1)

Gohar, K., Z.H. Zidan, A. Kansouah and S. Allam (1978).

The effect of sublethal doses of Dursban, Cyolane and Lannate on the amino acids and fat contents of the larvae of the cotton leafworm, Spodoptera littoralis (Boisd.).

4th Conf. Pest Control, NRC, Cairo, 1978.

Gomaa, E.A.A., M.M.I. Amir and H.H. Metaweh (1987).

Morphogenetic activity of dithamathrin on the desert locust Schistocerca gregaria (Forst), and biology of malformed adults.

J. Egypt Soc. Ent. (under publication).

- Gong, X.W., and G.L. Meng (1987).
Preliminary observations on the bionomics of pupation in the first generation of Pectinophora gossypiella (Saund.).
Insect Knowledge, 21 (5) : 208-210.
- Graham, H.M., P.A. Glick and M.T. Ouye (1967).
Temperature effect on reproduction and longevity of laboratory-reared adult pink bollworm (Lepidoptera: Gelechiidae).
Ann. Ent. Soc. Amer., 60 : 1211-1213.
-
- and D.F. Martin (1965).
Mating frequency of female pink bollworm collected from light traps.
Ann. Ent. Soc. Amer., 58 (4) : 595
- Gutierrez, A.R.P., M.A. Santos, W.J. Dos, A. Villacorta and K.D. Gallagher (1986).
Analysis of diapause induction and remination in Pectinophora gossypiella in Brazil.
Environ. Entomol., 15 (3) : 494-500.
- Hasegawa, K. and Yamashita, O. (1965).
Studies on the mode of action of the diapause hormone in the silkworm, Bombyx mori L. VI. The target organ of the diapause hormone.
J. Exp. Biol., 43 : 271-277.
- Hassanein, M.H. and A. Galal (1969).
Studies on the diapause of the pink bollworm, Pectinophora gossypiella (Saund.).
Bull. Soc. Ent. Egypté, 53 : 69-78.
- Henneberry, T.J., H.M. Flint and L.A. Bariola (1977).
Temperature effects on mating sperm transfer oviposition, and egg viability of pink bollworm.
Environ. Entomol., 6 (4)

- Harper, H.A., V.W. Rodwell and P.A. Mayer (1977).
The blood, lymph & cerebrospinal fluid.
Review of Physiological Chemistry, 17th ed. Lange
Medical Publication, Los Altos, California, U.S.A.
Middle East Edition, p. 188-216.
- Highnam, K.C. and L. Hill (1977).
The comparative endocrinology of the invertebrates.
Edward Arnold (Publishers) Ltd., England, 2nd ed.,
346 pp., ELBS edition.
- Hinds, W.E. and W.W. Yothers (1909).
Hibernation of the Mexican cotton boll weevil.
U.S. Dept. Agric. Bur. Ent. Bull., 77.
- Hunter, W.D. and W.E. Hinds (1904).
The Mexican cotton boll weevil.
U.S. Dept. Agric. Bur. Ent. Bull. 45.
- and W.D. Pierce (1912).
Mexican cotton boll weevil.
U.S. Senate Document No. 305.
U.S. Dept. Agric. Bur. Ent. Bull. 114.
- Ingram, R.L. (1955).
Water loss from insects treated with pyrethrins.
Ann. Ent. Soc. Amer., 48 : 481-485.
- Jeffery, P.L., L.A. Crowder and J.F. Watson (1974).
Amino acids and total nitrogen of diapause and non-
diapause larvae of the pink bollworm, Pectinophora
gossypiella.
Ann. Entomol. Soc. Amer., 67 (3)
- Kansouah, A., Z.H. Zidan, K. Gohar and S. Allam (1978).
Amino acids and fat contents of the pupal stage of
the cotton leafworm, Spodoptera littoralis (Boisd.)
as affected by sublethal doses of Dursban, Cyolane
and Lannate.
4th Conf. Pest Control, NRC, Cairo (1978).

Kuipers, J. (1962).

Her effect van sublethale op de epiroductie van de coloradokafer (Leptinotarsa decemlineata Say.).

Mededel Landbouwhogeschool Wageningen, 27 (3) : 908-919.

Laufer, H. (1960).

Blood protein in insect development.

Ann. N.Y. Acad. Sci., 89 : 490-515.

Loftin, U.C., K.B. McKinney and W.K. Hanson (1921).

Report on investigations of the pink bollworm of cotton in Mexico.

U.S. Dept., Agric. Bull., 918 pp.

Lowry, D.H., N.J. Rosebrough, A.L. Farr and H.J. Randall (1951).

Protein measurement with the folin phenol reagent. J. Biol. Chem., 176 : 367-388.

Lukefahr, M.J. and D.F. Martin (1966).

Cotton pigments as a source of resistance to the bollworm and tobacco budworm.

J. Econ. Entomol., 59 : 176-179.

-----, L.W. Noble and D.F. Martin (1964).

Factors inducing diapause in the pink bollworm.

Tech. Bull. U.S. Dept. Agric., 1304 (1+) 17 pp.

Mauney, J.R., D.L. Kitlock and L.A. Bariola (1972).

Limitation of late season green bolls : A possible new control for pink bollworms.

Proc. 26th Ann. Cotton Defoliation and Physiology Conf., Memphis, TN, 26 : 103-106.

Menaker, M. and G. Gross (1965).

Effect of fluctuation temperature on diapause induction in the pink bollworm.

Zool. Dept., Univ. of Texas, Austin, Texas.

- Metwally, A.G. (1961).
Studies on the pink bollworm, Pectinophora gossypiella (Saund.).
M.Sc. Thesis, Fac. of Agric., Ain Shams Univ.
- , S.M. Ahmed, Rashad M. Amira and S.M. (1987).
Effect of larvae exposure to sublethal doses of insecticides on pupae and adults of Pectinophora gossypiella (Saund.).
Ann. Agric. Sci. Moshtohor, 25 (2) : 1911-1920.
- Moore, R.F. (1983).
Effect of dietary gossypol on the boll weevil (Coleoptera : Curculionidae).
J. Econ. Entomol., 76 (4) : 696-699.
- Morgan, T.D. and G.M. Chippendale (1983).
Free amino acids of the haemolymph of the southwestern corn borers and the European corn borer in relation to their diapause.
J. Insect Physiol., 29 (10) : 735-740.
- Ohlendorf, W. (1926).
Studies of the pink bollworm in Mexico.
U.S. Dept. Agric. Bull., 1374.
- Omar, A.M.A. (1979).
Further studies on the effect of dietary lipids on the induction of diapause in the pink bollworm Pectinophora gossypiella (Saund.) (Lepidoptera : Gelechiidae).
Ph.D. Thesis, Fac. of Agric., Cairo University.
- Ouye, M.T. and H. Knutson (1957).
Reproductive potential, longevity and weight of house-flies following treatments of larvae with Malathion.
J. Econ. Entomol., 50 (4)

- Ouye, M.T., H.M. Graham, C.A. Richmond, and D.F. Martin
(1964).
Matin studies of the pink bollworm.
J. Econ. Entomol., 57 (3) : 222-225.
- Pearson, E.O. (1958).
The insect pests of cotton in Tropical Africa.
Commonwealth Inst. Ent. (London), 355 pp.
- Pons, W.A., G.L. Hoffpauir and T.H. Hopper (1953).
Influence of variety of cotton seed and environment
on its gossypol content.
J. Agr. & Food Chem., 1 : 1115-1118.
- Raina, A.K. (1974).
A non-diapause strain of pink bollworm from southern
India.
Ann. Ent. Soc. Amer., 67 : 685-686.
- and R.A. Bell (1976).
Occurrence of larval-pupal intermediates in hybrid
pink bollworms reared at low temperatures.
Ann. Ent. Soc. Amer., 69 (2) : 290-292.
- Reynolds, J.E.F. (1982).
Martindale.
The extra Pharmacopoeia Edition 28, published by
the Societ's Department of Pharmaceutical Sciences,
London, The Pharmaceutical press.
- Richmond, C.A., H.M. Graham and C.T. Perez (1972).
Temperatures for storing pupae and adults of the
pink bollworm.
J. Econ. Entomol., 65 (2)
- Rodwell, V.W. (1977).
Biosynthesis, catabolism of nitrogen and catabolism
of carbon skeletons of amino acid.
Review of Physiological Chemistry, p. 384-429.
Edited by Harper, H.A., V.W. Rodwell and P.A. Mayer,
17th ed., Middle East Edition, 702 pp.

Rustom, Z.M.F. (1960).

The chemical composition of the larvae of the pink bollworm, P. gossypiella (Saund.) before and during diapause.

Bull. Soc. ent. Egypté, 44 : 421-449.

Shaver, T.N. and M.J. Lukefahr (1969).

Effect of flavonoid pigments and gossypol on growth and development of the bollworm, tobacco, bollworm and pink bollworm.

J. Econ. Entomol., 62 (3)

Shotwell, O.S., G.A. Bennett, H.H. Hall, R.D. Stubblefield, J. Peters, C.H. Van Etten and R.W. Jackson (1965).
Amino acids in the haemolymph of diseased Popillia japonica (Newman) larvae.

J. Insect Physiol., 11 : 671-682.

Smith, R.F. (1971).

Juveno-memetic effects of Lannate in sublethal doses on western spruce budworm Choristoneura occidentalis.
J. Insect Path., 17 : 132-133.

Staal, J.B. (1972).

Biological activity and bioassay of juvenile hormone analogues.

In Merm, J. and Beroza, M. (ed.) : Insect Juvenile Hormone, Academic Press, New York & London.

Squire, F.A. (1937).

A theory of diapause in Platydera gossypiella (Saund.).
Trop. Agr., 14 : 299-301.

----- (1940).

On the nature and origin of the diapause in Platydera gossypiella (Saund.).

Bull. Entomol. Res., 31 : 1-6.

- Sukhiya, H.S. and G.J. Reddy (1983).
Time of initiating insecticidal application for the control of Pectinophora gossypiella (Saunders).
J. Res. Punjab Agric. Univ., 20 (1) : 47-56.
(R.A.E., 72 (1) : 3, 1984).
- Tsao, C.H. and W.L. Lowry (1963).
Insecticidal control of first instar pink bollworm larvae and observation of their dispersal on cotton plants.
J. Econ. Entomol., 56 (3) : 370-372.
- Taylor, T.H.C. (1936).
Report on a years investigation of Platydera gossypiella (Pink bollworm) in Uganda.
Ann. Rept. Uganda Dept. Agric., 1935-36 (2) : 19-33.
- Thomas, S.A. (1933).
Prothetely in an Elaterid larva (Ceoleop.).
Entomol. News, 44 : 91-96.
- Toyama, K. (1913).
Maternal inheritance and mendelism.
J. Genetics, 2 : 351-405.
- Vanderzant, E.S. (1965).
Axenic rearing of the boll weevil on defined diets : Amino acid, carbohydrate and mineral requirements.
J. Insect Physiol., 11 : 659-670.
- Wellso, S.G. and P.L. Adkisson (1961).
Gross morphology of the female reproductive organs of the pink bollworm.
Presented at the Meetings of the Southwestern Branch, Ent. Soc. Amer., at San Antonio, Texas, Feb. 20-21.
- Williams, C.M. (1952).
Physiology of insect diapause. IV. The brain and prothoracic glands as an endocrine system in the cercopia silk worm.
Biol. Bull. Woods Hole, 103 : 102-238.