

CONTENTS

I-Introduction.....	1
II-Review of Literature.....	5
1-Efficacy of controlled atmospheres of various carbon dioxide concentrations on stored product pests.....	5
2-Efficacy of controlled atmospheres of very high nitrogen content/or pure nitrogen on stored product pests.....	34
3-Efficacy of phosphine on stored product pests.....	37
4-Efficacy of combinations of phosphine and carbon dioxide against stored product- pests.....	66
III-Materials and Methods.....	77
-Storage facilities.....	77
a) Airtight steel bins.....	77
b) Closed metal drums (Barrels).....	77
c) Closed fiberglass bins.....	77
-Insects.....	81
-Insect cultures.....	81
-Cultures for supplying the developmental insect stages, required for the tests.....	81
-Methods of obtaining the developmental stages of the various insect species.....	82
-Preparation of the test-insects for various treatments.....	83
-Gases used.....	84
-Exposure procedure of the insects inside the bins.....	84
-Purging of the gases inside the bin.....	85

II

-Determination of gas concentrations.....	85
-Bioassay tests.....	86
-Calculation of the joint action of the combined treatments.....	86
IV-Results and Descussions.....	90
1-Efficacy of CA of 30% CO ₂ against the tested insect species at 18 ± 2 ⁰ C and 55 ± 5% R.H. inside the various storage facilities.....	90
1a) Efficacy of CA of 30 ± 5% CO ₂ against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed steel bins at varying exposure periods.....	90
1b) Efficacy of CA of 30 ± 5% CO ₂ against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed fiberglass bins at varying exposure periods.....	93
1c) Efficacy of CA of 30 ± 5% CO ₂ against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed metal drums at varying exposure periods.....	96
2-Efficacy of CA of 30% CO ₂ against the tested insect species at 28 ± 2 ⁰ C and 55 ± 5% R.H. in the various storage facilities.....	99
2a) Efficacy of CA of 30 ± 5% CO ₂ against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed steel bins at varying exposure periods.....	99
2b) Efficacy of CA of 30 ± 5% CO ₂ against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed fiberglass bins at varying exposure periods.....	102
2c) Efficacy of CA of 30 ± 5% CO ₂ against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed metal drums at varying exposure periods.....	105

IV

-Loss of 60% CO ₂ concentrations inside the various storage facilities at different exposure periods during summer and winter times.....	128
5-Efficacy of 100 ppm of phosphine against the tested insect species at 18 ± 2 ⁰ C and 55 ± 5% R.H. inside the various storage facilities.....	133
5a) Efficacy of 100 ppm of phosphine against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed steel bins at varying exposure periods.....	133
5b) Efficacy of 100ppm of phosphine against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed fiberglass bins at varying exposure periods.....	137
5c) Efficacy of 100 ppm of phosphine against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed metal drums at varying exposure periods.....	140
6-Efficacy of 100 ppm of phosphine against the tested insect species at 28 ± 2 ⁰ C and 55 ± 5% R.H. inside the various storage facilities.....	143
6a) Efficacy of 100 ppm of phosphine against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed steel bins at varying exposure periods.....	143
6b) Efficacy of 100 ppm of phosphine against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed fiberglass bins at varying exposure periods.....	146
6c) Efficacy of 100 ppm of phosphine against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed metal drums at varying exposure periods.....	149

III

-Loss of 30% CO ₂ concentrations inside the various storage facilities at different exposure periods during summer and winter times.....	108
3-Efficacy of CA of 60% CO ₂ against the tested insect species at 18 ± 2 ⁰ C and 55 ± 5% R.H. in the various storage facilities.....	108
3a) Efficacy of CA of 60 ± 5% CO ₂ against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. in the closed steel bins at varying exposure periods.....	108
3b) Efficacy of CA of 60 ± 5% CO ₂ against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed fiberglass bins at varying exposure periods.....	113
3c) Efficacy of CA of 60 ± 5% CO ₂ against the tested insect species at 18 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed metal drums at varying exposure periods.....	116
4-Efficacy of CA of 60% CO ₂ against the tested insect species at 28 ± 2 ⁰ C and 5 ± 5% RH inside the various storage facilities.....	119
4a) Efficacy of CA of 60 ± 5% CO ₂ against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed steel bins at varying exposure periods.....	119
4b) Efficacy of CA of 60 ± 5% CO ₂ against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed fiberglass bins at varying exposure periods.....	122
4c) Efficacy of CA of 60 ± 5% CO ₂ against the tested insect species at 28 ± 2 ⁰ C, 55 ± 5% R.H. inside the closed metal drums at varying exposure periods.....	125

7-Efficacy of CA of 99% N ₂ against the tested insect species at $18 \pm 2^{\circ}\text{C}$, $55 \pm 5\%$ R.H. inside the closed steel bins at varying exposure periods...	152
8-Efficacy of CA of 99% N ₂ against the tested insect species at $28 \pm 2^{\circ}\text{C}$, $55 \pm 5\%$ R.H. inside the closed steel bins at varying exposure periods...	155
9-Efficacy and combined action of 100 ppm of phosphine and $30 \pm 5\%$ carbon dioxide against the tested insect species at $18 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed steel bins at varying exposure periods.....	158
10-Efficacy and combined action of 100 ppm of phosphine and $30 \pm 5\%$ carbon dioxide against the tested insect species at $18 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed fiberglass bins at varying exposure periods.....	162
11-Efficacy and combined action of 100 ppm of phosphine and $30 \pm 5\%$ carbon dioxide against the tested insect species at $18 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed metal drums at varying exposure periods.....	162
12-Efficacy and combined action of 100 ppm of phosphine and $30 \pm 5\%$ carbon dioxide against the tested insect species at $28 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed steel bins at varying exposure periods.....	168
13-Efficacy and combined action of 100 ppm of phosphine and $30 \pm 5\%$ carbon dioxide against the tested insect species at $28 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed fiberglass bins at varying exposure periods.....	175

14-Efficacy and combined action of 100 ppm of phosphine and $30 \pm 5\%$ carbon dioxide against the tested insect species at $28 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed metal drums at varying exposure periods.....	179
15-Efficacy and combined action of 100 ppm of phosphine and $60 \pm 5\%$ carbon dioxide against the tested insect species at $18 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed steel bins at varying exposure periods.....	179
16-Efficacy and combined action of 100 ppm of phosphine and $60 \pm 5\%$ carbon dioxide against the tested insect species at $18 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed fiberglass bins at varying exposure periods.....	183
17-Efficacy and combined action of 100 ppm of phosphine and $60 \pm 5\%$ carbon dioxide against the tested insect species at $18 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed metal drums at varying exposure periods.....	189
18-Efficacy and combined action of 100 ppm of phosphine and $60 \pm 5\%$ carbon dioxide against the tested insect species at $28 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed steel bins at varying exposure periods.....	196
19-Efficacy and combined action of 100 ppm of phosphine and $60 \pm 5\%$ carbon dioxide against the tested insect species at $28 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed fiberglass bins at varying exposure periods.....	196
20-Efficacy and combined action of 100 ppm of phosphine and $60 \pm 5\%$ carbon dioxide against the tested insect species at $28 \pm 2^{\circ}\text{C}$ and $55 \pm 5\%$ R.H. inside the closed metal drums at varying exposure periods.....	200

VII

V-Conclusion.....	207
VI-Summary.....	211
VII-References.....	217
IX-Arabic Summary.	