



References

Reference

Abdel-Nasser A., Abdel-Azim., Taal E., El-Badawy S et al. (1998):

Depression and depressive symptoms in rheumatoid arthritis patients: An analysis of their occurrence and determinants. Br. J. of Rheuma 37: 391-397.

Anderson KO., Bradley LA., Young LD., et al. (1985) :

Rheumatoid arthritis. Review of psychological factors related to etiology, effects and treatment. Psychol Bull, 98:358-387.

Anderson., Ferrie., Baldwin. (2008) :

Evidence-based guidelines for treating depressive disorders with antidepressants: a revision of the 2000 British Association for Psychopharmacology guidelines. Journal of Psychopharmacology 22: 343-396.

Andrea W., Floris W., Rinie Geenen., Johannes W.(2003):

Pain coping and social support as predictors of long-term functional disability and pain in rheumatoid arthritis . Behaviour Research and Therapy, Volume 41, Issue 11, November 2003, Pages 1295-1310.

Andrew M H and Dominic YY. (2004):

Evaluation of a community rehabilitation service for people with rheumatoid arthritis. Patient Education and Counseling, Volume 55, Issue 1, October 2004, Pages 62-69.

Ang DC., Chori H., Kroenke K., Wolf F. (2005):

Comorbid depression is an independent risk factor for mortality in patients with rheumatoid arthritis. *J Rheumtol*; 32:1013-1019.

Angelopoulos N., Drosos AA., Kostouitsag E., Liskos A. (1988):

Personality and psychopathology in patients with primary Sogren's syndrome. *Terapevticheskii Arkhiv*, 60:49-52.

Anirban Ghosh., Biswadip Ghosh., Shantashil Pain. (2011):

Comparison between DAS28, CDAI and HAQ-DI as tools to monitor early rheumatoid arthritis patients in eastern India. *Indian Journal of Rheumatology* 2011 September. Volume 6, Number 3; pp. 116–122.

Anna Gramling and James R.(2012):

Initial Management of Rheumatoid Arthritis. Review Article. *Rheumatic Disease Clinics of North America*, Volume 38, Issue 2, May 2012, Pages 311-325.

Anne Mayoux.(2008).

Dynamic exercise therapy for rheumatoid arthritis . *Joint Bone Spine*, Volume 75, Issue 6, December 2008, Page 753.

Arnett FC., Edworthy SM., Bloch DA., McShane DJ., Fries JF., Cooper NS., et al.(1988):

The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis rheum*: Mar; 31(3):315-24.

Asaad T and Khla O.(2001):

Psychometric sleep assessments instruments: Arabic version for sleep evaluation. El Nahda. El Fagala. Egypt.

Asgary E., Veronica L., Catherine M. (2011):

Evidence-based program, Perspectives of service providers and administrators. *Professional Psychology: Research and Practice*, Vol 42(2), Apr 2011, 169-175.

Barlow JH., Wright CC., Turner AP., Bancroft GV.(2005):

A 12-month follow-up study of self-management training for people with chronic disease. *British Journal of Health Psychology* 10 (Pt 4):589-99.

Belzer K and Schneier FR. (2004):

Comorbidity of anxiety and depressive disorders: issue in conceptualization, assessment and treatment. *J Psychiatr Pract* 2004; 10:296-306.

Bernd T., Kivanc C., Gamze A., Ylker Y. (2008):

A comparative evaluation of health related quality of life and depression in patients with fibromyalgia syndrome and rheumatoid arthritis. *Rheumatology International* .28(9):859-865.

Berry H and Huskisson EC. (1972) :

Isotopic indices as a measure of inflammation in rheumatoid arthritis. *Ann. rheum. Dis.* (1972),523- 533.

Berry S and Romero R. (2006) :

Treatment of rheumatoid arthritis with a recombinant human tumor necrosis factor receptor. *Rheumatology* (January 2006) 45 (1): 1-5.

Bugajska J., Brzosko M., Jedryka-Góral A., Głuszko P. (2010):

Psychological stress in Rheumatoid arthritis patients: a comparative Polish–German study , Volume 9, Issue 4, February, Pages 211-215.

Carissa A.low ., Amy Lynn., Amy H.KAO. (2009):

Association between CRP and depressive symptoms in rheumatoid arthritis . *Biological psychology* volume 81, issue 2,page 131-134.

Carol Eustice. (2006):

Sleep Disturbance Evaluated in Rheumatoid Arthritis Patients. study published in the October 2006 issue of the Journal of Rheumatology.

Carol Eustice. (2011):

Depression and Rheumatoid Arthritis -Common comorbidities. Updated October 03, 2011. Health's Disease and Condition content is reviewed by the Medical Review Board.

Christin Miller . (2006):

Stress and Depression in Rheumatoid Arthritis. Health Central Sunday, February 26.

Chung-E Tseng., Jill P. Buyon., Mimi Kim.(2006):

The effect of moderate-dose corticosteroids in preventing severe flares in patients with serologically active, but clinically stable, systemic lupus erythematosus, *Arthritis & Rheumatism* ,Volume 54, Issue 11, pages 3623–3632, November 2006.

Costa., Antonio Filpi Coimbra da; Marco Antonio Alves ., Jose Angelo. (2008):

Depression, anxiety and disease activity in rheumatoid arthritis. *Rev. Revista Brasileira de Reumatologia*, vol.48, n.1, pp. 7-11.

Covic T., Adamson B., Spencer D., Howe G.(2005):

A biopsychological model of pain and depression in rheumatoid arthritis :a 12-month longitudinal study .*rheumatology* 2005;42:1287-94.

Creeds., Murphy F., Jayson M. (2005):

Psychiatric disorder and illness behaviour in rheumatoid arthritis. *Br J Rheumatol.* Oct;27(5):357-63.

Curtis R., Groarke A., Coughlan R., Gsel A. (2005):
Psychological stress as a predictor of psychological adjustment and health status in patients with rheumatoid arthritis. *Patient Educ Couns*, Vol.59, No.2 (November 2005), pp.192-198.

Daly D. (1942):

Central system in acute disseminated lupus erythematosus. *Journal of Nervous and Mental Disease*.

David B., Graham E., Greg C., Karen D. (2008):

Switching to Another SSRI or to Venlafaxine With or Without Cognitive Behavioral Therapy for Adolescents With SSRI-Resistant Depression, PubMed Central.

David Mandel. (2011):

The role of the rheumatologist in the management of rheumatic disease. www.dmandelmd.com/rheumatologist.htm.

Dicken C., McGown L., Clark-Carter D., Creed F. (2002):

Depression in rheumatoid arthritis :a systemic review of literature with meta-analysis. *Psychosom Med* 2002;64:52-60.

Fransen J (2005) :

The Disease Activity Score and the EULAR response criteria. *Clin Exp Rheumatol* 2005; 23 (Suppl. 39): S93-S99.

Goeb V., Jouen F., Gillber . (2009):

Diagnostic and prognostic uses of anti-CCP. *Turkish Journal of rheumatology Cilt 26, Sayı 3, Sayfa(lar) 193-198.*

Goldberg and Williams . (1988):

A user guide to the General Health Questionnaire Winder, Berkshire.

Helen Graves., David L Scott., Heidi Lempp., John Weinman.

(2009):

Illness beliefs predict disability in rheumatoid arthritis. *Journal of Psychosomatic Research*, Volume 67, Issue 5, November 2009, Pages 417-423.

Ja A and Bhadauria D. (2008):

Evaluation of efficacy of fluoxetine in the management of major depression and arthritis in patients suffering from rheumatoid arthritis. *WorldWideScience.org* .

Janca A., Ustrin B., Isaac M et al. (1994):

ICD-10 Symptom checklist for Mental Disorders Division of Mental Health World Health Organization. Geneva Version, 101.

Jeanette Raymond. (2010):

Disclosing Emotions Reduces Inflammation in Rheumatoid Arthritis, Ph.D., Body-Mind Psychotherapy *GoodTherapy.org* .

Jerant AF., Von Friederichs-Fitzwater MM., Moore M. (2005):

patients' perceived barriers to active self-managements of chronic conditions . *patient Educ conus* 2005;57:300-7

Katsumata Y., Harigai., M.Kawaguchi ., Hara M. (2007):

Diagnostic reliability of CSF test for acute confusional state in patient with SLE. *J Rheumatol.* 2007 Oct;34(10):2010-7. Epub 2007 Sep 15:

Katsumata Y., Harigai., M.Kawaguchi ., Hara M. (2010):

Diagnostic reliability of MRI for CNS syndrome in SLE. *PubMed Central.* Published online 2010 January 23.

Katz PP and Yelin EH .(2001):

Prevalence and correlate of depressive symptoms among persons with rheumatoid arthritis. *J Rheumatol*; 20:690-696.

Kimura A (2008) :

proteomic analysis of antibodies in SLE. *Nature Biotechnology* (26: 669–675).

Kirsch M., Hülseweh B., Nacke C., Rülker T., Schirrmann T.(2008):

Chronic stress and regulation of cellular markers of inflammation in rheumatoid arthritis: implications for fatigue. *Brain, Behavior, and Immunity* Volume 22, Issue 1, January, Pages 24-32.

Kotzin BL and Kozora E. (2001):

Anti-DNA meets NMDA in neuropsychiatric lupus. *Nature medicine* Vol.7, No.11, ISSN 1175-6.

Kurien B and Scofield R.(2006):

Autoantibody determination in the diagnosis of SLE. *Second J Immunol* 64,227-235.

Lars Klareskog., Anca Irinel C., Stephen A. (2009):

Rheumatoid arthritis management. *The Lancet*, Volume 373, Issue 9664, Pages 659 –665.

Lindberg NE., Lindberg E., Theorell T., Larsson G. (2002):

psychotherapy in rheumatoid arthritis--a parallel-process study of psychic state and course of rheumatic disease. *Z Rheumatol.*;55(1):28-39.

Lisa Emrich. (2009):

Rheumatoid Arthritis and Health Care Reform, health central sites .

Lisitsyna T A., Veltishchev D U., Seravina O F., Kovalevskaya O B. (2009):

Prevalence of Psychiatric Disorders in Rheumatoid Arthritis Patients .
Ann Rheum Dis 2009;68 (Suppl3):148.

Lok EYC., Scott Cunningham., Steve Smith et al. (2010):

Prevalence and Determinants of Psychiatric Disorders in Patients With
Rheumatoid Arthritis Psychosomatics, [http://www.mdlinx.com/
rheumatology/news](http://www.mdlinx.com/rheumatology/news).

*Loreto Carmona., Marita Cross., Ben Williams., Marissa
Lassere. (2010):*

Rheumatoid arthritis. Best Practice & Research Clinical
Rheumatology.24:733–745.

Lorig KR., Sobel DS., Ritter PL., Laurent D., Hobbs M. (2001):

Effect of a self-management program on patients with chronic disease.
Effective Clinical Practice; 4: 6, 256-262.

Low B., Willand L., Eick W et al. (2004):

Psychiatric comorbidity and work disability in patients with
inflammatory rheumatic disease . Psychosom Med 2004;66:58:55-62.

*Luyster FS., Chasens ER., Wasko MC., Dunbar-Jacob J.
(2011):*

Sleep quality and functional disability in patients with rheumatoid
arthritis. J Clin Sleep Med .2011;7:49–55.

Macrae R. (2004):

Radiographic evaluation of rheumatoid arthritis. In :Macrae's
examination .The clinical orthopedic examination (ed) 5th edition;5:16:74.

Majithia V., Stephen A., GeraKi A. (2007):

Rheumatoid Arthritis: Diagnosis and Management. The American
Journal of Medicine, Volume 120, Issue 11, November, Pages 936-939.

Malinow KL., Molina R., Gordon B., et al.(1985):
Neuropsychiatric dysfunction in primary Sjogren's syndrome. *Ann Intern Med*, 103(3):344-50.

Marcella Prete., Vito Racanelli., Liboria Digiglio., Angelo Vacca.(2011):
Extra-articular manifestations of rheumatoid arthritis: An update Review Article. *Autoimmunity Reviews*, Volume 11, Issue 2, December 2011, Pages 123-131.

Martin Winkler. (2008):
Rheumatoid Arthritis and Depression; Causes of Suicide. First version: 22 Jul 2008. Latest revision: 28 Aug.

Masayo Kojima., Toshishisa Kojima., Takeshi Oguchi. (2009):
Psychological factors , disease status and quality of life in patient with RA. *Journal of Psychosomatic Research*. Volume: 67, Issue: 5, Publisher: Elsevier Inc., Pages: 425-431.

Matthias Englbrecht., Laure Gossec., Anita DeLongis., Marieke Scholte., Tore K.(2012):

The Impact of Coping Strategies on Mental and Physical Well-Being in Patients with Rheumatoid Arthritis Original Research Article. *Seminars in Arthritis and Rheumatism*, Volume 41, Issue 4, February 2012, Pages 545-555.

Melissa M. Norberg, Gretchen J. Diefenbach, David F. Tolin (2008)

Quality of life and anxiety and depressive disorder comorbidity Original Research Article. *Journal of anxiety Disorders*, Volume 22, Issue 8, December 2008, Pages 1516-1522.

Melzack R and Wall PD. (1962):

On the nature of cutaneous sensory mechanisms. *Brain*, 85:331-356.

Michael R., Mary Davis ., Alex Zautra . (2008):

Behavioral co morbidities in RA *Psychiatric Times* . Volume .25 No,9 August 1.

Muellen AD and Lefkowitz AM.(1956):

Personality structure and dynamics of patients with rheumatoid arthritis . *J.Clin Psychol*;12:143 .

Nagyova I., Stewert RE ., Macejova Z ., Van Dijk JP ., Heuvel WJ. (2005):

The impact of pain on psychological well-being in Rheumatoid arthritis :the mediating effect of self-esteem and adjustment to disease. *Patient Educ Conus* 2005;58:55-62.

Newman and Mulligan . (2000):

The psychology of rheumatic disease . *Baillieres Best Pract Res Clin Rheumatol*;14:773-86.

Nicola J Gullick and David L Scott. (2011):

Co-morbidities in established rheumatoid arthritis *Best Practice & Research Clinical Rheumatology*. Volume 25, Issue 4 , Pages 469-483, August 2011.

Nierenberg AA (2001):

Current perspectives on the diagnosis and treatment of major depressive disorder. *Am J Manag Care* 2001;7(supp.11):S353-66.

Nishimura., Diane van der W., Tom Huizinga. (2007):

Diagnostic accuracy of anti-cyclic citrullinated peptide antibody and rheumatoid factor for rheumatoid arthritis .*Ann Intern Med*. Jun 5;146(11):797-808.

Okasha A. (1988):

Okasha's Clinical Psychiatry. PP: 1143. Anglo Egyptian Bookshop, Cairo.

Oslen and Stein. (2004):

Division of Rheumatology, Vanderbilt University School of Medicine, Nashville, USA. Comment in N Engl J Med. Aug 26;351(9):937-8.

Papakostas G., Shelton R., Smith J., Fava M.(2008):

Augmentation of antidepressant with antipsychotic medication for treatment resistant depressive disorder .A meta-analysis. Journal of clinical psychiatry, 68(6), 826-831.

Parker JC., Smarr KL., Slaughter JR et al. (2003):

Management of depression in rheumatoid arthritis: a combined pharmacologic and cognitive –behavioral approach. Arthritis Rheum 2003;49:766-77.

Perry M., Nicassio A ., Sarah R., Ormseth b., Morgan Kay C. (2011):

The contribution of pain and depression to self-reported sleep disturbance in patients with rheumatoid arthritis. by PM Nicassio – 2011 PAIN- 153 (2012) 107–112.

Plebani M., Pittani M., Celadin. (2009):

Recent advances in diagnostic technology in autoimmune disease. Autoimmunity Reviews 8 (2009) 238–243.

Prevo MLL., Vanthof MA., Kuper HH., et al. (1995):

Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. Arthritis Rheum. 1995;38:44–8.

Rahman A and Isenberg DA. (2008):

Management of systemic lupus erythomatosus. *N Engl J Med.* 2008; Feb 28;358(9):929-39.

Rindfleisch JA and Muller D. (2005):

Diagnosis and treatment of RA. *Rheumatology*; 2005;72 (6), 1037-1047.

Rogers MP and Helfgott. (1998) :

Psychiatric issue in rheumatology . In: , Meddison PJ, Isenberg DA, Woo P, Glass DN(eds); Oxford Textbook of rheumatology, Second edition, Section 1;chapter 3.12:pp 188-196.

Rush A., Trivedi H., Wisniewski S. (2006):

Bupropion-SR, Sertraline, or Venlafaxine-XR after failure of SSRIs for depression. *New England Journal of Medicine*, 354, 1231–1242.

Sam Norton., Amanda Sacke., Adam Young., John Done (2011):

Distinct psychological distress trajectories in rheumatoid arthritis: Findings from an inception cohort. 2011; Epub.

Scharloo M., Kaptein A., Weinman J., Hazes J., Willems L.(1998):

Illness perceptions, coping and functioning in patients with rheumatoid arthritis, chronic obstructive pulmonary disease and psoriasis . *Journal of Psychosomatic Research*, Volume 44, Issue 5, May 1998, Pages 573-585.

Shadick . (2010):

Changing the Physician-Patient Dynamic. Brigham and Women's hospital Biomedical research institute. *Rheumatology News* ,page 6.

Sharp L., Sensky T., Timberlake N., Ryan B. (2003):

Long-term efficacy of a cognitive behavioral treatment from randomized controlled trial for patients recently diagnosed rheumatoid arthritis , *Rheumatology* 2003;42:435-41.

Sharp L., Sensky T., Timberlake N., Ryan B., Brewin CR.(2001):

A blind, randomized, controlled trial of cognitive-behavioral intervention for patients with recent onset rheumatoid arthritis: preventing psychological and physical morbidity. *Pain* 2001;89:275-8.

Sheehy C., Murphy E., Barry M. (2006):

Depression in rheumatoid arthritis: underscoring the problem. *Rheumatology* 2006;45:1325-1327.

Sochet BR., Lisansky ET., Chubart AF et al. (1969):

A medical psychiatric study of patients with rheumatoid arthritis. *Psychosomatic*, 10:271.

Taylor S. (2006):

The future of selective serotonin reuptake inhibitors (SSRIs) in psychiatric treatment. *Med Hypotheses* ;66 (1):14-21.

Thea P M. (2003):

Rehabilitation of people with rheumatoid arthritis. *Best Practice & Research Clinical Rheumatology*. Vol. 17, No. 5, pp. 847-861, 2003.

Timonen M., Viilo K., Hakko H et al. (2003):

Suicides in persons suffering from rheumatoid arthritis. *Rheumatology* ;42:287-91.

Tomasevic-Todorovic S., Filipovic D., Boskovic K. (2010):

The effect of pain on cognitive abilities of patients with rheumatoid arthritis-*Clinical Neurophysiology*. Vol 121, Issue 4: page 18-20.

Treharne G., Lyons AC., Kitaz GD.(2008):

Suicidal ideation in patient with rheumatoid arthritis. Researches may help identify patients at high risk *Br Med J* 2008;321:1290.

Tseng CE., Buyon JP., Kim M., et al (2006):

The effect of moderate-dose corticosteroids in preventing severe flares in patients with serologically active, but clinically stable, systemic lupus erythematosus: *Arthritis Rheum* 2006; 54:3623–3632.

Tudor G., Weinberger M., de Vellis R., Beard A. (2008):

Communication about depression during rheumatoid arthritis patient visits. *Arthritis Care & Research*. Volume 59, Issue 2, pages 186–191, 15.

Turesson C., Matteson EL. (2008):

Extra-articular features of rheumatoid arthritis and systemic involvement. *Clin Rheumatol* 2008; 27:841-844

Turesson C., Mc Clelland RL., Christianson TJ., Matteson EL. (2007):

Sever extra-articular disease manifestations in patients with rheumatoid arthritis. *Ann Rheum Dis* 2007;66:70-5.

VanDyke MM., Parker JC., Smarr KL., Hewett JE., Johnson GE. (2006):

Anxiety in rheumatoid arthritis". *Arthritis and Rheumatism* (2006 Jun 15), pp. 408-412.

Venables PJW., MA., MB BChir. (2011):

Rheumatoid arthritis treatment, The content on the Up To Date website.

Vital E and Emery P. (2005):

Advances in the treatment of early rheumatoid arthritis." *Am Fam Physician* 72 Sep 15 2005; (6): 1002, 1004.

Vivian C., Leonard B., Robert J., Todd P. (2005):

Rehabilitation of orthopedic and rheumatologic disorders. *Archives of Physical Medicine and Rehabilitation*, Volume 86, Supplement 1, March 2005, Pages 61-68.

Walker D., Griffiths M., Dewer P et al. (1985):

Association of MHC antigens with susceptibility to and severity of rheumatoid arthritis in multicase families. *Ann. Rheuma. Dis.* 44: 519-529.

World health organization (1994):

International Classification of Diseases (ICD) ICD-10 was endorsed by the Forty-third World Health Assembly in May 1990 and came into use in WHO Member States as from 1994.

Y azici Y and Lockshin MD. (2000):

Patients with secondary Vasculitis in *Manual of Rheumatology and Orthopedics. Journal of Rheumatology* 2000;27:2817-2821.

Zautra AJ., Davis MC., Reich JW., Nicassario P. (2008):

Comparison of cognitive behavioral and mindfulness meditation interventions on adaptation to rheumatoid arthritis for patients with and without history of recurrent depression. *Clin Psychol.* 2008;76:408-21.
