

CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITTERATURE	3
 2.1 Relationship between agriculture practice and the infestation by seedling pests, <u>Aphids</u> and <u>Thrips</u>	 3
2.1.1 Density of plants and rate of infestation.	4
2.1.2 Other agriculture practice and rate of infestation	4
 2.2 Relationship between agriculture practice and rate of infestation by the cotton leafworm .	
2.2.1 Density of plants	6
2.2.2 Other agriculture practice and rate of infestation	6
 2.3 Relationship between agriculture practice and rate of infestation by bollworms	8
2.3.1 Density of plants and infestation .	8
2.3.2 Other agriculture practice and rate of infestation	11
 METHODS AND TECHNIQUE	15
3.1 Aphis and Thrips	16
3.2 Cotton leafworm, egg-masses <u>Spodoptera littoralis</u>	17
3.3 The cotton bollworms	17
3.4 Determination of yield loss	18
3.5 Statistical analysis	19

RESULTS and DISCUSSION	20
4.1 Effect of hill spacing on the rate of infestation of cotton seedlings with <u>Aphis gossypii</u> (Glov.)	20
4.1.1 Season 1981	20
4.1.2 Season 1982	24
4.2 Effect of row spacing on the rate of infestation of cotton seedlings with <u>Aphis gossypii</u> (Glov.)	26
4.2.1 Season 1981	26
4.2.2 Season 1982	30
4.3 Effect of planting on both sides of the row on the rate of infestation of cotton seedlings with <u>Aphis</u> <u>gossypii</u>	32
4.3.1 Season 1981	32
4.3.2 Season 1982	35
4.4 Effect of hill spacing on the rate of infestation of cotton seedlings with <u>Thrips tabaci</u> (lind.)	37
4.4.1 Season 1981	37
4.4.2 Season 1982	41
4.5 Effect of row spacing on rate of infestation of cotton seedlings with <u>Thrips tabaci</u> (lind.)	43
4.5.1 Season 1981	43
4.5.2 Season 1982	47

4.6	Effect of planting on both sides of the row on the rate of infestation of cotton seed- lings with <u>Thrips tabaci</u> (lind.).....	49
4.6.1	Season 1981	49
4.6.2	Season 1982	52
	DISCUSSION	54
4.7	Effect of density of the cotton plants on popu- lation density of the cotton leafworm, egg-masses <u>Spodoptera littoralis</u> (Boisd.)	55
4.7.1	Fluctuations of the egg-masses population in cotton fields	55
4.7.2	Effect of hill spacing on the abundance of the cotton leafworm, <u>Spodoptera litt-</u> <u>oralis</u> egg-masses	58
4.7.2.1	Season 1981	58
4.7.2.2	Season 1982	62
4.7.3	The effect of row spacing on rate of infes- tation of the cotton leafworm egg-masses	65
4.7.3.1	Season 1981	65
4.7.3.2	Season 1982	68
4.7.4	The effect of planting on the two sides of the ridge on the rate of infestation of the cotton leafworm egg-masses .	71
4.7.4.1	Season 1981	71
4.7.4.2	Season 1982	74

CONTENRS (cont.)	Page
DISCUSSION	76
4.8 The relation between Density of cotton plants and Rate of infestation by bollworms	79
4.8.1 Effect of hill spacing on the rate of infestation of the cotton bollworms .	79
4.8.1.1 Season 1981	79
4.8.1.2 Season 1982	85
4.8.2 Effect of row spacing on the rate of infestation of the cotton bollworm .	90
4.8.2.1 Season 1981	90
4.8.2.2 Season 1982	95
4.8.3 Evaluation of infestation by bollworm in relation to planting on both sides of the ridge	100
4.8.3.1 Season 1981	100
4.8.3.2 Season 1982	105
4.8.4, YIELD LOSS	110
DISCUSSION	113
5 SUMMARY	115
6 REFERENCES	118
7 ARABIC SUMMARY	124