

REFERENCES

Abdel-Salam, N.M. (1980).

Ecological studies on Genus Agrotis (Lepidoptera : Noctuidae).

M.Sc. Thesis, Ain Shams Univ.

Abul-Nasr, S., El-Sherif, S. and Naguib, M.A. (1972).
Oviposition behaviour of the cotton leafworm in clover fields.

Z. Angew. Ent., 19 (3) : 310-314.

(1973).

Seasonal fluctuations of moth of the cotton leafworm, S. littoralis (Boisd.) during the cotton season.

Bull. Soc. ent. Egypt, 57 : 413-418.

Abul-Nasr, S., Moussa, M.A. and Naguib, M.A. (1966-a).

Ecological study on the cotton leafworm, Prodenia litura (Fab.), Part I. The egg-masses in cotton fields.

Bull. Soc. ent. Egypt, 50 : 149-162.

(1966-b).

Ecological study of the cotton leafworm, Prodenia litura (Fab.), Part II. The larvae and pupae in cotton fields.

Bull. Soc. ent. Egypt, 50 : 173-180.

Bishara, I. (1928).

Preliminary experiments with dusting and spraying against insect pests of cotton.

Bull. Min. Agric. Egypt, No. 77, 11 pv.

Bishara, I. (1934).

The cotton leafworm, Prodenia litura (F.) in Egypt.
Bull. Soc. ent. Egypt, 18 : 223-404.

Bourne, A. (1936).

Value of electric traps against orchard insects.
Mass. Agric. Exp.Sta. Bull., 327 : 47-49.

Campion, D.G. (1976).

A comparison of the catches of moth of the cotton leafworm, Spodoptera littoralis in light and pheromone traps.

Miscellaneous Report No. 21, Center for Overseas Pest Research, College House, Wrights, Lane, London W 855 J, U.K.

El-Borolossy, F.M. (1959).

Studies on the effect of some major weather conditions on the activity of cotton leafworm and bollworms, and the estimation and forecasting of changes in their populations.

Ph.D. Thesis, Fac. Agric., Cairo Univ., Egypt.

El-Saadany, G. and Abd El-Fattah, M.I. (1975).

Über die nächtliche flugaktivität einiger an Baumwolle in Ägypten Schädlicher Lepidoptern.
Arten Anz. Schädlingskde, Pflanzenschutz, umveltschutz,
48 : 109-110.

Frost, S.A. (1952).

Light traps for insect collection survey and control.
Bull. 550, Agric. Expt. State College, Pennsylvania.

Hanna, H.M. (1972).

The flight activity of Agrotis ipsilon (Hufn.) at two levels in the vicinity of Assiut (Lepidoptera : Noctuidae).

Bull. Soc. ent. Egypt, 56 : 93-102.

Hanna, H.M. and Atries, I.E. (1968).

On the time of flight of certain nocturnal Lepidoptera
as measured by a light trap.

Bull. Soc. ent. Egypt, 52 : 535-545.

----- (1969-a).

The flight activity of certain nocturnal Lepidoptera
in relation to temperature and humidity.

Bull. Soc. ent. Egypt, 53 : 1-6.

----- (1969-b).

The effect of moonlight on certain nocturnal Lepidoptera.

Bull. Soc. ent. Egypt, 53 : 7-12.

Harvey, G.E.R. and Palm, C.E. (1935).

A preliminary report on the responses of the European
corn borer to light.

J. Econ. Ent., 28 : 670-675.

Hassan, A.S., Moussa, M.A. and Nasr El-Sayed A. (1960).

Factors affecting longevity and reproductive poten-
tials of moths of cotton leafworm, Prodenia litura(F.).

Bull. Soc. ent. Egypt, 44 : 383-386.

Hassanein, M.H. (1956).

Nocturnal activity of insects indicated by light traps.

Bull. Soc. ent. Egypt, 40 : 463-479.

-----, Khalil, F.M. and El-Naby, A.A. (1971).

Abundance and population density of three lepidopt-
erous insects in Upper Egypt.

Bull. Soc. ent. Egypt, 55 : 79-83.

Hosny, M.M. and Khattab, A.A.S. (1960).

Phenological studies on the abundance of seven species
of insect pests as indicated by catches in an Ultra-
violet trap in Giza Area.

Agric. Res. Rev., 78 (1) : 129.

Hosny, M.M., Nasr El-Sayed A. and El-Shafei, S.M. (1978).
Catches of S. littoralis (Boisd.) male moths in
light and pheromone traps in the Delta and in
Middle Egypt.

4th Conf. Pest Control, Cairo, 1-6.

Isshak, R.R. and Abdel-Megeed, M.I. (1975).

Horizontal distribution of egg-masses of cotton
leafworm, Spodoptera littoralis (Boisd.) in cotton
fields.

Z. ang. Ent., 78 : 65-69.

Jonescu, M., Romascu, E. and Coicev, A. (1963).

Investigations on the bionomics and control of
larvae of Agrotis segetum.

Anal. Sec. Prot. Pl. Inst. Cent. Cerc. Agric., 1 :
115-123.

Khattab, A.A. (1965).

Biological and ecological studies on Laphygma
exigua (H.b.) and its normal distribution at the
different regions during the various seasons.

M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt.

Larsen, E.B. (1943).

The importance of the master factors for the activity
of Noctuidae. Studies on the activity of insects.

I. Soertryk of Ent. Medd., 23.

Moussa, M.A., Zaher, M.A. and Kotby, F. (1960).

Abundance of the cotton leafworm, P. litura (F.),
in relation to host plants and their effect on the
biology.

Bull. Soc. ent. Egypt, 46 : 241-251.