

- (50) Lazarevskii, M.A. (1963): "How to study grape sorts" Ezotshene Sartov vinograda in Rus. Ezdat. Rostovskova uneversteta.
- (52) Levy, J.F. (1965): Identification et ctude par analyse foliare de quelques carences nlementaires de la vigne. Vignes et vins, No. 138:18-24.
- (53) Licui, R. (1969): Determination of bud fruitfulness in different vine varieties. Poljopr. Zananst. Smotra, 25(5): 1-24(Hort. Abst., 41:8690).
- (54) Marazova, J.S. and Negrul, A.M. (1972): practicals in viticulture Praktikum Povinogrodorstro, Ezd. Kolos. Moskow.
- (55) Melnik, C.A.; B.K. Kosareva (1965): Determination of nitrogen fertilizer required by using leaf analysis. Report, Lenin Akademia for Agriculture Science No.6 (1965).
- (56) Minesey, F.A., M.W. Taha and M.G. Maughice (1965): Seasonal changes in nitrogen in some grape varieties grow under different nitrogen levels in relation to yield and fruit quality. Alex. Jour. Agri. Res. 12(2): 1-24.
- (57) Minessy, F.A., M.O.El- Karouri, M.T. Biely and A. El-Fahl. (1971): Studies on the nutritional status of fruit trees in the Northern Prövine, Sudan. Sudan. Agric. Jour, Vol. 6, 1971.
- (58) Michel Dubois , K. A. Gilles , J. K. Hamilton , P. A. Rebers , and Smith , 1956 : Colorimetric Method for Detrmination of sugars and Related substance . Anal. Chem. Vol. 28, No. 3:350 -354
- (59) Mume, L. and J. Dulac. (1945): Carence potassique chez la vigne decelee (diagnosed) pha la controle chimique de la feuille, avant l'apparition de la brunissage. Comptes Rendus. Aad. of Sciences, paris Vol. 221:116-117.

- (60) Mume, L. and J. Dulac. (1948): Nouvelle observations sur la nutrition de la vi, ne controlee par l'analyse chimique de la feuille. Compt. Rend. Acad. Agf. 34:861-864.
- (61) Muthukrishnan, C.R. and Srinivesan, C. (1974): Correlation between yield, quality and petiole Nutrients in grape. Vitis. 12(4):227-285.
- (62) Nastev, D. (1964): (Results obtained hitherto on the fertility of the vine variety. Variety Sultanina.)
- (63) Olsen, S.R, Cole, C.V., Watanable, F.S. and Dean, L.A. (1954): Estimation of available phosphorus in soils by extraction with sodium bicarbonate. U.S. Dept. Agric. Cir- 939.
- (64) Papov, T. (1972): The effects of length of the fruiting shoot on the biological uptake of nitrogen, phosphorus and potassium by certain vine varieties. Hort. Abst. 42(3)- 5720.
- (65) Partridge, N.L. (1925): In "Small fruit Culture" shoemaker, T.S., 3rd ed., 1954: Mc Grow Hill Book Company, INC., New York, Toronto, London, p.84.
- (66) Piper, C.S. (1950): Soil and Plant Analysis. Inter Science Publications, Inc. New York.
- (67) Polyak, D.; Kopcsal, P.M.; Angyan, F. (1974): Leaf analysis of grape vines of the Tokay region. Tokajhegyalia szoloinek levelanalizise. Hort. Abs. No. 3075.
- (68) Pregl, E. 1945 Quantative Organic Micro, Analysis 4 th. Ed.j. Chndril, London
- (69) Ravikovitch, S., and N. Bidner. (1937): The deterioration of grape vines in salin soils- Empire. Jour. Exp. Agric. 5:197-203.

- (70) Romano, Gallo, J. and Wilson R. Correa. (1962): Foliar mineral composition of different scion rootstock combinations for ten grape varieties. *Bragantia* 21:pt. 1a, 397-410. Abstract only seen. *Chem. Abst.* 60:85 I.
- (71) Saini, S.S; R.V. Singh. (1975): Effect of nitrogen, Phosphorus and potassium on the nutritional status of young vines. *Indian Journal of Agricultural Sciences* 4,(V2) 1-10 (Rn, 18, ref.) Indian Agricultural Research institute, New Delhi, India, *Hort. Abst.* 1979 Vol. (43) 3.
- (72) Sharaf, M.M. (1979): Effect of nitrogen and other mineral nutrients content in the different organs of vines as influenced by soil type and age of grape plant. Ph.D. Thesis, Bulgaria.
- (73) Shaulis, Nelson. (1954): Potash deficiency in the vineyard and its cure. N.Y. Agr. Exp. Sta- Farm Research. 20(2):4.
- (74) Shaulis and Kimball (1956): The association of the nutrient composition of Concord grape petiole with deficiency symptom, growth and yield. *Proc. Amer. Soc. Hort. Sci.* Vol. 68:141-156.
- (75) Smith, C.B., Harold K. Fleming and Harry J. Pearbaugh, (1957): The Nutritional Status of Concord Grape Vines in Erie County, Pennsylvania as Indicated by Petiole and Soil Analysis. *Proc. Amer. Soc. Hort. Sci.* 70:189-196.
- (76) Snedecor, G.W. and W.G. Cochran. (1972): Statistical methods 6th. Ed. The Iowa state Univ. Press, Ames, Iowa, U.S.A. pp. 593.
- (77) Stanimirovic, P. (1970): The use of foliar diagnosis for determining the nutrient requirements of vines. *Hort. Abst.* 40:5915.

- (78) Stoev . K, D(1952) Biological Characteristics of lateral shoots of grape vines . Agrobiological N. 6
- (79)Taha, M.W. (1958): Effect of severity of Pruning on the Nutrient Element Content of Concord Grape leaf Petioles, Vine Vigor, Yield, and Friuit Quality.
- (80)Taha, N.W. (1965): Effect of nitrogen fertilization from different sourses on nitrogen content of leaf Petioles, yield and fruit quality of some grape varieties - Alex- Jour. Agric. Res. 12(2): 129-150.
- (81)Ulrich, A. (1944): Potassium content of grape leaf petioles and blades contrasted with soil as an indicator of the potassium status of the plant. Proc. Amer. Soc. Mort. Sci. 41:204-212.
- (82)Wiliams, W.O. (1946): California vineyard fertilizer expermentation. Amer. Soc. Hort. Sci. Proc. 48:269-278.
- (83)Williams, L.E. (1987): Growth of Thompson seedless' grape vines II. Ntrogen distribution. Joournal of the American Society for Horticultural Science 112(2) 330-333 {En, 23 ref.} California University, Davis, CA 95616, USA.
- (84)Winkler, A.J. and E.M. Shamsettin (1937): Fruit bud and flower formation in the Sultanina grape. Hilgardia Vol. 10, No. 15, May 1937.
- (85)Winkler, A.J., Cook, J.A., Kliewer, W.M., and Līder, I.A., (1974): General Viticulture. Unv. of California Pres. U.S.A.