

VI - REFERENCES CITED

- Abbassy, M.A., El. Shazli, A. and El-Gayar, F. (1977).
A new antifeedant to Spodoptera littoralis
Boisd. (Lepid, Nectuidae) from Acokanthera
spectabilis Hook. (Apocynaceae). Zeit. Ang.
Ento. 83 (3) 317-322.
- Ahmed, S.M.; Shalaby, F.F. and El-Khishin, M.K. (1978)
Antifeedant properties of some plant extracts
for the larvae of Spodoptera littoralis
(Boisd) 4 th conf. pest control NRC, Cairo.
- All, J.N.; Benjamin, D.M. and Matsumura, F. (1975). Influence
of Semipurified constituents of juvenile
jack pine, Pinus banksiana, foliage and other
pine-derived chemicals on feeding of Neodiprion
Swainei and N. rugifrons larvae. Annals of the
Entomological Society of America 68 (6) 1095-1101.
- All, J.N. and Benjamin, D.M. (1975). Olfactory response of
Neodiprion rugifrons larvae to inhibitory
extracts of juvenile jack pine, pinus bank-
siana. foliage. Journal of the Georgia
Entomological Society 10 (4) 300-305.
- Attri, B.S. (1977). Utility of neem oil extractive as feed-
ing deterrent to locust. Indian Journal of
Entomology. 37 (4) 417-418.
- Bailey, L.H. (1935). The standard cydopedia of horticul-
ture the Macmillan company, New York, Being-
Vols. I and II of original Edition, pp 3639.
- Bailey, L.H. and Bailey, E.Z. (1949). Hortus second.
A. Concise Dictionary of Gardening, General
Horticulture and cultivated plants in North
America. The Macmillan company, New York,
PP 778.
- Balbaa, S. I. Chemistry of crude drugs, Laboratory manual.
El Shaab Printing House, Cairo 1976 .

- Barton, D.H.R.; Cheung, H.T.; Cross, A.D.; Jackman, L.M. and Martin-Smith, M. (1961a). The constitution of clerodin. *Proc. Chem.Soc.* , 76-77.
- Barton, D.H.R.; Cheung, H.T.; Cross, A.D.; Jackman, L.M. and Martin-Smith, M. (1961b). Diterpenoid bitter principles. Part III. The constitution of clerodin. *J. Chem. Soc.*, 5 obl.5073.
- Bernays, E.A. and Chapman, R.F. (1975).. The importance of chemical inhibition of feeding in host plant selection by *parallelus* (Zetterstedt) . *Acarida* 4 (2) 83-93.
- Bernays, E.A. and Chapman, R.F. (1976). Antifeedant properties of seedling grasses. *Symp. Biol. hung.* 16:41-46.
- Bernays, E.A. and Chapman, R.F. (1977). Deterrent chemicals as a basis of oligophagy in Locust migratoria (L.) *Ecol. Ento.* 2:1-18.
- Butterworth, J.H. and Morgan, E.D. (1968) Isolation of a substance that suppresses feeding in Locusts *chem. comm.* 23-24.
- Butterworth, J.H. and Morgan, E.D. 1971. Investigation of the locust feeding inhibition of the seeds of the neem tree Azadirachta indica. *J. Insect physiol.* 17,969-977.
- Capinera, J.L. and Stermitz, F.R. (1979). Laboratory evaluation of Zanthophylline as a feeding deterrent to range Caterpillar, migratory grasshopper, alfalf weevil, and greenbug. *Journal of chemical Ecology*, 5(5) 767-771.
- Chauvin, R. (1946) Sur la substance qui, dans les feuilles de Melia azederach, repousse Les criquets. *C.r. Acad. Sci. Paris*, 222, 412-414.
- Gill, J.S. and LEWIS, C.T. (1971) Systemic action of an insect feeding deterrent. *Nature*, 232,402-403.

- Gombos, M.A. and Gasko, K. (1977) Extraction of natural antifeedants from the fruits of *Amorpha fruticosa* L. *Acta Phytopathologica Academiae Scientiarum Hungaricae*. 12 (314) 349-357.
- Gour, A.C. and Prasad, S.K. 1970-Effect of organic matter and inorganic fertilizer on soil and plant nematodes. *Indian J.Ent.*, 32 (2): 186-88.
- Granich, M.S.; Halpern, B.P. and Eisner, T. (1974) Gymnemic acids: Secondary plant substance of dual defensive action? *Journal of Insect physiology* 20 (2) 435-439.
- Heal, R.E.; Rogers, E.F., Wallace, R.T. and Starnes, O. 1950. A survey of plants for insecticidal activity. *Lloydia*, 13:89-162. (c.f.)
C. Insecticides from plants, a review of Literature 1941-1953. by M. Jacobson *Agriculture hand book no. 154* (1958) USDA.
- Higgins, R.A. and Pedigo, L.P. (1979). A laboratory antifeedant simulation bioassay for phytophagous insects. *Journal of Economic Entomology* 72 (2) 238-244.
- Hosozawa, S.; Kato, N. and Munakata, K. (1973). Diterpenoids from *Caryopteris divaricata*. *Phytochemistry* 12, 1833-1834.
- Hosozawa, S.; Kato, N. and Munakata, K. (1974a) Diterpenes from *caryopteris divaricata*. *Phytochemistry*, 13, 1019-1020.
- Hosozawa, S.; Kato, N.; Munakata, K. and Chen, Y.L., 1974. Antifeeding active substances for insects in plants. *Agr. Biol. Chem.* 38, 1045-1048.
- Hosozawa, S., Kato, N. and Munakata, K., 1974 b. Diterpenoids from *Clerodendron calamitosum*. *phytochemistry*, 308-309.

- Hosozawa, S.; Kato, N. and Munakata, K. (1974 c). Absolute configuration of caryoptin and 3-epicaryoptin. An exception in the exciton chirality method. *Tetrahedron Letters*, 3753-3756.
- Hosozawa, S.; Kato, N.; Munakata, K. and Chen, Y.L. (1974b) Antifeeding active substances for insect in plant-Agr. Biol. Chem., 38 (5), 1045-1048.
- Hunter-Jones, P. 1961. Rearing and breeding locusts in the laboratory .
Anti-locust Research Center, London.
- Ikeda T.; Matsumura, F. and Benjamin, D.M. (1977). Chemical basis for feeding adaptation of pin sawflies Neodiprion rugiformis and Neodiprion Swainel *Science* 2. USA 197 (4302)497-499 .
- Jones, C.G. (1979). An automatic feeding detector (AFD) for use in insect behaviour studies. *Entomologia Experimentalis et Applicata*, 25 (L) 112-115.
- Ju-Lurie (1975). Handbook of analytical chemistry (English translation)Mir publishers, Moscow 488pp, p:332-337.
- Kato, N.; Shibayama, S.; Munakata, K. and Katayama, C. (1971) Structure of the diterpene clerodendrin A. *Chem. Comm*, 1632-1633.
- Kato, N., Takahashi, M. Shibayama, M. and Numakata, K. (1972). Antifeeding active substances for insects in Clerodendron tricotomum Thunb. *Agr. Biol. Chem.*, 36,2579-2582.
- Khan, A.M. ; Adhami; Siddiqui, Z.A. and Saxena, S.K. (1967). Effect of different oil-cakes on hatching of larvae and on the development root-knot caused by Meloidogyne incognite. *International symp. plant pathology*, New Delhi, India , 65 pp.
- Kim, M., Koh, H.S.; Ichikawa, T., Fukami, H. and Ishii, S. (1975). Antifeedant of barnyard grass against the brown planthopper, Nilaparvata lugens (Stål) (Homoptera: Delphacidae). *Applied Entomology and Zoology*, 10 (2) 116-122.

- Kim, M.; Koh, H.S.; Obata, T.; Fukami, H. and Ishii, S. (1976). Isolation and identification of trans-aconitic acid as the antifeedant in barnyard grass against the brown planthopper, Nilaparvata lygens (Stål) (Homoptera: Delphacidae). Applied Entomology and Zoology 11 (1) 53-57.
- Kumar, A.; Tewari, G.D. and Pandey, N.D. (1979). Anti-feeding and insecticidal properties of bitter gourd, Momordica charantia Linn. against Athalia proxima klug-Indian Journal of Entomology, 41 (2) 103-106.
- Lavie, D., Jain, M.K. and Shpan-Gabrie Lith, S.R. 1967. Alocust phagorepellent from two Melia species chem. comm. 910-911.
- Ladd, T.L., Jr., Jacobson, M. and Buriff, C.R. (1978). Japanese beetles: extracts from neem tree seeds as feeding deterrents. Journal of Economic Entomology, 71 (5) 810-813.
- Leheta, M.F. (1979). Classification of colour patterns of hoppers of the desert locust, (Schistocerca gregaria Forsk.), with reference to colour variations in transiens forms. FAO, plant production and protection Division Desert Locust project, Progress Report, No.AGP/DL/TS/20. pp.10-39.
- Mathur, A.C. and Sharma, A.K. (1978) Resistance of Aristolochia sp. to insect attacks: a chemical basis. Indian Journal of Horticulture, 35 (4) 406-408.
- Matsuda, K. (1978). Feeding stimulation of flavonoids for various leaf beetles (coleoptera:chrysomelidae). Applied Entomology and Zoology, 13 (3) 228-230.
- Meisner, J.; Kehat, M.; Zur, M. and Elzick, C. (1978) Response of Earias insulana Boisd. larvae to neem (Azadirachta indica A.Juss) Kernel extract. phytoparasitica, 6 (2)-38.

- Meisner, J. and Skatulla, U. (1975). phytostimulation and phagodeterreny in the larvae of the gypsy moth, porthetria dispar L. phytoparasitica, 3 (1) 19-26.
- Mishra, S.D. and Prasad, S.K. 1974. Effect of soil amendments on nematodes and crop yield. I. oil seed cakes. organic matter, inorganic fertilizers and growth regulators on nematodes associated with wheat and their residual effect on mung. Indian J. Nematol, 4 (1): 1-9.
- Mishra, S.D. and Prasad, S.K. 1977. Effect of neem cake water extract on Root nematode (Meloidogyne incognita) infesting tomato. Indian J. Ent. 39 (3) 228-231.
- Mong Ting Tan and Sudderuddin, K.I. (1978). Effects of a neem tree (Azadirachta indica) extracts on diamond-back moth (plutella xylostella L.) Malaysian Applied Biology, 7 (1) 1-9.
- Morgan, E.D. and Thornton, M.D. 1973. Azadirachtin in the fruit of Melia azedarach. Phytochemistry, 12, 391-392.
- Munger, R. and Siegler, E.H. (1937) Stomach poison method for cockroaches. Soap and Sanit. Chem. 13 (16) 94.
- Narayanan, C.R.; Singh, R.P. and Sawaikar, D.D. (1978 reed. 1980). Phagodeterreny of various fractions of neem oil against Schistocerca gregaria F.: Pesticides, 12 (11) 31-32.
- Nielsen, J.K. (1978). Host plant discrimination within cruciferae: feeding responses of four leaf beetles (Coleoptera: Chrysomelidae) to glucosinolates, cucurbitacins and cardenolides. Entomologia Experimentalis et Applicata, 24 (1) 41-54.

- Pandey, N.D.; Mahendra Singh and Tewari, G.C. (1977). Antifeeding, repellent and insecticidal properties of some indigenous plant materials against mustard sawfly, Athalia proxima klug. Indian Journal of Entomology, 39 (1) 60-64.
- Patel, H.K.; Patel, V.C.; Chari, M.S.; Patel, J.C. and Patel, J.R. (1968). Neem seed paste suspension-A sure deterrent to Hairy Caterpillar (Am-sacta moorei But). The Madras Agricultural Journal, 55 (11) 509-510.
- Paul, I.C.; Sim, G.A., Hamor, T.A. and Robertson, J.M. (1962) The structure of clerodin: x-ray analysis of clerodinbromo-lactone. J. Chem. Soc. 4133-4145.
- Paul, C.F.; Dixit, R.S. and Agarwal, P.N. (1963). Evaluation of the insecticidal properties of the seed oil and leaf extract of the common Indian Neem, Azadirachta indica Linn. Science and Culture, 29 (8) 412-413.
- Pendse, G.S.; Ghokale, V.G., Phalnikar, N.L. and Bhide, B.V. 1946. Investigation of new plant larvicides with special reference to Spilanthes aermella. Jour. Univ. Bombay, Sect, A. Sci 15:26-30. (c.f.)
C.F. Insecticides from plants by M. Jacobson (1958). A review of literature from 1941-1953. Agric. Hand book No. 154 USDA.
- Ranganathan, S.K.; Koshi, T. and Sitaraman, N.L. 1949. Methyl anthraquinone (tectoquinone)-a synergist for 2,2-bis - cp. chlorophenyl 1,1,1-trichloroethane (DDT). Nature (London) 164 =1095.
- Rao, P.J. and Mehrotra, K.N. (1977). phagostimulants & Antifeedants from Calotropis gigantea for Schistocerca gregaria F. Indian Journal of Experimental Biology Vol. 15, NO.2, pp. 148 150.

- Rao, P.J. and Mehrotra, K.N. (1978) Feeding deterrent from *calotropis gigantea* for locusts. Indian Journal of Entomology (1978) 40 (1).
- Russel, G.B.; Sutherland, O.R.W.; Hutchins, R.F.N. and Christmas, P.E. (1978). Vestitol: a phytoalexin with insect feeding-deterrent activity. Journal of Chemical Ecology. 4(5) 571-579.
- SIM, G.A.; Hamor, T.A.; Paul., I.C. and Robertson, J. M. (1961). The Structure of Clerodin. Proc. Chem. Soc., 75-76.
- Singh, R.P. (1974). *Crinium bulbispermum*-an antifeedant to *Schistocerca gregaria* Forskal. Entomologist's Newsl., New Delhi, 4:34.
- Singh, R.P. and Pant, N.C. (1978). Antifeedants property of Sub-family Amaryllidoideae. Entomologists, Newsletter, 8 (3)8 .
- Singh, R.S. and Sitaramaiah, K. (1966). Incidence of root-knot of okra and tomatoes in oil cake amended soil. pl. Dis. Repr., 50:668-72.
- Steets, R. (1975). The effect of crude extracts of the meliaceous plant *Azadirachta indica* and *Melia Azederach* on various insect species. Die Wirkung von Rohextrakten aus den Meliaceen *Azadirachta indica* und *Melia azederach* auf Verschiedene Insektenarten. Zeit. Ang. Ento. 77 (3) 306-312 .
- Sudhakar, T.R.; Pandey, N.D. and Tewari, G.C. (1978). Antifeeding property of some indigenous plants against mustard sawfly, *Athalia proxima* klug. (Hymenoptera: Tenthredinidae,) Indian Journal of Agricultural Sciences, 48 (1) 16-18.
- Sutherland, O.R.W.; Mann, J. and Hillier, J.R. (1975). feeding deterrents for the grass grub *Costelytra zealandica* (Coleoptera: Scarabaeidae) in the root of a resistant pasture plant, *lotus pedunculatus*. New Zealand Journal of Zoology, 2 (4) 509-512.

- Tackholm, V. (1974). Students' flora of Egypt. Cairo University, printed by Cooperative Printing Company, Beirut, pp 888.
- Trial, H.; JR. and Dimond, J.B. (1979). Emodin in buck thorn: a feeding deterrent to phytophagous insects. Canadian Ethomologist, 111 (2) 207-212.
- Vignerón, J.P. (1978) Antifeeding substances of natural origin. Substances antiappetantes d'origine naturelle. Ann-Zool. Ecol. Anim., 10 (4) 663-694.
- Volkonsky, M. (1937) Insect-repellant action of extracts of the leaves of Melia azederach. Archs inst. pasteur, Alger. 15, 427-418.
- Wagner, M.R.; Ikeda, T.; Benjamin, D.M. and Matsumura, F. (1979) Host derived chemicals: the basis for preferential feeding behaviour of the larch sawfly, pristiphora erichsonii (Hymenoptera: Tenthredinidae), on tamarack, Larix laricina. Canadian Entomologist, 111 (2) 165-169.
- Warthen, J.D. and JR. (1979) Azadirachta indica: a source of insect feeding inhibitors and growth regulators. Agricultural Reviews and Manuals, Science and Education Administration, United states Department of Agriculture, ARM-NE-4, 11+21 pp.
- Whittaker, R.H. 1970. The biochemical ecology of higher plants in chemical Ecology, E. Sondheimer and J.B. Simeone Eds. Academic Press, N.Y. pp.43-70
- Whittaker, R.H. and P.P. Fecny. 1971. Allelochemicals: Chemical interaction between Species. Science 171:757-770.
- Wolcott, G.N. 1946. Factors in the natural resistance of Woods to termite attack. Caribbean Forster 7:121-134.

- Wolcott, G.N. 1946 . What to do about polilla. puerto Rico Univ. Agr. Expt. Sta. Bull 68:29 pp.
- Woodhead, S. and Bernays, E. (1977) changes in release rates of cyanide in relation to palatability of Sorghum to insects. Nature, UK, 270 (5634) 235-236.
- Woodhead, S. and Bernay , E.A. (1978). The chemical basis of resistance of Sorghum bicolor to attack by locusta migratoria. Entomologia Experimentalis et Applicata, 24 (2) 123-144.
- Yoshihara, I. Sogawa, K.; Pathak, M.D . , Julianio, B.O. and Sakamura, S. (1979). Soluble silicic acid as a sucking inhibitory substance in rice against the brown plant hopper (Delphacidae, Homoptera). Entomologia Experimentalis et Applicata, 26 (3) 314-322.