




# REFERENCES

## 7. REFERENCES

- A.O.A.C. (1975):** Official methods of analysis of Association of Official Analytic Chemists. Pub. A.O.A.C. INC. Suite 400, 22201 USA fifteenth Ed. 1990; PP: 62-63, 236 and 877-878.
- Abbas, M. S. T. and Y. A. A. El-Deeb (1993):** On the natural enemies of the major pests infesting cotton in Egypt. Egyptian Journal of Agricultural Research, 71(I): 131-138.
- Abbott, H. A. (1986):** The influence of molasses on the physical properties and spray ability of two pesticide formulations. pesticide-Science, 17(5): 526-534.
- Abd- El-Khaliq and M. K. Abbasi (2006):** Effects of integrated use of organic and inorganic nutrient sources with effective microorganisms (EM) on seed cotton yield in Pakistan. Bioresource- Technology, 97 (8): 967-972.
- Abd- El-Naser, S.; E. D. Ammar and S.M. Farrag (1979):** Rates of infestation by *Pectinophora gossypiella* and *Earias insulana* on flowering sites of the cotton plant. Deutsch Entomologische zeitschrift, 26 (113): 165-172.
- Abo- Elhagag, G. H. (2) (1998):** Effect of spraying cotton plants during the early season against cotton aphid on cotton; pests, natural enemies and some crop characters in southern Egypt. Assiut Journal of Agricultural Sciences, 29 (4): 91-100.

- Agnihotri, N. P.; K. P. Srivastava; V. T. Gajbhiye and H. K. Jain (1986):** Relative efficacy of some synthetic pyrethroids and other commonly used insecticides against bollworms and their residues in cotton. *International Journal of Tropical Agriculture*; 4(2): 168-174.
- Ali, K. A. H. (1998):** Studies on some arthropodous natural enemies on the pink bollworm *Pectinophora gossypiella* (Saunders). M. Sc. Thesis, Faculty Agriculture, Moshtohor, Benha Branch, Zagazig University.
- Ali, K. A. H. (2003):** Trials for mass-rearing and release of some predaceous insects for controlling the pink and spiny bollworm. Ph.D. Thesis, Faculty Agriculture, Moshtohor, Benha Branch, Zagazig University.
- Bishara, I. (1968):** Bollworm attack in relation to manuring of cotton. Ministry of Agriculture, Egypt, Tech. Bull. 1: 35pp.
- Brown, J. D. and O. Lillard (1946):** Rapid determination of potassium and sodium in plant material and soil extract by flame photometry. *Proc. Amr. Soc., Hort., Sci.*, 73, 813.
- Butter, N. S. and H. S. Sukhija (1987):** Efficacy of flucythrinate (pay off 10 EC) against bollworms (*Pectinophora gossypiella*, *Earias* spp. and *H. armigera*) infesting cotton. *Journal of Research, Punjab Agricultural University*, 24(4): 615-622.

- Castella, J. C.; Dollon and S. Savary (2005):** Path coefficient analysis to assess yield losses due to a multiple pest complex in cotton in Thailand. *International Journal of Tropical Insect Science*, 25 (1): 39-49.
- Damo, M. C. and Solsoloy (1997):** Efficacy of *Bacillus thuringiensis* chemical insecticide mixtures for cotton bollworm control. *Philippine Entomologist*, 11(1): 57-64.
- Deshpande, R. R.; S. S. Shaw; R. N. Ganguli and K. C. Mandloi (1988):** Comparative toxicity and economics of some pyrethroid and non pyrethroid insecticides against cotton bollworms. *Agricultural Sciences-Digest- Karnal*, 8 (3): 131-134.
- Dhawan, A. K.; G. S. Simwat and A. S. Sidhu (1990):** Management of bollworms on cotton with synthetic pyrethroids. *Journal of Insect Science*, 3 (2): 158-161.
- Dunn, D.; S. G. Stevens and B. Phipps (1999):** The use of concentrated molasses soluble as a nitrogen and sulfur fertilizer in cotton production. *Proceedings- Beltwide-Cotton- Conference, Orlando, Florida, USA 3-7 January*, (2): 1275-1276.
- El-Gemeiy, M. M. and A. I. Afify (1992):** The efficacy of *Bacillus thuringiensis Berliner* against the spiny bollworm *Earias insulna* by adding a feeding stimulant. *Bull Soc. Ent. Egypte*, 63: 37-41.

- El-Gemeiy, M. M. and S. M. Naguib; M.A. Shalaby and M.E. Wahba (1999):** Efficacy evaluation of biocide Ecotech PRO and the organophosphorus insecticide isometaphos for controlling the bollworms, *Pectinophora gossypiella* and *Earias insulna*. Annals of Agriculture Science, Moshtohor. 37 (1): 673-684.
- El-Hammawi, A. E. and M. A. H. El-Ghandour (1974):** Preliminary studies on effect of foliar feeding during the flowering period on growth and yield of cotton plants. Egypt. J. physiol. Sci., 1: 71-77.
- El-Hattab, R. K. and A. A. Abdel-Raheem (1965):** The effect of sowing methods, nitrogenous fertilizers and number of plants per hill on some agronomic characters of cotton. Bull. Fac. Agric. Cairo Univ., 162 (1): 21-50.
- El-Heneidy, A. H.; M.S.T. Abbas and A.A. Khidr (1987):** Comparative population densities of certain predators in cotton fields treated with sex pheromones and insecticides in Menofia Governortae, Egypt. Bull. Ent. Soc. Egypt, Economic Series, 16: 181-190.
- El-Kady, W.A.; I. S. El-Hawary and S. E. D.A. Faizy (1988):** Effect of nitrogen and potassium fertilizers on seed cotton yield and insect pests infestation. J. Agric. Res., Tanta Univ., 14(2): 1426-1437.
- El-Nawawy, A. S.; M.A. Ashry; A.A. Zein; A. H. Masoudi, A. G. Abd-El-Hafez and R. El-Safty (1983):** Foliar fertilizers, insecticides and their mixtures as protectants of cotton bolls against pink bollworm. Mededlingenvan de faculteit land bouwwetenschappen Rijksuniversiteit-Gent., 48(2): 349-357.

- El-Sayed, A.A.A. (2001):** Studies on pink bollworm *Pectinophora gossypiella* and *Earias insulana* in cotton fields, Sharqia Governorate. M. Sc. Thesis, Fac. Agric. El-Zagazig Univ., Egypt. PP.132.
- Fadare, T.A. and N.A. Amusa (2003):** Comparative efficacy of microbial and chemical insecticides on four major lepidopterous pests of cotton and their natural enemies. African Journal of Biotechnology, 2 (11): 425-428.
- Gardner, B.R. and T.C. Tucker (1967):** Nitrogen effect on cotton 1-vegetative and fruiting characteristics. 2, Soil and petiol analysis. Proc. Amer. Soil, 31:780-791.
- Gomaa, M. (1999):** Biological and ecological studies on pink and spiny bollworms in Egypt. Ph.D. Thesis, fac. Agric., El-Azhar, Univ., pp. 188.
- Gormus,-O (2004):** Yield response of cotton to foliar fertilization of potassium in Cukurova Region. Department of Field Crops, Faculty of Agriculture, Cukurova University, Adana, Turkey, Turkish-Journal-of-Field-Crops. 2004; 9 (1): 35-42.
- Gupta, G. P. and K. N. Katiyar (1987):** Evaluation of new synthetic pyrethroids and formulations against bollworm complex in cotton. Pesticides, 21(4): 20-22.
- Gupta, G. P.; N. P. Agnihotri; K. N. Katiyar and H. K. Jain (1996):** Bioefficacy and persistence of synthetic pyrethroids in cotton. Pesticide Research Journal, 2 (2): 115-122.

- Habib, A.; A.M. Afify and D.S. Daoud (1976):** Seasonal abundance of some predators in cotton fields and its relation to certain pests. Bull. Soc. Ent. Egypte, 60: 191-196.
- Hafez, M.; A.M. Ali; Khalil, E.M. and F. Abd- El Kawi (1975):** Fluctuation of population of certian predators in cotton and clover fields in Assiut Governorate (Egypt). Agric. Res. Rew., Egypte, 53 (1): 74-51.
- Hamed, M.; R. A. Khan and F.F. Jamil (2006):** Toxicity of different insecticide mixtures against cotton bollworm. Pakistan Journal of Zoology; 38 (1): 39-42.
- Hango, K. and S. Uthamasamy (1989):** Influence of spacing and fertilizer levels on the incidence of bollworm boll rot complex in three varieties of cotton. Applied Agric. Res., 4(3): 173-178.
- Hango, K. and S. Uthamasamy (1992):** Effect of graded levels of fertilizer and methods of application on bollworms boll rot complex in cotton. Modras Agricultural Journal, 79(1): 51-54.
- Henderson, C.F. and E. W. Tilton (1955):** Tests withacaricides against the brown wheat mite. Journal Econ. Entomol., 48: 157-161.
- Henneberry, T. J. (2003):** History of pink bollworm (lepidoptera: Gelechiidae) control in the southwestern cotton growing areas of U.S.A. Recent research developments in plant science, (1): 111-123.

- Herbert, D.; P. J. Phipps and R. E. Strange (1971):** Determination of total carbohydrates, Methods in Microbiology 5 (8): 290-344.
- Hussain, T.; G. Jilani; M. Jamil and M. A. Haq (1999):** Role of EM to sustain crop production in Pakistan. Fifth international conference on Kyusei-Nature Farming, Bangkok, Thailand 23-26 October, 1997, 1999, 165-174.
- Ilango, K. and S. Uthamasamy (1989):** Biochemical and physical basis of resistance to bollworms complex in cotton varieties. Madras Agricultural Journal, 76 (2): 73-76.
- Jaglan, R. S. (1986):** Population build up of bollworm on ratoon cotton. Indian J. Plant Protection, 13 (2): 109-110.
- Jai-Singh and B.S. Sandhu (1995):** Economic threshold for chemical control of spotted bollworms (*Earias vittella* and *E. insulana*) and pink bollworm (*Pectinophora gossypiella*) on tree cotton (*Gossypium arboreum*). Indian Journal of Agricultural Sciences, 65(8): 613-616.
- Joginder, Singh, A. S. Sidhu and J. Singh (1995):** Schematic model of hirsutum cotton phenology and pink bollworm incidence for pest management. Indian, J. Ecol, 10(2): 310-315.
- Khoja, S. M. T.; G. N. Rezk; A. R. Madiha and H.E. Hanafy (2005):** Evaluation of *Bacillus thuringiensis* formulations against the pink bollworm *Pectinophora gossypiella*. Arab Universities Journal of Agricultural Sciences, 13 (3): 929-938.

- Khoja, S. M. T.; G.N. Rezk; A. R. Madiha and H.E. Hanafy (2006):** Effect of *Bacillus thuringiensis* commercial formulations against *E. insulana* (Lepidoptera: Noctuidae). Annals of Agricultural Science, Cairo, 51(1): 261-269.
- Khurana, S.; A. D. Khurana and A. N. Verma (1994):** efficacy of synthetic pyrethroids and conventional insecticides in controlling pink bollworm and their effect on plant phenology, yield and quality of seed cotton. Indian Journal of Entomology, 56(1): 28-35.
- Kim, M. J. and D. M. Oosterhuis (1998):** Changes in the cotton boll wall with age in relation to bollworm and boll weevil feeding habits. Proceedings Beltwide cotton conference, San-Diego, California, USA, 5-9 January-1998, (2): 1367-1368.
- Klein, C. D.; D. R. Johnson; H. B. Myers and Page (1996):** Evaluation of *Bacillus thuringiensis* and ovicide combinations for bollworm and tobacco budworm. Proceedings-Beltwide cotton conferences, Nashville, TN, USA, January 9-12, 2: 798-799.
- Kostandy, S. N. (1992):** Effect of deflowering on cotton boll infestation with the pink bollworm *Pectinophora gossypiella*. Ent. Bull. Soc. Soc. Egypt, (70): 45-49.
- Lohar, M. K. and Y. M. Nahyoon (1995):** Comparative efficacy of Hostathion, Sumicidin and Sevin XLR against cotton bollworms under field condition. Sarhad Journal of Agricultural, 11(3): 363-368.

- Lu- Zhengsong, L. U.; H. U. Siyan and D. Aping (1999):** A discussion on the technique of applying EM bacteria fertilizer to cotton. China- Cottons, 26 (7): 30-31.
- Luo, S. B.; J.P. Yan, C. J. Chai, S. P. Liang; Y.M. Zhang; Y. Zhanf and G. K. Le (1986):** Control of pink bollworm *Pectinophora gossypiella* with *Bacillus thuringiensis* in cotton field J. Biol. control, 2(4): 167-168.
- Mahar, A. N.; M. Munir; D.A. Darban; A. Q. Mahar and G. S. Markand (2004):** Comparative efficacy of different insecticides against *E. insulana* and *Earias vitella* on different cotton varieties in Sindh, Pakistan. Induc-cottons; 1(2): 80-87
- Makkar, A.W. and S. N. Kostandy (1995):** Relationship between spiny bollworm males attracted to pheromone traps and larval infestation in cotton bolls. Annals of Agric. Sci., Moshtohor, 33 (4): 1529-1538.
- Makram, E.A. (1988):** Effect of foliar nutrition, feeding rates and soil fertilization on growth, yield and some fiber properties of cotton cultivar pan 575. Annals of Agricultural Science, Cairo, 33(2): 917-930.
- McConnell, J. S.; W. H. Baker and C. S. Snyder (1999):** Foliar nitrogen fertilization of cotton in Southeastern Arkansas. Research Series Arkansas Agricultural Experiments Station, (463): 38-44.
- Men, U.B.; S.K. Mundiwale; O. R. Gangade and M.N. Bole (1985):** Chemical control of pink bollworm, *Pectinophora gossypiella*. PKV-Research Journal, 9(2): 81-82.

- Mirnoayedi, A. (2001):** The effect of experimental release of eggs and larvae of *Chrysoperla carnea* insect, Neuroptera, Chrysopidae on the reduction of the attack of *E. insulana* on cotton bolls, in Kermanshah, Iran. The first.
- Moawad, G. M.; Z.R. Sawires; M. A. El-Hamaky and M. F. Gergis (1991):** The impact of sex phermonoes and insecticide on the natural enemies in cotton fields in middle Egypt, Bull. Ent. Soc. of Egypt, Economic Series, (19): 237-242.
- Nagewekar, S. N.; K. R. Saini and V. G. Munshi (1984):** Effect of new insecticides on pink bollworm incidence, yield and fiber properties of American cotton H 777. Journal of the Indian Society for Cotton Improvement, 9 (1): 18-22.
- Nageib, M. A. (1980):** Effect of the predators' population in clover fields and their population in adjacent cotton fields. Proc. 1<sup>st</sup> Conf. Pl. Prot. Res. Ins., A.R.C., Cairo, 309-321.
- Nasef, M. A. M. (1990):** Control of certain cotton pests. M. Sc. Thesis, Fac. Agric. Pesticides, Zagazig University.
- Ounchaichon, K.; K. Jee-Rajunya and S. Sirisingh (1996):** Effect of EMS (effective microorganisms) on cotton bollworm. Kasetsart Journal, Natural Sciences, 30 (5): 83-86.
- Paech, K. and M. V. Tracey (1955):** Modern methods of plant analysis. Edited by K. Paech, N. Y. Tracey. Springer-Verlag Berlin Gottingen. Heidelberg. Vol. 4, 766 pp.

- Patel, M. C. and R.N. Vyas (2000):** Field bioefficacy of *Bacillus thuringiensis* var. *kurstaki* and neem based formulations against cotton bollworms. Indian Journal of Plant Protection, 28 (1): 78-83.
- Pawar, V. M.; G. D. Gadhav and S. P. Shirshikar (1987):** Bioefficacy and residues of permethrin in controlling bollworms in NHH-1 cotton. Journal of Maharashtra Agricultural Universities, 12 (3): 334-337.
- Pregl. F. (1945):** Quantitative organic micro analysis 4<sup>th</sup> ED. J. and Achurnil, London, 215.
- Ram. S.; M.P. Gupta and R.P. Maurya (1987):** Role of major plant nutrients (N, P, and K) in the management of insect pests of cowpea (*Vigna unguiculata* (L.) Walp.) International Journal of Tropical Agriculture; 7 (1): 90-96.
- Salem, M. S. (2002):** Evaluation to the efficacy of some recent approaches used for controlling pink and spiny bollworms in the field. M. Sc. Thesis, Fac. Agric. Moshtohor Univ, Zagazig., 137 pp.
- Sarug, D. G. and U.S. Sapute (1988):** Efficacy of some synthetic insecticides and *Bacillus thuringiensis* Var. *kenyae* against bollworms of cotton. Pak. Res. J., 12(2): 119-122.
- Sharma, G.C.; V. T. Gajbhiye; N. P. Agnihotri and H. K. Jain (1984):** Efficacy of some insecticides against cotton bollworms and persistence of their residues in cotton seed and lint. Pesticides, 18 (11): 10-13.

- Sharma, G. C.; V. T. Gajbhiye; N. P. Agnihotri; H. K. Jain and K. N. Katiyar (1986):** Efficacy of cypermethrin, decamethrin, permethrin and fenvalerate against cotton bollworms. *Pesticides*, 20 (4): 40-43.
- Sharma, P. K.; M. Singh and A. K. Dhawan (1989):** Management of bollworm in *Gossypium arboreum* L. through nutrient and irrigation in Punjab. *J. Res. Punjab Agric. Univ.*, 26 (2): 204-205.
- Simwat, G. S.; A. K. Dhawan and A. S. Sidhu (1987):** Effect of delayed sowing, higher nitrogen levels and plant population on incidence of bollworms and yield of seed cotton in F414 hirsutum cotton. *J. Res. Punjab Agric. Univ.*, 24(2): 247-254.
- Snedecor, G. W. and W. G. Cochran (1980):** Statistical methods. 7<sup>th</sup> Ed. Iowa State Univ. Press. Ames Iowa, USA.
- Sorour, F. A.; S. Abou-Khadrah and E. Abou-Ahmed (1967):** Effect of foliar spraying of urea, superphosphate and microelements on growth, flowering, boll setting and yield of cotton. *Egyptian Journal of Agronomy*, 11(1-2): 13-24.
- Srivastava, K. P.; V. T. Gajbhiye and H. K. Hain (1983):** Efficacy of some synthetic pyrethroids against the bollworms and their residues in cotton. *Indian Journal of Agricultural Sciences*, 53 (12): 1048-1051.
- Srivastava, K. P.; V. T. Gajbhiye; H. K. Jain; N. P. Agnihotri and Munshi (1983):** Efficacy of some synthetic pyrethroids against the bollworms and their residues in

cotton. Indian Journal of Agricultural Sciences, 53(12): 1048-1051.

**Stam, P. A. and H. Elmosa (1990):** The role of predators and parasites in controlling populations of *Earias insulana* and *Bemisia tabaci* on cotton in the Syrian Arab Republic. Entomophaga, 35(3): 315-327.

**Sukhija, H. S. and G. J. Reddy (1993):** Time of initiating insecticidal application for the control of *Pectinophora gossypiella* J. Res. Punjab Agric. Univ., 20 (1): 47-56.

**Sukhoruchenko, G. I.; V. B. Mitrofanov and E. V. Vikar (1986):** Comparative estimation of mortality factors of the cotton moth against a background of insecticides of different chemical groups. Problemy izbiratel nosti deistviya insektitsidov-I- akaritsidov-I-ee- znachenei-v-rastenii, 121-127.

**Taneja, S. L. and A. S. Dhindwal (1982):** Bollworm incidence as affected by sowing date, nitrogen application and plant population in up land cotton. Indian J. of Plant Protection. 10 (1-2):1-6.

**Yadav, G. S.; T. S. Kathpal; P. D. Sharma and G. Singh (1993):** Relative efficacy of synthetic pyrethroids against bollworm pests of cotton and their residues in cotton seed. Journal of Insect Science, 6 (1): 92-96.

**Youssef, E.A. (1997):** Damage and damage loss assessments of certain insect pests attacking cotton in Egypt. Ph.D. thesis of Agric., Ain Shams Univ., 160 pp.