




References

VII. REFERENCES

- Atallah, M.A; Aly, F.K.; Eshbah, M.M. (1989):** Pollen gathering activity of worker honeybees, on field crops and medicinal plants in Minia region. Middle Egypt. Proceedings of the fourth International Conference on Apiculture in Tropica Climates, Cairo, Egypt, 6-10 November 1988. 1989, 109-115; Bdo.
- Barth, O.M. (2004) :** MELISSOPALYNOLOGY IN BRAZIL: A REVIEW OF POLLEN ANALYSIS OF HONEYS, PROPOLIS AND POLLEN LOADS OF BEES. Sci. Agric. (Piracicaba, Braz.), v.61, n.3,p.342-350
- Betts, A. D. (1928) :** Pollen supply : broad rearing and nectar flow. Bee zld. 9: 23-25.
- Cakmak, I; L. Aydin; S. Camuzine and H. Wells (2002):** Pollen traps and walnut leaf smoke for Varroa control. American Bee Journal, May,367-370.
- Cano, C.B. (2003) :** Characterization of eucalyptus and citrus monofloral honey in São Paulo State by pollen and physical – chemical analysis. São Paulo; Rev. Inst. Adolfo Lutz, 62(1): 63
- Carvalho, C. A. L.; A. C. Moreti ; L. C, Marchini,.; R. M, Alves, and P.C.F, Oliveira. (2001):** POLLEN SPECTRUM OF HONEY OF “URUÇU” BEE. Rev. Brasil. Biol., 61(1): 63-67

- Chaubal, P.D. (1980) : Melittopalynological studies of shivaji university Campus. Part I : Floral calendar. Journal;- of- Shibaji- University. 1980-1981, 20: 89-94; Bb.**
- Crane, E.(1976) : Bee products : harvesting pollen from hives. Bee World, 57: 20-25.**
- D'Albore , G.R. (1977) : Origine géographique de la galee royale . Apidologie , 9 (1) , 1-17**
- Deans, A.S.C (1963): Beekeeping Technques, Oliver&Boyd. Edinburg and London . 160 p.p.**
- Diaz- Losada, E.; Ricciardelli, d'- Albore, G. and Saa- Otero, M.P. (1998): The possible use of honeybee pollen loads in characterizing vegetation. Grana. 1998, 37:3, 155-163; 26 ref.**
- Duff, S.R. and B. Furgala (1986): Pollen trapping honeybee colonies in Minnesota. American Bee Journal. 126: 755-758.**
- Erickson, E. TR, Young , A. M. and Erickson, B. J. (1988): Pollen collection by honeybees (Hymenoptera : Apidae) in Costarican Cacao (Theobroma Cacao) plantation . J. Apic. Res. 27(3): 190- 196.**
- Floyd E. Moeller (1972): Honeybee collection of corn pollen reduced by feeding pollen in the hive. American Bee Hournal, June 1972, 210-212.**
- Free , J. B. (1967) :Factors determining the collection of pollen by honeybee foragers. Anim , Behave . 15(1): 134-144.**

- Free, J. B. and Williams, I.H (1971):** The effect of giving pollen and pollen supplement to honeybee colonies on the amounts of pollen collected. *J. Apic. Res.* 10(2): 87-90.
- Freitas, B.M. (1994) :** Pollen identification of pollen and nectar loads collected by Africanized honey bees in the state of Ceara, Nrazil. *Proceedings of the fifth International Conference on Apiculture in Tropical Climates, Trinidad and Tobago, 7-12 September 1992.* 1994, 73-79; Bdo.
- Garg, A. (1996) :** Palynocontents of bee- collected pollen loads of autumn season in Bhimal, Indoa. *Taiwania*, 1996, 41: 3, 197-207, Bc.
- Garg, A. and Nair, P.K.K. (1993) :** Honey bees pollen loads as a bioindicator of bee pasturage in Bhimtal area of western Himalayas, India. *Hournal- of- Palynology.* 1993, 29: 89-107; Bb.
- Garg, A. and Nair, P.K.K. (1994):** Seasonal analysis of honey bee pollen loads from a temperature zone in kumaon Himalaya in Uttar Pradesh, India. *Journal – of – Palynology.* 1994, 30: 1, 1-34; Bb.
- Gimenes,M.(1991):** Some morphological adaptations in bees (Hymenoptera, Apoidea) for collecting pollen from *Ludwigia elegans* (Onagraceae). *Revista- Brasileira- de- Entomologia.* 1991, 35: 2, 314-422; Ba.
- Goyal, N. P. (1974):**Apis cerana indica as complementary to each other for the development of apiculture . *Bee world* 53(3) 98- 101.

- Hart , T. and Nabors ,R. (1999):** Pollen Traps as a method of Varroa control. American Bee J. vol. 139 (5):366-367.
- Hart, T. and Nabors, R.(2000):** Pollen traps and drone pupae destruction as a method of Varroa control. Apiculture Research. Vol, 140. No. 2,P,151.
- Harter, B; Leistikow, C; Wilms, W; Truylio, B and Engles, W. (2002):** Bees collecting pollen from flowers with poricidal anthers in a south Brazilian Araucaria forest: a community study. Journal of Apiculture Research 40 (1-2): 9-16.
- Haydak, M.H.(1970):** Honeybee nutrition. Ann. Rev. Entmol. 15: 143-156.
- Hodges, D. (1952):** The pollen loads of the honeybee, Bee Research Association Limited.
- Horskins, K. and Turner, V.B. (1999):** Resource use and foraging patterns of honeybees, *Apis mellifera* and native insects on flowers of *Eucalyptus costata*. Australian – Journal- of – Ecology. 1999, 24: 3, 221-227; 25 ref.
- Hyde , h. A and Adams , K. F. (1958) :** An atlas of airborne pollen Erains: vii-ix. macmillan & Co Ltd. London.
- Jhajj, H. S. and Eoyal, N.P.(1979):** Comparative behaviour of pollen foragers of *Apis cerano indica* and *Apis mellifera* . J. Apic. Res. 18(4): 279-284.
- Jones, G. D., V. M. Bryant, (2004):** The use of ETOH for the dilution of honey.– Grana 43: 174–182. ISSN 0017-3134.

- Kaya ,Z. ; R.Binzet and N.Orcan (2005):** Pollen analysis of honeys from some regions in Turkey. *Apicta* 40 , 10-15
- Kerkvliet, J.D. and Beerlink, J.G. (1991):** pollen analysis of honeys from the coastal plain of Surinam. *Journal of Apiculture Research* 30 (1): 25-31.
- Kerkvliet, J.D. and H.J. Meijer, (2000) :** Adulteration of honey: relation between microscopic analysis and $\delta^{13}C$ measurements. *Apidologie* 31 (2000) 717–726.
- Khattab, M.M.(1979):** Effect of ecological factors on honeybee activities. M. Sc. Thesis, Fac. Of Agric. Cairo Univ.
- Khattab, M.M.(1981):** Ecological and physiological studies on royal jelly of honeybees (*Apis mellifera* L.). PH.D, Zagazig Univ. (199) pp.
- Khattab, M.M.(2000):** Honeybees for people treatments. NPHD.Fac. Agric. Moshtohor, Egypt, 270 pp.
- Kirk, W. (1994):** A colour guide to pollen loads of the honey bee. International Bee Research Association; Cardiff; UK> 1994, 54pp.; Bd.
- Klungness, L.M.; Peng, Y.S. (1983):** A scanning electron microscopic study of pollen loads collected and stored by honeybees. *J. Of Apicultural, Research*. 1983, 22:4, 264-271; B.
- Loper, G. M.(1986): Cotton pollen :** Honeybee avoidance and absence of gossypol. *Jour., Economic Entomology*, 79(1):103-106.

- Mahmood, A.N.; Waller, G.D.; Hagler, J.R. (1990):** Dispersal of upland and Pima cotton pollen by honey bees (Hymenoptera: Apisae) visiting upland male-sterile flowers. *Environmental- Entomology*. 1990, 19:4, 1034-1036; Bb.
- Marcella , B. Bernardini (1978) :** Origine Eeographic of Royal jelly . *Apidologia* , 9(1): 1-17.
- Marchini, L.C.; Moreti, A.C. de, C.C.; Teixeira, E.W.; Oliveira, P.C.F., de; de C.C. Moreti, A.C.; and de, Oliveira, P.C.F. (2000):** Identification on pollen loads by Africanized honeybees (*Apis mellifera* L.) of different colonies places in the same surrounding. *Ecossistema*. 200, 25: 1, 48-51, 14 ref.
- Maurizio, A. (1975):** **Microscopy of honey:** PP 240-257 from honey: acomprehensive survey . ed. E. Crane. London : Heinemann in co-operation with IBRA.
- McLellan, A.R. (1974):** some effects of pollen traps on colonies of Honeybees. *Journal of Apiculture Research* 13 (2): 143-148.
- Michener, C. D.(1979) :** The biography of the bees. *Ann. Missouri Bot. Garden* 66: 277-347.
- Moeller, F. E. (1972):** Honeybee collection of Corn pollen reduced by feeding pollen in the hive. *Am. Bee J.* 112: 210-212.
- Nabors, R.A (1997):** Trapping pollen collections of the honeybee, *Apis mellifera* L., to determine pollen flow periods. *American Bee Journal*. Vol 137 No. 3.

- Nagamitsu, T. and Inoue, T. (1999):** Differences in pollen sources of *Apis cerena* and *Apis mellifera* at a primary beech forest in central Japan. *Journal of Apiculture Research* 38 (1-2): 71-78.
- Nolan, W. J.(1925):** A two years brood rearing curve for a single colony. *J. Econ. Ent.* XVI, 117-129.
- O'Neal, R. Jand Waller , E. D. (1984):** On the pollen harvest by the honeybees (*Apis mellifera* . L) Near Tucson, Arizona (1976-81). U.S.D.A. Agricultural Research service car Hglyden Bee Research Center, Tucson, Arizona . 81-109.
- Percival, M. S. (1947):** Pollen collection by *Apis mellifera*. *New Phytologist* 46: 142-165.
- Pernal, S.F. and Currie, R.W. (2001):** The influence of pollen quality on foraging bahviour in honeybees (*Apis mellifera* L.) Bevahioral – Ecology and Sociobiology. 2001, 51: 1, 53-68.
- R. D Albore , M. Battaglini and D Ambrosio M. (1977):** Considerations on pollens of Royal Jelly . *Annali Dell istituto spermentale La Zoologia Agraria* Vol. V, 1976. 1977 pp. 203- 213 , 2 figg. 3, ta - vv f.t. pubblicato il 4 aprile 1977.
- Ranjhan, S.K. and Krishna, G. (1980):** Laboratory Manual for Nutrition Research. Vikas Puplishing House Pvt LTD, New Delhi India.

- Rao, S.P.C.; Atluri, J.B. and Rao, C.S. (1995):** *Sterculia foetida* as a pollen source for the Indian honey bee, *Apis cerana indica*. Indian- Bee – Journal. 1995, 57: 1, 18-19; Bj.
- Rashad, S. E. and R. L. Parker (1956) :** Major pollen sources in the Manhattan, Kansas area and the influence of weather factors upon pollen collection by honeybees. Proc.Tenth Int, Congr. Entomo. 4: 1037-1046.
- Real, L.R. (1983):** *Polination biology*. Academic Press., London. 338 pp.
- Roubik, D. W. & C. D. Michener, (1980) :** The seasonal cycle and nests of *Epicharis zonata*, a bee whose cells are below the wet-season water table (Hymenoptera, Anthophoridae). *Biotropica* 12(1): 55-60.
- Roubik, D. W. (1982b) :** The ecological impact of nectar-robbing bees and pollinating hummingbirds on a tropical shrub. *Ecology* 63: 354-360.
- Schmalzel, R.J. (1980) :** The diet breadth of *Apis* (Hymenoptera: Apidae). M. S thesis, Univ. Arizona. Tucson. 79 pp.
- Schwarz, H.F. (1948):** Stingless bees (Meliponidae) of the western hemisphere. *Bulletin of the American Museum of Natural History* 90: i- xvii + 1-546.
- Seeley, T. D. (1985):** *Honeybee ecology a study of adastatim in social life* Princeton : Princeton univ. press.
- Seiji, M.C.; Aira, M.J.; Iglesias, M.I. and Jato, M.V. (1994):** Foraging activity of the honey bee on *Actinidia deliciosa*

Chev. as shown by pollen analysis. Grana, 1994, 33: 4-5, 286-291; Bb.

Shannon, C.E. and W. Weaver, (1949): The mathematical theory of communication. Univ. Illinois Press. Urbana. 117 pp.

Shimanuki, H.(1987): Varroa mites found in the United States. American Bee Journal. 125(11), 745-746.

Simpson, J.(1955): The significance of the presence of pollen in the food of the worker larvae of honeybees. Quarterly J. of Microscopic Science. 96: 117-120.

Smith, M. V. and A. Adie (1963) : A new design in pollen traps. Can. Bee.J. 74: 4,5,8.

Snedecor, G.W. and Cochran, W.G. (1971): Statistical Methods. Iowa State Univ. Press, Ames, USA. Sixth edition, 4 th printed.

Spencer-Booth, Y. (1960): Feeding pollen, pollen substitutes and pollen supplements to honeybees. Bee World 41: 253-263.

Synge, A. D. (1947): Pollen collection by honeybees (*Apis mellifera*). J. Anim. Ecol.. 16: 122-138.

Thajj, H. S. and Eoyal, N. P. (1979): Comparative behavior of pollen foragers of *Apis cerana indica* and *Apis mellifera*. J. Apic Res. 18(4):279-284

Thans, P; Kalpana, T; Ramanujam, C; (1991): Pollen analysis of rock bee summer honeys from the Prakasam district of the Andhra Pradesh, India. Journal of Apiculture Research 30 (1): 33-40.

- Todd , F.E. & Reed , C.B. (1970) :** Brood measurement as a valid index to the value of honeybees as pollinators . J. Econ. Ent. 63(1) : 148-149.
- Todd, F. E. and R. K. Bishop, (1946) :** The role of pollen in the economy of the hive. U. S. Bur. Ent. and Plant Quar. E-536, Rev., 7 pp.
- Traynor , J.(2001) :**Evaluating Pollen Production of plants (with sample calculations for almonds) American Bee J.Vol.141 No. 4:287-288.
- Traynor. J. (2001):** Evaluating pollen production of plant. American Bee Journal Vol. (4) No.4.
- Vaissiere, B.E. (1988):** A novel method for quantification of pollen production and pollen loads on bees and stigmas using a coulter counter. American- Bee- Journal. 1988, 125: 12, 810; Bj.
- Villanueva, G.R. (1989):** Important plant species for apiculture in Ejido plan del Rio, Veracruz, Mexico. Proceedings of the Fourth International Conference on Apiculture in Tropical Climates, Cairo, Egypt, 6-10 November 1988. 1989, 138-145, Bdo.
- Villanueva, Gutierrez, R. (1995):** Food resources of European and Africanized honeybees (*Apis mellifera* L.) in the eastern Yucatan Peninsula, Mexico. 1995, vii + 167 pp.; Bdo.
- Vogel, S. (1974) :** Ölblumen und ölsammelnde Bienen. Tropische und Subtropische Pflanzenwelt [Mainz], no. 7: 283-547.

- Von Der Ohe, W; L.P, Oddo; M.L.Piana ; M. Morlot and P. Martin (2004):** Harmonized methods of melissopalynology. *Apidologie* 35 (2004) S18–S25.
- Von Frish, K (1967):** The dance language and orientation of bees. Harvard univ. Press. Cambridge.
- Wahl, O. (1963) :** Vergleichende untersuchungen über nährwert von pollen, hefe, soja-mehl, und trockenmitch für die honigbiene (*Apis mellifera*); ein beitrage zum eiweiss und vitamin stoffwechsel der biene. 2. Bienenforsch. 6: 209-280.
- Waller, G. D. (1980) :** A modification of the Ontario Agricultural college pollen trap. *American Bee Journal* 120(2): 119-121.
- Waller, G. D; D.M. Caron and G. M. Loper (1981):** Pollen patties maintain brood rearing when pollen is trapped from honeybee colonies. *Amer. Bee J.*, 121(2): 101-103,105.
- Winston, M.L. (1987):** The biology of the honeybee. Harvard Univ. Press. London. 281 pp.