

CONTENTS

	Page
I. INTRODUCTION	1
II. REVIEW OF LITURATURE	3
1. Plant height	3
2. Stem diameter	4
3. Third leaf angle and chlorophyll content	5
4. Varietal performance	6
5. Flowering date	6
6. Effect of sowing dates on the rate of infestation by corn borers	7
a- <i>Sesamia cretica</i> Led	7
b- <i>Ostrinia nubilalis</i> , Hub	12
c- <i>Chilo Agamemnon</i> , Bels	15
7. Susceptible of varieties to infestation	17
a- <i>Sesamia cretica</i> Led	17
b- <i>Ostrinia nubilalis</i> , Hub	22
8. Damage caused by corn borers on maize yield	28
9. Leaf feeding infestation	36
10. Number of cavities per plant	36
11. Number of holes per 100 internodes	38
12. Number of larvae per plant	39
13. Infested ears	41
14. Stored corn	43
III- MATERIALS AND METHODS	45
1. Reading on borers	46
a- Anatomy of stalk	46
b- Number of infested plants by borers	47
c- Number of died hears	47
2. Agronomic characters	47
IV- RESULTS AND DISCUSSION	51
Part I:	51
1. Botanical characters	51
1.1. Plant height and stem diameter	51
1.2. Third leaf angle	56

	Page
1.3. Leaf chlorophyll content	56
1.4. Husk length	56
1.5. Not covered ears %	56
1.6. Moisture percentage	60
2. Yield and Yield components	60
2.1. Effect of season	60
2.2. Effect of planting dates	63
2.3. Varietal performance	63
2.4. Interaction between planting dates and season	66
3. Store insects	66
Part II: Evaluation of certain corn genotypes for infestation by stem borers	68
1. The effect of planting date on the degree of infestation with <i>Sesamia cretica</i>	68
1.1. Dead heart symptoms	68
1.2. Mean percentage of pink borers infestation	70
1.3..Relationship between different plant height and degree of <i>S. cretica</i> infestation	72
1.4..Relationship between plant stem diameter and mean percentage of infestation	75
1.5. Effect of <i>S. cretica</i> infestation on different maize genotypes at 50% flowering date	78
1.6..Effect of the third leaf angle of genotypes on percentage of <i>S. cretica</i> infestation	80
1.7..Relationship between leaf characters and <i>S.</i> <i>cretica</i> infestation at 32 days old	83
1.8..Effect of leaf chlorophyll content on the percentage of dead heart caused by <i>S. cretica</i>	87
2. The effect of corn planting date on the degree of infestation with <i>Ostreia nubilalis</i> and <i>Chilo</i> <i>Agamemnon</i>	89
2.1..Relative susceptibility of certain genotypes of maize to natural infestation with <i>O. nubilalis</i> and <i>C. agamemnon</i>	89

	Page
2.2..Relationship between mean percentage of infestation, plant height and stem diameter at 75 days	91
3. Effect of three borers infestation on different genotypes at harvest time	93
3.1..Relationship between plant height and phenomenons of infestation at harvest time	93
3.2..Relationship between stalk diameter and some phenomenons of infestation at harvest time	99
3.3..Relation between borers infested levels and weight of ears harvest time	104
3.4..Field evaluation of maize genotypes at harvest time for their susceptibility to <i>S. cretica</i> , <i>O. nubilalis</i> and <i>C. agamemnon</i> under natural infestation	107
3.5. Relationship between number of borers larvae and ear weight at harvest time	110
4..Susceptibility of maize grains to infestation with some storage insects	111
4.1. <i>Sitotroga cerealella</i>	113
4.2. <i>Sitophilus oryzae</i>	113
4.3. <i>Rhizopertha dominica</i>	114
V- SUMMARY	115
VI- REFERENCES	129
VII- ARABIC SUMMARY	