



REFERENCES

VI. REFERENCES

- Abd El-Gayed, A.A. (1995):** Studies on some maize insect pests and their natural enemies. M.Sc. Thesis, Fayoum Fac. Agric., Cairo Univ.
- Abd El-Rahim, M.M.; M.I. Abdel-Fattah; A.I. Farag and M.A.Z. El-Naggar (1991):** Relative susceptibility of maize cultivars to infestation by corn borers, *Sesamia cretica* Led. and *Ostrinia nubilalis* (Hub.) in relation to the sowing date. Minufya, J. Agric. Res. 16 (1): 843-857.
- Abou-Khadrah, S.H. (1968):** The effect of time of planting and varieties on the yield of corn and its chemical constituents. M.Sc. Thesis, fac. Agric., Cairo Univ.
- Abukar, M.M.; G. Burgio and E. Tremblay (1986):** Evaluation of post harvest losses caused by insects to maize in three districts of southern somalia. Bollettino del laboratorio di Entomol. Agraria "Filippo Silvestri", Italy. 43 : 51-58.
- Ahmed, M.K. and M.T. Kira (1960):** Studies on corn borers and their control. Egyptian Agricultural Organization, 44: 1-78 (In Arabic).
- Ahmed, M.M.M. (2000):** Studies on the important insect pests of maize plants and their natural enemies at Kafr El-Sheikh District. Ph.D. Thesis, Fac. Agric. Tanta Univ.

- Ahmed, M.T. (1980): Preliminary field observations on the preference of stem borers to different grainous host plants. Bull. Soc. Ent. Egypte. 63: 219-222.
- Ali, M.A.; M.M. Metwally; M.M. Abd El-Ghaffar and A.S. Metwally (1989): Susceptibility of four local maize varieties to corn borer infestation under natural condition in Egypt. Acta phytopath. et Entomol. Hung., 24 (1-2): 13-16.
- Al-Jassany, R.F.; H.C. Ali and M.A. Younis (1987): Estimation of the percentage of infestation by stem corn borer, *Sesamia cretica*. Led. for several varieties of maize Iraqi. J. Agric. Sci. ZANCO., 5(1): 223-229.
- Aly, A.M. (1983): Studies on the occurrence of suckers in maize. M.Sc. Thesis, Fac. Agric., Moshtohor, zagazig Univ.
- Anglade, P.; F. Baca; H. Berger; H.C. Chiang; D. Mustea and D. Hadzistevic (1981): Assessment of the susceptibility of maize inbred lines to the European corn borer (*Ostrinia nubilalis* Hbn.) as part of the programme of IWGO, 1977-1980. Proble de protectia plantelor 9 (2): 147-177.
- ; A. Monteagudo and IWGO Co-operators (1980): Assessment of the susceptibility of maize lines to the European corn borer, *Ostrinia nubilalis* as part of the programme of IWGO, 1975-1976. EPPO Bull, 10 (3) 365-369.

- Anonymous (1977):** Estimates of damage by the European corn borer grain corn in the United States in 1976. Coop Plant Pest Rep. 2: 375-376.
- Atiyeh, R.; M. Aslam and R. Baalbaki (1996):** Nitrogen fertilizer and planting date effects on insect pest populations of sweet corn. Pakistan J. Zool., 28 (2): 163-167.
- Awadallah, W.H. (1984):** Evaluation of certain local exotic corn material for relative susceptibility to infestation with *Ostrinia nubilalis* (Hubner), in Egypt. Bull Soc. Ent. Egypt, 65: 65-72.
- ; **A.A. Galal and W.D. Guthrie (1982):** Genetic analysis of resistance to the European corn borer, *Ostrinia nubilalis* (Hubner) in diallel crosses of maize in Egypt. Alex. J. Agric. Res., 30 (1): 345-356.
- Baker, J.E. (1988):** Development of four strains of *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) on barley, corn (maize), rice and wheat. J. Stor. Prod. Res. 24 (4): 193-198.
- Ba-Momen, A.M. (1981):** Differential yield response of corn varieties and hybrids to different sowing dates. M.Sc. Thesis, Fac. Agric., Cairo Univ.
- Barbulescu, A.L. (1979):** Results obtained at Fundulae on the resistance of corn to European corn borer *Ostrinia nubilalis* Hb. attacks Proc. of the COMECON Symp., Cluj-Napoca, September 26-27.

- Barbulescu, A.L. and T. Sarca (1983):** Testing of some hybrid combinations between maize lines tolerant to the European corn borer (*Ostrinia nubilalis* Hb.) Proble Prot. Plant. Vol. Xj Nr. 1.
- Barry, B.D. (1969):** European corn borer and corn leaf aphid resistance in corn hybrids. Ohio Agric. Res. and Dev. Center. Res. Circular 174, October 1969: 1-8.
- Barry, D. and L.L. Darrah (1991):** Effect of research on commercial hybrid maize resistance to European corn borer. J., Econ. Ent., 84 (3): 1053-1059.
- Bigler, F.; M. Waldburger and S. Bosshart (1990):** Yield losses in grain and silo maize caused by the European corn borer, *Ostrinia nubilalis* Hbn. Land wirtschaft Schweiz, 11 (3): 591-596.
- Bisher, M.A. (1967):** Effect of sowing and cultivation methods on the growth and yield of some maize varieties at different planting dates. M.Sc. Thesis, Fac. Agric., Ain Shams Univ.
- **(1973):** Physiological response of maize plants to planting dates. Ph.D. Thesis, Fac. Agric., Ain Shams Univ.
- ; **R.M.M. Abdallah and A.A.E. El-Sayed (1977):** Effect of planting date on grain yield of some maize varieties at different locations in A.R.E. Agric. Res. Rev. 55: 125-135.

REFERENCES

- Cagan, L. and M. Grencik (1990):** Harmfulness of the European corn borer (*Ostrinia nubilalis* Hbn.) on grain maize. Rostlinna Vyroba. 36 (2): 203-212.
- Calvin, D.D.; M.C. Knapp; K. Xingquan; F.L. Poston and S.M. Welch (1988):** Influence of European corn borer (Lepidoptera : pyralidae) feeding on various stages of field corn in Kansas. J. Econ. Ent. 81 (4): 1203-1208.
- Cardoso, M.J.; HWL de. Carvalho; M.de. L.dos. Leal; Mx. dos. Santos; HWL de Carvalho and Mx. dos. Santos (2000):** Behaviour, adaptability and stability of hybrids of corn in the state of Piaui in the agricultural year of 1998 Revista cientifica Rural 5(1): 142-149.
- Cartea, M.E.; R.A. Malvar; P. Revilla; A. Ordas and A. Alvarez (1994):** Seasona; occurrence and response of maize inbred lines to pink stem borer in the northwest of Sppain. Maydica, 39 (3): 191-196.
- Likewise and Cheu, S.P. (1940):** Studies on the planting date of corn in relation to European corn borer infestation. Kwangsi Agric. 1, No. 6: 373-383 (c.f. El-Sherif, S.E.I. Ph.D. Thesis, Fac. Agric. Alexandria Univ. 1965).
- Dobie, P. (1974):** The laboratory assessment of the inherent susceptibility of maize varieties to post harvest infestation by *Sitophilus zeae* maize Motsch. (Coleoptera. Curculionidae). J. Stored prod. Res. (10): 183-197.

- El-Hattab, A.H.; E.M.S. Geith; Sh.A. Shaban and A.A. Bedeer (1986): Plant growth and yield of some maize varieties as affected by some micronutrient and their time of application. Proc. 2nd Conf. Agron., Alex., Egypt. Vol. 1: 277-293.
- El-Naggar, M.A.Z. (1991): Ecological factors influencing some insect pests diversity in maize fields. M.Sc. Thesis, Fac. of Agric., Menoufia Univ.
- (1997): Further studies on greater sugarcane borer *Sesamia cretica* Led. and european corn borer *Ostrinia nubilalis* Hub. Ph.D. Thesis, Fac. Agric., Minoufia Univ. Egypt.
- El-Naggar, Z. (1967): Studies on corn borers and their biological control. M.Sc. Thesis, Assiut University.
- El-Saadany, G.B. (1965): Ecological and biological studies on some maize pests. M.Sc. Thesis, Fac. of Agric., Ain Shams Univ.
- (1969): Further ecological and biological studies on certain corn pests. Ph.D. Thesis, Fac. of Agric., Ain Shams Univ.
- El-Sawaf, S.K.; S.M. Hammad and S. El-Sherif (1964): Preliminary study of the susceptibility of certain corn varieties to infestation with the corn borers in Egypt. Alex. J. Agric. Res., 12 (1): 103-129.
- El-Sherif, S.E.I. (1962): Studies on the corn borers of Alexandria District M.Sc. Thesis, Fac. Agric., Alexandria Univ.

- El-Sherif, S.E.I. (1965):** Studies on the corn borers in the Alexandria district. Ph.D. Thesis, Fac. Agric., Alexandria Univ.
- El-Sherif, S.I. and F.F. Mostafa (1987):** Susceptibility of maize germplasm with the greater sugar-cane, *Sesamia cretica* Led. (Lepidoptera: Noctuidae). 10-Evaluation of the commercial cultivars Giza 2, DC202 and DC204. Arab J. Pl. Prot. 5: 8-13.
- Emara, N.M. (1997):** Ecological studies on certain natural enemies of maize and sorghum pests. M.Sc. Thesis, Fac. of Agric., Cairo Univ.
- Everett, T.R.; H.C. Chiang and E.T. Hibbs (1958):** Some factors influencing populations of European corn borer (*Pyrausta*). *O. nubilalis* (Hub) in the North Central States. Minn. Agric. Exp. Sta. Tech. Bull: pp. 63
- Farghaly, H.T. (1969):** The biology and ecology of certain natural enemies of corn borers. Ph.D. Thesis, Fac. Agric., Ain Shams Univ. p. 182.
- Felt, E.P. (1923):** European corn borer. 35th. Report of the state entomologist, 1921. New York State Mus. Bull. No. 247-248, 129 pp. Albany, N.Y. (c.f. El-Sherif, S.E.I. Ph.D. Thesis, Fac. Agric., Alexandria Univ. 1965).
- Ficht, G.A. (1931):** Some observations of the planting date of corn and its relation to European corn borer population. J. Econ. Ent. XXIV. No. 2: 380-386, Geneva, N.Y. (c.f. El-Sherif, S.E.I., Ph.D. Thesis, Fac. Agric. Alexandria Univ., 1965).

- Frey, R.J. and Y. Malonado (1967):** Relative productivity of homogenous and interogenous oat cultivars in Optimum and suboptimum. Environments. Crop Sci., 7: 520-530.
- Frolov, A.N. and M.A. Chumakov (1990):** Early lines of maize resistance to European corn borer. Seleksiya-I-Semenovodstvo Moskva, 5: 29-30. (c.f.R.A.E., 79, Abst. No. 9882).
- Galal, A.A.; R.M. Abdallah; W.H. Awadallah and N.S. Selim (1980):** Evaluation of some local and exotic materials for the resistance in maize to the European corn borer, *Ostrinia nubilalis* (Hub) and studying the effect of nitrogen fertilizer and number of plant per unit area on the infestation level of the same pest. Proc. 1st Conf. Pl. Prot. Res. Ins., I: 187-196.
- Ghidiu, G.M.; E.C. Berry and P. Du Bose (1979):** Relationship between entrance holes and stalk cavities caused by first and second generation European corn borer in Field corn J. Econ. Ent. 72: 85-86.
- Gouda, A.S. (1982):** Effect of planting density and nitrogen fertilization on growth and yield of some maize varieties. M.Sc. Thesis, Fac. Agric., Zagazig Univ.
- Grier, S.L. and D.W. Davis (1980):** Infestation procedures and heritability of characters used to estimate ear damage caused by second-brood European corn borer (*Ostrinia nubilalis* Hubner) on corn. J. of Am. Soc. for Hort. Sci. 105 (1): 3-8.

- Grier, S.L. and D.W. Davis (1983):** Optimizing European corn borer (Lepidoptera: Pyralidae) infestation of maize ears for kernel damage evaluations. J. Econ. Ent., 76: 429-431.
- Guthrie W.D.; F.A. Onkogu; W.H. Awadallah and J.C. Robbins (1980):** European corn borer: Evaluation of resistance in husk silk tissue of inbred lines of corn. J. Econ. Ent. 1.73. 178-180.
- Hamza, Z.A. (1969):** A comparison of open-pollinated varieties with some hybrids of corn under different environmental conditions. M.Sc. Thesis, Fac. Agric., Ain Shams Univ.
- Hanna, H.M. and T.E. Atris (1969):** Field observation on *Chilo agamemnon* Bles. (Lepidoptera: cerambidae). Bull. Soc. Ent. Egypte, 53: 551-556.
- Hassanien, M.H. (1961):** Studies on certain corn borers in Egypt , U.A.R. the 4th Arab Sci. Cong U.A.R. (1960) .
- **and A.I. Badawy (1959):** The effect of weather factors on corn borers distribution in Egypt. Conf. Cereal'Crops, Cairo, Part II, PP .1560.
- Hervey, G.E.R. (1928):** Life history, habits and control of the European corn borer in Erie and Chautauqua countries. Circ . N.Y. Agric. Expt. Sta., No. 157, Geneva, N.Y. (c.f. El- Sherif, S.E.L. Ph.D. Thesis, Fac. Agric., Alexandria Univ., 1965).

- Hsu, S.L.; W.K. Peng and F.K. Hsieh (1988):** Loss assessment of corn infested with Asian corn borer, *Ostrinia furnacalis* (Guenee). Memoirs of the college of Agriculture. National Taiwan Univ. 28 (2): 27-31.
- Huber, L.L.; C.R. Neiswander and R.M. Salter (1928):** The European corn borer and its environment Ohio Agr. Exp. St. Bul. 429, 196 pp.
- Hudon, M. and M.S. Chiang (1991):** Evaluation of resistance of maize germplasm to the univoltine European corn borer *Ostrinia nubilalis* (Hubner) and relationship with maize maturity in Quebec Maydica 36: 69-74.
- Hudon, M. and M.S. Chiang (1985):** Resistance and tolerance of maize germplasm to the European corn borer *Ostrinia nubilalis* (Hubner) and its maturity in Quebec. Maydica. 30, (3): 329-337.
- Isa, A.L. (1959):** Two short notes on the susceptibility of certain corn varieties to infestations with corn borers and the possibility of controlling these pests through the establishment of a host free period. The Agric. Res. Rev. 37 : 378-383.
- **and W.H. Awadallah (1975):** Biological studies on corn borers in Egypt. 1-Seasonal distribution. Ibid., 53 (1): 53-64.
- ; **W.H. Awadallah and A.M. Tantawi (1969):** The susceptibility of some corn varieties to infestation with corn borers and other corn pests. Ibid., 47 (3): 23-28.

- Isa, A.L.; W.H. Awadallah and H.N. Wanas (1969):** Distribution of overwintering corn borer larvae in residues of maize plants. *FAO plant protection Bul.* Vol. 17 (5): 112-113.
- Ismail, I.I. (1968):** Studies of ecology, biology and control of corn borers in Giza region, Ph.D. Thesis, Fac. of Agri., Cairo Univ., Egypt.
- Ismail, I.I.; M.T. Kira and M.A. Hussein (1974):** Corn yield response to borers infestation and planting dates in Giza region. *Fac. of Agric. Cairo Univ. Res. Bull.* 25: 77-85.
- Jackson, R.D. and D.C. Peters (1963):** Biological observations on the European corn borer in South Western Missouri. *J. Econ. Ent.* 56, No. 6: 741-747, Menesha, Wis. (c.a. El-Sherif, S.E.I. Ph.D. Thesis Fac. Agric. Alexandria Univ., 1965).
- Jarvis, J.L. and W.D. Guthrie (1980):** Resistance of maize plant introductions to sheath-collar feeding by 2nd O. generation European corn borers. *Maydica* 25: 25-32.
- **and W.D. Guthrie (1988):** Effect of first generation European corn borer on yield and plant height of pop corn. *J. Agric. Entomol.* 5 (3): 179-183.
- ; **T.R. Everett; T.A. Brindley and F.F. Dicke (1961):** Evaluating the effect of European corn borer populations on corn yield. *Iowa State J. Sci.*, 36: 115-132.

- Jarvis, J.L.; W.A. Ruseel, J.E. Campbell and W.D. Guthrie (1991): Level of resistance in maize to prevent or reduce yield losses by second-generation European corn borers. *Maydica* 36 : 267-273.
- ; W.D. Guthrie and E.C. Berry (1983): Time and level of infestation by second generation European corn borers on a resistant and a susceptible maize hybrid in relation to yield losses. *Maydica*, 28 (4): 391-400.
- ; ----- and J.C. Robbins (1986): Yield losses from second generation European corn borers (Lepidoptera: Pyralidae) in long-season maize hybrids planted early compared with short-season hybrids planted late. *J. of Econ. Ent.* (79)1: 243-246.
- Jonhson, G.R. and R.J. Frey (1967): Heritabilities of quantitative attributes of Oat *Avena* spp. at varying levels of environmental stress. *Crop. Sci.*, 7: 43-46.
- Kaan, F.; P. Anglade; A. Boyat and A. Panouille (1983): Resistance to the European corn borer, *Ostrinia nubilalis* Hbn. (Lep. Pyralidae) in a diallele of 14 early inbred lines of maize, *Zea mays* L. *Agronomie*, 3 (6): 507-512.
- Keenan, W.N. (1926): The spread and degree of infestation of the European corn borer in 1925. 56th Ann. Rept. Ent. Soc. Ontario: 75-77 (c.a. El-Sherif, S.E.I., Ph.D. Thesis, Fac. Agric., Alexandria Univ., 1965).

- Keller, N.P.; G.C. Bergstrom and R.I. Carruthers (1986):**
Potential yield reduction in maize associated with an anthracnose European corn borer pest complex in New York. *Phytopatho.* 76 (6): 586-589.
- Kim, S.K.; W.D. Guthrie; A.R. Hallauer; W.A. Russell; J.L. Brewbaker and C.S. Hong (1989):** Evaluation of tropical and subtropical corn lines for resistance to second-generation European corn borer (Lepidopta: Pyralidae). *J. Econ. Entomol.* 82. 82 (4) 1245-1250.
- Kira, M.T.; I.I. Ismail, and A.K. Azab (1974):** Determination of the number of annual generations of the pink borers, *Sesamia cretica* Led. in Giza region, Egypt. *Bull. Fac. of Agric., Univ. Cairo*, 25 (1): 87-93.
- Klenke, J.R. (1986):** Recurrent selection for European corn borer resistance in a maize synthetic. Dissertation Abs. International, Bll. Sci. and Engi. 46 (12).
- Klenke, J.R.; W.A. Russell and W.D. Guthrie (1986a):** Grain yield reduction caused by second generation European corn borer in B59 corn syntheric. *Crop Sci.* 26 (5): 859-863.
- ; ----- and ----- (1986b):
Recurrent selection for resistance to European corn borer in a corn synthetic and correlated effects on agronomic traits. *Ibid.*, 26: 864-868.
- Kwolek, W.F. and T.A. Brindley (1959):** The effects of the European corn borer *Pyrausta nubilalis* (Hubn.) on corn yield. *Iowa State Cill. J. Sci.*, 33: 293-323.

- Labatte, J.M. and B. Got, (1991):** Modelling damage on maize by the European corn borer, *Ostrinia nubilalis*. Annals of Applied Biol. 119 (3): 401-413.
- Lisowicz, F. (2000)** Maize protection against aphids and the European corn borer. Ochrona Roslin (2000) 44 (3): 35-36.
- Lutfallah, A.F.; M.R. Sherif and K.A. Moafy (1991):** Further studies on the chemical control of the European corn borer, *Ostrinia nubilalis* (Hubner) on maize. Fourth Arab Congress of Plant Protection, Cairo 1-5 Dec., 160-163.
- Lynch, R.E. (1980):** European corn borer yield losses in relation to hybrid and stages of corn development. J. Econ. Ent., 73 (1): 159-164.
- **and W.D. Guthrie (1980):** Relative susceptibility of corn hybrids to European corn borer damage. Maydica 25: 117-126.
- Lynch, R.E.; J.F. Robinson and E.C. Berry (1980):** European corn borer: yield losses and damage resulting from a stimulated natural infestation. J. Econ. Ent., 73 (1): 141-147.
- Mahmoud, A.S. (1967):** The effect of varieties and dates of planting on the yield and its components in maize. M.Sc. Thesis, Fac. Agric., Cairo Univ.
- Malvar, R.A.; M.E. Cartea; P. Revilla; A. Ordas; A. Alvarez and J.P. Mansilla (1993):** Sources of resistance to

pink stem borer and European corn borer in maize.
Maydica, 38 (4): 313-319.

Mannaa, S.H. (1999): Differential response of corn yield and infestation by the pink borer, *sesamia cretica* Led., as affected by certain agricultural practices in Assiut. Assiut J. of Agric Sci. 30 (5): 7-106.

McMillian, W.W.; N.W. Widstrom; Dean Barry and E.B. Lillehoj (1988): Aflatoxin contamination in selected corn germplasm classified for resistance to European corn borer (Lepidoptera: Noctuidae). J. of Ent. Sci. 23 (3): 240-244.

Metawi, A.A.E. (1996) . Genetic analysis of resistance to corn borers in some inbreds of maize .M.Sc. Thesis , Fac. Agric ., Kafr El - Sheikh , Tanta Univ ., Egypt.

Metwally, A.S. (1988): Relative susceptibility of certain maize varieties to infestation with Lepidopterous insects. M.Sc. Thesis, Fac. Agric., Al-Azhar, Univ., Egypt.

Metwally, A.S. (1995): Studies on the relationship between some geotypes of maize and infestation level of European Corn Borer, *Ostrinia nubilalis*, (Hubn.) in Egypt. Ph.D. Thesis, Fac. Agric., Minofiya Univ., Egypt.

Metwally, S.M.I. (1967): Insect fauna of maize plants with special references to the interrelationship between major pests and their predators and the effect of certain insecticides on the populations. M.Sc. Thesis, Fac. Agric., Cairo Univ., pp. 198.

REFERENCES

- Metwally, S.M.I. (1976):** The relationship between the planting dates of maize and rates of infestation by corn borers. J. Agric. Res, Tanta Univ. 2: 327-331.
- Mohamed, S.M. (1997).** Breeding studies on resistance to corn borer (*Sesamia cretica* Led.) in maize (*Zea mays* L.). Ph . D. Thesis, Fac. Agric., Cairo Univ., Egypt
- Mourad, S.B.; M.A. El-Lakany; A.I. El-Agamy and F.H. Soliman (1986):** Response of some varieties and hybrids of maize to plant densities and nitrogen fertilization. 1. Growth and flowering traits. Al-Azhar J. Agric. Res. VI: 401-412.
- Moustafa, F.F. (1981):** Biological and ecological studies on the pink borer, *Sesamia cretica* Led. (Lepidoptera: Noctuidae). Ph.D. Thesis, Fac. of Agric., Cairo Univ.
- Nawar, A.A.; M.E. Ibrahim and M.B. Attia (1992):** Grain yield, yield components and infestation rates of corn borers and aphid of maize genotypes as influenced by nitrogen fertilization. Egyptian J. of Agronomy 17(1-2): 41-58.
- Ordas, A.; A. Alvarez and R.A. Malvar, (1988):** Variability of resistance to the second generation of European corn borer in maize. Maydica. 33 (1): 27-36.
- Parag-Agarwal; S.S. Verma; S.N. Mishra and P. Agarwal (2000):** Phenotypic stability for different quantities traits in maize hybrids. Indian J. of Agric. Res. 34(2): 107-111.

- Patch, L.H.; G.T. Bottger and B.A. App. (1938):** Comparative resistance to the European corn borer of two hybrid strains of field corn at Toledo, Ohio. J. Econ. Ent., 31: 337-340.
- ; **G.W. Still; B.A. App. and C.A. Crooks (1941):** Comparative injury by the European corn borer to open-pollinated and hybrid field corn. J. Agric. Res., 63: 355-368.
- Pereverzev, D.S.; A.S. Khromenko and I.P. Chuchmii, (1991):** Damage to maize by European corn borer in countries of the Danube basin. Vestnik. Sel Skokhozyaistvennoi Nauki Moskva. No. 2: 78-82.
- Poos, F.W. (1927):** Biology of the European corn borer (*Pyrausta nubilalis* Hubn.) and two closely related species in North Ohio. J. Sci., xxvii, No. 2: 47-88 (c.a. El-Sherif, S.E.I., Ph.D. Thesis, Fac. Agric. Alexandria Univ., 1965).
- Prury, R.A. and E.A. Wellington (1980):** Carleton's histological technique. 5th Ed. Oxford Univ. Press, New York, and Toronto.
- Ragab, M.E. (1988):** Insects attacking corn ears in the field and their natural enemies. Ph.D. Thesis, Fac. of Agric., Mansoura Univ.
- Raghip, M.M. (1979):** Studies on nitrogen fertilization and plant population densities on yield and its components in corn. M.Sc. Thesis, Fac. Agric., Al-Azhar Univ.

- Rahman, S.; Inayatullah; M. Shahid and F. Gul (2000):** Relative resistance of some maize cultivars to *Sitophilus oryzae* (L.) in store. Sarhad J. Agric. 16(1): 77-79.
- Rushdi, M.M.; A.A. Abdallah; Galal and W.H. Awadallah (1982):** Genetic variability inter and intra populations of maize and expected gain from selection for resistance to European corn borer (*Ostrinia nubilalis* (Hubner)). Ann.of Agric. Sci. Moshtohor 17: 3-12.
- Russell, W.A.; W.D. Guthrie and R. Grindeland (1974):** Breeding for resistance in maize to first and second brood of the European corn borer. Crop. Sci., 14: 725-727.
- Saleh, M.M.E. (1986):** Ecological studies on certain corn borers in Egypt. M.Sc. Thesis, Fac. of Agric., Cairo Univ.
- Salem, M.S.; A. Roshdy and F.I. Gaballa (1983):** Grain yield of maize in relation to variety, plant population and nitrogen application. Ann. Agric. Sci. Moshtohor, 20: 91-105.
- Salwau, M.I. (1985):** Effect of some agricultural treatment on the yield and technological properties of maize varieties. Ph.D. Thesis, Fac. Agric. Moshtohor Zagazig Univ.
- Sapunaru, T.; Tr. Mihail and C. Brezuleann, (1998):** Ursache, C.; Lisnic, Tr. New data on the behaviour of some maize lines and hybrids in relation to attack by the European corn borer *Ostrinia nubilalis* Hb. At S.C.A.

Podu-Iioaiei. Probleme de protectia plantelor 26 (2):
221-227.

- Sayers, A.C.; R.H. Johnson; D.J. Arnt and M.K. Bergman (1994): Development of economic injury levels for European corn borer (Lepidoptera: Pyralidae) on corn grown for seed. J. Econ. Ent., 87 (2): 458-465.
- Shahoudah, S.K. (1964): Studies on the pink borer of sugarcane, *Sesamia cretica* Led. and its control by chemicals. M.Sc. Thesis, Fac. of Agric., Cairo Univ.
- Shalaby, G.A.M. (1996): Studies on certain maize pests in Kafr El-Sheikh governorate. M.Sc. Thesis, Fac. of Agric., Zagazig Univ., Egypt.
- Sherif, M.A. and A.F. Lutfallah (1991): Assessment of losses in corn yield due to late natural infestation by *Ostrinia nubilalis* (Hubner) in Egypt (Lepidoptera: Pyralidae) Bull. Soc. Ent. Egypt 70: 25-30
- Showers, W.B.; J.F. Witkowski; C.E. Mason; F.L. Poston; S.M. Welch; A.J. Keaster; W.D. Guthrie and H.C. Chiang (1983): Management of the European corn borer. North central Reg. Publ. No. 22: 48-59.
- Snedecor, G.W. and Cochran, W.G. (1967): Statistical methods, six edition. Iowa State Univ. Press, Ames. Iowa, U.S.A.
- Steel, R.G.D. and Torrie, J.H. (1980): Principle and procedures of statistics, a biometrical approach. Mc Graw-Hill Book Company, Second Edit.

- Storey, C.L.; D.B. Sauer and D. Walker (1983): Insect populations in wheat, corn and oats stored on the farm. J. Econ. Ent. 76 (6): 1323-1330.
- Tantawi, A.M. (1989): Resistance of certain maize cultivars to the pink borer, *Sesamia cretica* Led. infestation. Proc. 1st Int. Conf. Econ. Ent., Cairo, Egypt, I : 271-275.
- Thome, C.R.; M.E. Smith and J.A. Mihm (1994): Yield reduction in a maize diallel under infestation with southwestern corn borer. Crop Sci. 34 (6): 1413-1435.
- Tigar, B.J.; P.E. Osbourne; G.E. Key; S.M.E. Flores and A.M. Vazquez (1994): Insect pests associated with rural maize stores in Mexico with particular reference to *prosthephanus truncatus* (Coleoptera: Bostrichidae). J. Stor. Prod. Res. 30 (4): 267-281.
- Umeozor, D.C.; J.W. Vanduyne; G.G. Kennedy and J.R. Bradley (1985): European corn borer (Lepidoptera: Pyralidae) damage to maize in eastern North Carolina. J. Econ. Ent., 78: 1488-1494.
- ; J.W. Van Duyn and J.R. Bradely (1987): Distribution of early instars of second-generation European corn borer on the maize plant. J. Agric. Ent. 4 (1): 48-54.
- Viereck, A. (1983): Influence of tissue hardness in the resistance of maize genotypes against *Ostrinia nubilalis* (Hbn) Z.P. flam zenzuchtung 90: 75-84.

- Willcocks, F.C. (1925):** The insects and related pests of Egypt. (c.f. El-Sherif, S.E.I., Ph.D. Thesis, Fac. Agric., Alexandria Univ., 1965).
- Yousef, M.M. (1968):** Effect of planting date and population density on growth and yield of maize. M.Sc. Thesis, Fac. Agric., Ain Shams Univ.
- Zangheri, S. and Masuti (1967):** Effect of treatment against *Ostrinia nubilalis* on the yield of two hybrid maize varieties. Giornate sitopatologiche Bologna, 345-552.