




REFERENCES

REFERENCES

- Abdalla, E.F. and Sammour, E.A. (1991):** Comparative study on the efficacy of neem seed extract and Diflubenzuron against the Egyptian cotton leaf worm *Spodoptera littoralis*. Bull. Of Entomological Society of Egypt Econ. Series No. 19, 207-214.
- Abdul-Kadir, A. and Connolly, J.D. (1991):** Preparation of 3-deacetyl-azaderachin using azadirachin isolated from *Azadirachta indica* seed. J. of Tropical Forest Sci., 3 (4): 342-347. {c.f. Hort. Abstr., 63 (11):8678}.
- Abo-Lila-MMS and Shoreit-MN (1997):** Trials for controlling varroa mite, *Varroa jacobsoni* Oud treated with formic acid in colonies of the honey bee in Egypt. Assiut-Journal-of-Agricultural-Sciences, 28: 2, 189-199.
- Abbott, W.S. (1925):** Methods for computing the effectiveness of insecticides. J. Econ. Entomol., 18(2): 265-273.
- Abdel-Zaher, R.T. (1998):** Using some plant extracts and their mixtures with pesticides to control cotton pests. M.Sc. Thesis, Fac. Agric., Moshtohor, Zagazig Univ.
- Abrol and Kakroo (1996):** Studies on seasonal activity and control of wax moths (*Galleria mellonella* L. and *Achoria grisella* F) attacking combs of four honeybee species. Mysore-Journal-of-Agricultural-Sciences. 1996, 30: 4, 365-373; 21 ref.
- Adler, C. (1992):** Vertical dispersion of adult *Sitophilus granarius* (L.) (Coleoptera: Curculionidae) in a wheat column flushed with modified atmospheres. J. Stored Prod. Res. 28: 201-209.

- Ahmad-R (1991):** Incidence of *Tropilaelaps clareae* on *Apis* species in Pakistan and its control with formic acid. Pakistan-Journal-of-Zoology. 1991, 23: 4, 363-364.
- Ahmed, M.E.R. (1985):** Biochemical studies on natural products from desert plants. Ph.D. Thesis, Fac. Agric., Cairo Univ.
- Ali, A.; Rasul, G.; Ulfat, M. and Ahmed, R. (1989):** Effect of fumigants on insect pests of stored grain. J. of Agric. Research Labor. 27(1): 77-81.
- Anuj-Bhatnagar; Agarwal-AP and Bhatnagar-A (1997):** Antifeedant activity of neem formulations against the larvae of castor semilooper, *Achaea janata* Linn. Plant-Protection-Bulletin-Faridabad, 49: 1-4, 17-18.
- Artemenko-LP; Kuz'-menko-VT and Kostritsa-PG (1988) :** Control of *Varroa jacobsoni* infestation among bees in the Crimea. Veterinariya,-Kiev, 63: 52-55.
- Barbattini-R; Greatti-M; D'-Agaro-M; Sabatini-AG; Colombo-R and Marcazzan-GL (1994):** Use of formic acid to control *Varroa jacobsoni*: verification of its efficacy and residues in honey. Ape-Nostra-Amica, 16: 4, 4-9.
- Bell, C. H. (1987):** The response of insects to fumigant concentration gradients. Proceedings of the fourth International Working Conference on Stored Product Protection, Tel Aviv, Israel, 21-26 September 1986 [edited by Donahaye, E. Navarro, s.]. 1987, 316-326; 13 ref. Bet Dagen, Israel; Agricultural Research Organization.

- Bracey-S and Fischer-F (1989):** Initial results of the field treatment of honey bee colonies infested with *Varroa jacobsoni* using formic acid in hot climates. American-Bee-Journal, 129: 11, 735-737.
- Buhlmann-G (1991):** Dose, concentration and method of application of formic acid in Dadant boxes. Schweizerische-Bienen-Zeitung., 114: 9, 505-511.
- Buscarlet, L.A. (1993):** Study on the influence of temperature on the mortality of *T. confusum* . du val. exposed to carbon dioxide or nitrogen. Zeitschrift fur Naturforschung. Section-C, Biosciences. 48: 7-8.
- Calderon, M. and Navarro, S.(1980):** Synergistic effect of CO₂ and O₂ mixtures on two stored grain insect pests. Developments in Agric. Engineering I: controlled atmosphere J. PP. 79-84, Italy.
- Calderone-NW and Nasr-ME (1999):** Evaluation of a formic acid formulation for the fall control of *Varroa jacobsoni* (Acari: Varroidae) in colonies of the honey bee *Apis mellifera* (Hymenoptera: Apidae) in a temperate climate. Journal-of-Economic-Entomology, 92: 3, 526-533.
- Calderone-NW and Spivak-M (1995):** Plant extracts for control of the parasitic mite *Varroa jacobsoni* (Acari: Varroidae) in colonies of the western honey bee (Hymenoptera: Apidae). Journal-of-Economic-Entomology, 88: 5, 1211-1215.
- Calderone-NW (1999):** Evaluation of formic acid and a thymol-based blend of natural products for the fall control of *Varroa jacobsoni* (Acari: Varroidae) in colonies of

Apis mellifera (Hymenoptera: Apidae). Journal-of-Economic-Entomology, 92: 2, 253-260.

Chakraborti-S and Chatterjee-ML (1996) : Effects of neem derivatives on survival, growth and development of the rice moth, *Coreyra cephalonica* (Stainton). Pest-Management-and-Economic Zoology, 4: 1-2, 37-43.

Champagne, D.; Isman, M.B. and Neil, G.H. (1989): Insecticidal activity of phytochemical and extracts of Meliaceae. Pp. 95-109, In Arnason, J.T.; Philogene, B.Jr. and Morand, P.(Eds.), Insecticides of plant origin, American Chemical Society Washington D.C.

Chiesa-F (1991): Effective control of varroaosis using powdered thymol. Apidologie, 22: 2, 135-145.

Childs, D.P. and and Overby, J.E. (1983): Mortality of the cigarette beetle in high-carbon dioxide atmosphere. J. Econ. Ent. 76:544-546.

Clark-A (1994): Control of varroa mites in British Columbia with either formic acid or Apistan. American-Bee-Journal, 134: 12, 829.

Contessi-A (1996) : New method of preserving combs with carbon dioxide. Rivista-di-Apicoltura, 4: 9, 16-19.

Cox-RL; Moffett-JO and Ellis-M (1987): Honey production, colony size and mite infestation levels of honeybee colonies treated with menthol. American-Bee-Journal, 127: 12, 842.

Das-SK and Isahaque-NMM (1999) : Control of red pumpkin beetle, *Aulacophora foveicollis* (Lucas). Journal-of-the-

Agricultural-Science-Society-of-North-East-India, 12: 1, 112-117.

Deshpande, S.G.; Sinha, B.; Nanda, B. and Sharma, R.N. (1993): Antifeedant activity in the total extract and fractions of the plant *cassine glauca* (Leguminaceae){Celastraceae} against the castor semi-looper *Achoea janata* (Noctuidae: Lepidoptera). Rajahmundry, India Society of Tobacco Science, 381-384. {c.f. Hort. Abstr., 63(11):8682}.

Desmarchelier, J.M. and Wohlgemuth, R. (1984): Response of several species of insects to mixtures of phosphine and carbon dioxide. Development in Agric Engineering. Controlled atmosphere and fumigation in grain storage, edited by Rippetal, B.E. pp. 75-81.

Desmarchelier, J.M. (1984): Effect of carbon dioxide on the efficacy of phosphine against different stored product insects. Mitteilung aus der Biologischen Bundesanstalt für Land-und Forstwirtschaft- Berlin- Dahlem Heft 220 Juli, 1984.

Dimetry, N.Z. and Abdalla, E.F. (1988): Feeding deterrent effects of various extracts of the bead tree, *Abrus precatorius* seed towards *Spodoptera littoralis*. Bull. Of the Entomological Society of Egypt. Economic Series No. 17, 85-92.

Dominguez-DA; Band-JM (1992): Cuban research in beekeeping technology: conservation of combs by chemical methods. Actualidad-Apicola. No. 61, 15-17.

- Duff-SR and Furgala-B (1992)** : Some effects of menthol and fluvalinate on mite-free honey bee (*Apis mellifera* L.) colonies. American-Bee-Journal., 132: 7, 476-477.
- Edward G. Jay, George E. Cantwell, Gordon C. Pearman, JR. and J.V. Thompson (1972)**: Control of the Greater Wax Moth, *Galleria mellonella*(L.), in comb honey with carbon dioxide, Part II, Field studies. American Bee Journal 9,342-344.
- Eguaras-M; Quiroga-S and Garcia-O (1996)**: The control of *Varroa jacobsoni* (Acari: Gamasida) by means of organic acids. Apiacta, 31: 2, 51-54.
- Eischen-FA (1998)**: Trials (and tribulations) with formic acid for varroa control. American-Bee-Journal, 138: 10, 734-735.
- EL-Deeb, H.A.H. (1981)**: Studies on some factors affecting the efficiency of atmospheric fumigation with methyl bromide and phosphine against the rice weevil, *Sitophilus oryzae* (L.) and the granary weevil, *S. granarius* (L.). Ph. D. Thesis, Fac. Of Agric., Cairo Univ., Egypt.
- EL-Lakwah, F.A.; Reichmuth, C.; Khattab, M.M. and Darwish, A.A.(1989 b)**: Sorption of phosphine by larvae and pupae of *Sitotorga cerealella* (Olivier); Lepidoptera. Egypt. J. Appl. Sci., 4(2): 1-21.
- EL-Lakwah, F.A.; Wohlgemuth, R. and Khattab, M.M. (1989a)**: Efficiency of phosphine and combination of phosphine with carbon dioxide against Khapra beetle larvae *Trogoderma granarium* (Everts). Anz. Schadlingsked-Pflakzenschuts, Umweltschutz. 62: 85-88.

- EL-Lakwah, F.A.; Abd EL-Gawad, A.; Meuser, F.; Wohlgemuth, R. and Darwish, A.A. (1991c):** Efficiency of phosphine alone and in mixtures with carbon dioxide against the adults of *Tribolium castaneum* and *Sitophilus oryzae*. Egypt. J. Appl. Sci., 4(2): 527-545.
- EL-Lakwah, F.A.; Meuser, F.; Abd EL-Gawad, A; Wohlgemuth, R. and Darwish, A.A.(1991d):** Efficiency of phosphine alone and in mixtures with carbon dioxide against Angoumois Grain Moth *Sitotroga cerealella* (Olivier), (Gelechiidae, Lepidoptera). J. of plant diseases and protection 98 (1): 92-102.
- Ellis-MD and Simonds-C (1987) :** Menthol fumigation of caged honey bees to prevent the spread of tracheal mites in package and queen bee shipments. American-Bee-Journal, 127: 12, 844.
- El-Meniawi-FA; Hashem-M; El-Mesieri-SM and Rawash-IA (1999):** Physiological effects of neem seed extract on the reproductive activities of the cotton leafworm, *Spodoptera littoralis* Boisd. Alexandria-Journal-of-Agricultural-Research, 44: 1, 79-96.
- EL-Nomrossy, A.M.Z.(1988):** Screening of naturally occurring toxicants in plant sources. Ph. D. Thesis, Fac. Of Agric., Cairo Univ.
- EL-Sayed, E.I. (1985a):** Evaluation of the insecticidal properties of the common Indian neem (*Azadirachta indica* A. Juss) seeds against the Egyptian cotton leaf

worm *Spodoptera littoralis*. (Biosd) Bull. Ent. Soc. Egypt. Econ. Ser., 13:39-47.

EL-Sayed, E.I. (1985b): Evaluation of the insecticidal properties of the common Indian neem (*Azadirachta indica* A. Juss) seeds against the Egyptian cotton leaf worm *Spodoptera littoralis*. (Biosd.) Bull. Ent. Soc. Egypt. Econ. Ser., 13:39-47.

Farag, A.A.A.A.(1984): Toxicological and chemical studies on plant extracts against *Tribolium castaneum* (Hbast) and *Spodoptera granarium* (L.). Ph. D. Thesis, Fac. Agric., Cairo Univ.

Ferguson-JL (1986): Menthol crystals reduce tracheal mite populations. American-Bee-Journal, 126: 11, 760.

Frilli-F; Milani-N; Barbattini-R; Greatti-M; Chiesa-F; Iob-M; D'-Agaro-M; Prota-R (ed.) and Floris-I (1992): The effectiveness of various acaricides in the control of *Varroa jacobsoni* and their tolerance by honeybees. Proceedings of 'The current state and development of research in apiculture', 25-26 October 1991, Aula Magna Facolta di Agraria, Sassari, Italy, 59-77.

Garza-Q-C and Dustmann-JH (1991): Formic acid against the tracheal mite [*Acarapis woodi*]. Deutsches-Imker-Journal, 2: 11, 455-456.

George E. Cantwell; E.G. Jay; G.C. Pearman, JR. and V. Thmpson (1972): Control of the Greater Wax Moth, *Galleria mellonella* (L.), in comb honey with carbon dioxide, Part I, Laboratory studies. American Bee Journal 8,302-303.

- Gerard-PJ and Ruf-LD (1995):** Effect of a neem (*Azadirachta indica* A. Juss, Meliaceae) extract on survival and feeding of larvae of four keratinophagous insects. *Journal-of-Stored-Products-Research*, 31: 2, 111-116.
- Goodman-RD; Williams-p; Oldroyd-BP; Hoffman-J (1990):** Studies on the use of phosphine gas for the control of greater wax moth (*Galleria mellonella*) in stored honey bee comb. *American-Bee-Journal*. 130: 7, 473- 477.
- Greatti-M; Barbattini-R and D'-Agaro-M (1993):** Treatment of varroosis. Formic acid as a control method against *Varroa jacobsoni*. *Obiettivi-e-Documenti-Veterinari*, 14: 6, 37-43.
- Greatti-M; Iob-M; Barbattini-R and D'-Agaro-M (1992):** Effectiveness of spring treatments with lactic acid and formic acid against *Varroa jacobsoni*. *Apicoltore-Moderno*, 83: 2, 49-58.
- Gretti M. and D-Agrao-O (1992):** Experiments on the control of *Galleria mellonella* and *Achoroia grisella* using carbon dioxide. *Apicoltore-Moderno*, 83:4, 123-128.
- Guirguis, M.W. and Gohar, K.M. (1989):** Toxicity and antifeedant activity of wild plant extracts against some insect pests. *Agric. Res. Rev.*, 67 (1): 1-12.
- Hashem, H.O.; Kheirallah, A.M.; Swidan, M.H.; and Osman, W.F. (1994):** Effect of alcoholic fruit extract of *Melia azadirachta* on the ovaries, fecundity and fertility of cotton leaf worm Moths *Spodoptera littoralis*. *Alex. J. Agric. Res.*, 39 (1): 211-227.

- Herbert-EW Jr.; Shimanuki-H and Matthenius-JC Jr. (1987):** The effect of two candidate compounds on *Acarapis woodi* in New Jersey. American-Bee-Journal, 127: 11, 776-778.
- Hoppe-H; Ritter-W and Stephen-EWC (1989):** The control of parasitic bee mites: *Varroa jacobsoni*, *Acarapis woodi* and *Tropilaelaps clareae* with formic acid. American-Bee-Journal, 129: 11, 739-742.
- Ibrahim, S.M., Ibrahim, A.A. and Fayad, y.H. (1984):** Studies on mass rearing of the wax moth, *Galleria mellonella* L. and its parasite (*Apanteles galleriae* W.) with some biological notes on the parasite. Agric. Res. Rev. 62(1): 449-454
- Imdorf-A; Bogdanov-S; Kilchenmann-V and Maquelin-C (1994):** Apilife-VAR - a varroa treatment substance with thymol as the active ingredient. Schweizerische-Bienen-Zeitung, 117: 6, 326-333.
- Imdorf-A; Bogdanov-S; Kilchenmann-V and Maquelin-C (1995):** Apilife VAR: a new varroacide with thymol as the main ingredient. Bee-World, 76: 2, 77-83.
- Imdorf-A; Kilchenmann-V; Bogdanov-S; Bachofen-B and Beretta-C (1995):** Toxic effects of thymol, camphor, menthol and eucalyptol on *Varroa jacobsoni* Oud. and *Apis mellifera* L. in a laboratory test. Apidologie, 26: 1, 27-31.
- Imti-B; Zudir-T (1997):** Effect of neem; *Melia azedarch* (Linn) and *Azadirachta indica* (A. Juss) on the incidence of *Sitophilus oryzae* (Linn) (Coleoptera; Curculionidae)

- on stored paddy. Plant-Protection-Bulletin-Faridabad, 49: 1-4, 44-47.
- Jones, R.N. (1983):** Toxicity of fumigant-carbon dioxide mixtures to the red flour beetle. J. Econ. Ent. 31:298-309.
- Kashi, K.P. and Bond, E.J. (1975):** The toxic action of phosphine: Role of CO₂ on the toxicity of phosphine to *S.granarius* and *T.confusum*. J.Stored prod. Res:11:9-15.
- Khadr, G.AA; EL-Monem, E.M.A. and Taha, M.A. (1986):** Effect of *Melia azzaderach* extract on *Spodoptera littoralis* (Boised.) larvae in the laboratory. Bull. Of the Ent. Soc. Of Egypt Economic Series No. 15, 235-243.
- Klinar-Z (1985):** Honeybee losses caused by volatile preparations used against Varroa. Veterinarski-Arhiv, 55: Supplement, S50-S51.
- Kshirsagar-KK; Salvi-SR; Mahindre-DB and Chauhan-RM (1983):** Menthol as an effective acaricidal fumigant. Proceedings of the Second International Conference on Apiculture in Tropical Climates, New Delhi, February 19-March 4, 1980, 454-456.
- Kumar, B.H. and Thakur, S.S.(1988):** Certain non-edible seed oils as feeding deterrents against *Spodoptera littoralis*. J. of The Oil Technologists Association of India, 20(3): 63-65.
- Lee-ML; Choi-MY; Choi-KS. (1991):** Effects of controlling *Galleria mellonella* in honey bee combs with aluminium phosphide, its residues and safety to honey bees. Korean-Journal-of-Apiculture. 6:1, 12-15.

- Lima-CPF-de; De-Lima-CPF; Ripp-BE (ed.) (1984):** Minimal fumigant requirements for long-term air-tight storage of grain. Controlled atmosphere fumigation in grain storage. Proceedings of an international symposium 'Practical aspects of controlled atmosphere and fumigation in storages ' held from 11 to 22 April 1983 in perth, Western Australia. 1984, 665-671.
- Liu-TP (1995) :** A possible control of chalkbrood and nosema diseases of the honey bee with neem. Canadian-Beekeeping, 18: 5, 107-109.
- Locatelli, D.P. and Daolio, E.(1991):** Effectiveness of CO₂ in a vacuum autoclave at different temperatures... on *Rhizopertha dominica* (F.) and *Sitophilus oryzae* (L.). Tecnica-Molitoria. 42(7): 605-610.
- Lodesani-M; Bergomi-S; Pellacani-A; Carpana-E and Rabitti-T (1990):** A comparative study on the efficacy of some products for controlling Varroa, and determinations of their residues. Apicoltura, No. 6, 105-130.
- Lupo-A and Gerling-D (1990):** comparison between the efficiency of summer treatments using formic acid and Taktic against *Varroa jacobsoni* in beehives. Apidologie., 21: 3, 261-267.
- Manehasa-M (1987):** Treatment of varroa disease with a combination of tobacco smoke, acetic acid and Milbol. Buletini-i-Shkencave-Zooteknike-e-Veterinare, 5: 2, 55-61.
- Mann-DD; Jayas-DS; Muir-WE (1997):** Conducting a successful CO₂ fumigation in welded-steel hopper bin.

ASAE Annual International Meeting, Minneapolis, Minnesota, USA, 10-14 August, 1997. Paper -American-Society-of-Agricultural-Engineers, No. 976064, 9 pp.

Mladan-V; Erski-Biljic-M; Jakic-D and Simic-M (1985) :

The application of some preparations for the eradication of varroa disease. Veterinarski-Glasnik, 39: 4-5, 489-496.

Moffett-JO; Cox-RL; Ellis-M and Rivera-R (1987) :

Effect of winter treatments with menthol on populations of tracheal mites, *Acarapis woodi*, of honey bees in Tamaulipas, Mexico and Nebraska. American-Bee-Journal, 127: 12, 848-849.

Morner, J. ; Hyvonen, A. and Simonsson, M.,(1987):

Experiments with carbon dioxide against stored products pests. Vaxtskyddsnotiser. 51(3): 87-92.

Morse(1975): Honeybee pests, predators and diseases. pp. 106-123.

Murugan-K; Jeyabalan-D; Kumar-NS; Babu-R;

Sivaramakrishnan-S and Nathan-SS (1998):

Antifeedant and growth-inhibitory properties of neem limonoids against the cotton bollworm, *Helicoverpa armigera* (Hubner). Insect-Science-and-its-Application, 18: 2, 157-162.

Mutinelli-F; Cremasco-S; Irsara-A; Baggio-A; Nanetti-A

and Massi-S (1996): Organic acids and Api Life VAR in the control of varroosis in Italy. Apicoltore-Moderno, 87: 3, 99-104.

- Navarro-S; Jay-EG (1987):** Application of modified atmospheres for controlling stored grain insects. Monograph,-British-Crop-Protection-Council., No. 37, 229-236; In Stored Products Pest Control, Proceedings of a Symposium held at the University of Reading, 25-27 March, 1987.
- Nelson-DL (1995) :** Control of the honey bee tracheal mite with menthol and formic acid. Canadian-Beekeeping, 18: 6, 136-137.
- Osman, M.Z. (1993):** Effects of neem seed extract on growth and development of larvae of *Pieris brassicae*. J. of Appl. Ento., 115 (3):254-258. {c.f. Rev. of Agric. Ento., 82 (3):2220.
- Osman, M.Z. and Bradley, J. (1993):** Effects of neem seed extracts on pholeastor (*Apanteles*) glumeratus (*Braconidae*), aparasitoid of *Pieris brassica* (Lep.: *Pieridae*). J. of Appl. Ento., 115 (3): 259-265. {c.f. Rev. of Agric. Ento., 82 (3):2221}.
- Pandian-RS and Devi-TS (1998) :** Repellent action of plant oils on mosquito. Insect-Environment, 4: 2, 58.
- Pathak-PH and Krishna-SS (1991):** Postembryonic development and reproduction in *Corcyra cephalonica* (Stainton) (Lepidoptera: *Pyralidae*) on exposure to eucalyptus and neem oil volatiles. Journal-of-Chemical-Ecology, 17: 12, 2553-2558.
- Price, N.R. (1985):** Review, the mode action of fumigants. J. Stored Prod. Res. 21:157-164.

- Prokopy-RJ and Powers-PJ (1995):** Influence of neem seed extract on oviposition and mortality of *Conotrachelus nenuphar* (Col., Curculionidae) and *Rhagoletis pomonella* (Dip., Tephritidae) adults. Journal-of-Applied-Entomology, 119: 1, 63-65.
- Rademacher-E; Polaczek-B and Schricker-B (1995):** Formic acid: a new method of application of the product to hives. Vida-Apicola, No. 70, 17-20.
- Raffa, K.F. (1987):** Influence of host plant on deterrence by azadirachtin of feeding by fall armyworm larvae (Lepidoptera: Noctuidae). J. of Ent., 80 (2): 384-387.
- Rashad, Amira M., Suzan, H.; Taher, Sonia M. Naguib and Abd EL-Salam, N.M. (1991):** Effect of *Melia azadirachta* extract on *Earias insulana* and *Pectinophora gossypiella*. Egypt. J. Biol. P. Cont., 1 (1):117-120.
- Rickli-M; Imdorf-A and Kilchenmann-V (1991):** Treatment of varroa disease using compounds of essential oils. Apidologie, 22: 4, 417-421.
- Saleh, M.A.; EL-Bolok, M.M.; Abdel-Salam, K.A. and Ibrahim, N.A.(1986a):** Plant extracts affecting insect feeding, growth and metamorphosis. Bull. Of Fac. Agric., Cairo Univ., 37(1): 529-539.
- Schuster-H (1997):** Comparison of different methods for varroa treatment with formic acid. Imkerfreund, 52: 7, 4-12; Bj.
- Senguttuvan-T; Kareem-AA and Rajendran-R (1995):** Effects of plant products and edible oils against rice moth

Corcyra cephalonica Stainton in stored groundnuts. Journal-of-Stored-Products-Research, 31: 3, 207-210.

Sharma-SD; Desh-Raj and Raj-D (1999): Evaluation of different methods of using formic acid against *Acarapis woodi* Rennie infesting *Apis mellifera* L. Agricultural-Science-Digest-Karnal, 19: 3, 167-171.

Sharma-SD; Kashyap-NP; Desh-Raj and Sharma-OP (1994): Control of ectoparasitic mite, *Tropilaelaps clareae* Delfinado and Baker infestation with formic acid and sulphur. International-Journal-of-Tropical-Agriculture, 12: 1-2, 96-100.

Smirnov-AM; Kolchko-RT and Luganskii-SN (1994): Technology for using formic acid against Varroa infection. Problemy-Veterinarnoi-Sanitarni-i-Ekologii, 93: 1, 108-109.

Srivastava-RK and Krishna-SS (1990): Temperature- and exposure duration-related eucalyptus oil odour effects on egg hatchability and subsequent postembryonic development in *Dysdercus koenigii* (F.) (Heteroptera: Pyrrhocoridae). Journal-of-Advanced-Zoology, 11: 2, 103-105.

Srivastava-RK and Krishna-SS (1992): Effect of exposure of eggs or first instar nymphs of *Dysdercus koenigii* (F.) (Heteroptera: Pyrrhocoridae) to eucalyptus oil odour on the insects postembryonic development and/or reproductive potential. Phytophaga-Madras, 4: 1, 47-52.

Srivastava-RK; Gurusubramanian-G and Krishna-SS (1995): Postembryonic development and reproduction in

- Dysdercus koenigii* (F.) (Heteroptera: Pyrrhocoridae) on exposure to eucalyptus oil volatiles. Biological-Agriculture-and-Horticulture, 12: 1, 81-88.
- Srivastava-SK and Krishna-SS (1987):** Eucalyptus oil odor experienced by fifth instar nymphs of *Dysdercus koenigii* (Fabr.) (Heteroptera: Pyrrhocoridae) and its subsequent effect on the insect's reproduction. Zeitschrift-fur-Angewandte-Zoologie, 74: 4, 467-470.
- Su-TianYun; Mulla-MS and Su-TY (1999):** Oviposition bioassay responses of *Culex tarsalis* and *Culex quinquefasciatus* to neem products containing azadirachtin. Entomologia-Experimentalis-et-Applicata, 91: 2, 337-345.
- Thyagaraj-NE (1999):** Effect of neem based insecticides against the mosquito (*Culex quinquefasciatus* Say) larvae. Insect-Environment, 5: 2, 78-79.
- Tremblay, A.(1978):** Controlling wax moths in Honeycombs. Extension service, Oregon state Univ., Corvallis, pp. 240-241.
- Valladares-GR; Ferreyra-D; Defago-MT; Carpinella-MC and Palacios-S (1999):** Effects of *Melia azedarach* on *Triatoma infestans*. Fitoterapia, 70: 4, 421-424.
- Veen-J-van; Calderon-Fallas-RA; Cubero-Murillo-A and Arce-Arce-HG (1998):** *Varroa jacobsoni* in Costa Rica: detection, spread and treatment with formic acid. Bee-World, 79: 1, 5-10.

- Vincent, L.E. and Lindgren, D.L. (1975):** Toxicity of methylbromide at various temperatures and exposure period to the metamorphic stages of *Trogoderma variable*. J. Econ. Ent., 68 (1):53-56.
- Wang-ZhaoLong; Gao-HongMing; Zhang-BiAo; Wu-XiaoXia; Wang-ZL; Gao-HM; Zhang-BA and Wu-XX (1999):** Exploitation of insecticides from medicinal plant resource - investigation and screening of botanical insecticides. Research-progress-in-plant-protection-and-plant-nutrition, 203-208.
- White, N.D.G. and Jayas, D.S. (1991):** Control of insects and mites with carbon dioxide in wheat stored at cool temperatures in non air tight bins. J. Econ. Ent. 84: 1933-1942.
- Winks, R.G. (1984):** The toxicity of phosphine to adults of *T. castaneum*. Time as a dosage. J. Stored Prod. Res. 20: 45-56.
- Xie-YS; Fields-PG and Isman-MB (1995):** Repellency and toxicity of azadirachtin and neem concentrates to three stored-product beetles. Journal-of-Economic-Entomology, 88: 4, 1024-1031.
- Zimmerman-RJ; Sclar-DC; Randolph-T and Cranshaw-WS (1995):** Toxicity of neem-derived insecticides to various life stages of the elm leaf beetle. Journal-of-Arboriculture, 21: 4, 181-186.