

REFERENCES

VI. REFERENCES

Abbot , W. S. (1925) : A method of compating the effectiveness of insecticide
.J.Econ. Entomol. 18: 265-267.

Abd el-Megeed- M.I.; M . G Abbas ; S . M . EL-Sayes ;
and Moharam E. A .(1998) : Efficacy of certain
biocides against potato tuber moth, *Phthorimaea
operculella* under field and stora conditions . Annals.
Agric. Scie . 1: 309 - 317.

Abd-El Samie , M.B ; H.M. Salem and M. Mohle (1979)
: Identification and distribution of esterase and
phosphatase isozymes in various tissues of
susceptible and resistance to insecticides of
cotton leafworm , *Spodoptera littoralis* (Boisd.).
Proc. 3 rd. Arab .pesticide .conf ,Tant Univ., 450.

Abd el –Wahab – M.A.; M. M. Soliman; F. A. Abd El-
Galill; A. M. El- Ghareeb and M. A. Abd El-
Wahab(1987): Toxicity of certain insecticides to
potato tuber worm, *Phthorimaea operculella*
(Zeller) .Assiut J. Agric. Scie., 18(3) 247-254.

- Abedi, Z. H. and A. W.A. Brown (1960)** : Development and reversion of DDT- resistance in *Aedes aegypti*. Can, J. Genet. Cytol., 2: 252-261.
- Abo- El ghar, M. R.; A. M. shaaban ; M. M. Abd El-Hafez and M.A. El.Molla (1984)** : Cholinesterase , aliesterase and non- specific esterase activity in relation to insecticides resistance of *Spodoptera littoralis* (Boisd.). Z. pflanzenkr pflanzenschutz , 91: 476-482.
- Aldosari, S. A.; T. F. Watson ; S. Sivasubramanian; and A. A. Osman (1996)**: Susceptibility of field populations of beet armyworm (Lepidoptera : Noctuidae) to cyfluthrin, methomyl and profenofos and selection for resistance to cyfluthrin. J. Econ. Entomol. 89(6) :1359-1363.
- Ali, M. I. (1991)** :Efficacy of a granulosis virus on the control of potato tuber moth , *Phthorimaea operculella* (Zeller) (Gelechiidae : Lepidoptera) infesting potatoes in Bagladesh. Bandladesh. J. zodogy., 19: 141- 143.
- Allam, A. M. (1990)**: Effect of treatment with some growth regulators on activity of some metabolizing enzymes in resistant *Spodoptera*

littoralis (Biosd.) to insecticides . Ph. D. Thesis,
Fac. Agric. Al-Azhar Univ., Egypt.

**Allam, A. M.; E. I. Moarad; Y.F. Goneim; A. Mohanna
; F. A. Ayed and M. A. El- Guindy (1994):**
Relative susceptibility of Egyptian cotton
leafwoem , *Spodoptera littoralis* (Biosd.) larvae
to chlorpyrifos in relation to cross- resistance
pattern and esterase activity. J. product . Dev . 2:
169-183.

Andrew, R.; J. Herbas; J. Alcazar and V. Lion (1992) :
Integrated management of potato tuber moth in
cochabambas, Bolivia, CIP. Circular (CIP) . 19: 8-
9.

Anonymous (1981): Annual report of the international
potato center ; Lime, Peru .(in press) .1981,xii +
133pp.; many rig.; 35 ref.

Arx , R. Von and F. Gebhardt (1990): Effects of
agranulosis virus and *Bacillus thuringiensis* on
life – table parameters of the potato tuber moth ,
Phthorimaea operculella .Entomophaga 35: 151-
159.

**Bekhiet, H. K.M.; G. M. Moawad ; R. A. El-Bendewy
and A. M. Mabrouk (1997):** Control of the
potato tuber moth *P.operculella* (Zeller.) in

potato crop . Egypt. J. Agric. Res., 75(74) : 923-937.

Bozkurt, E. and H. Genc (1994) : Investigations on the host plant distribution , damage and biology of the potato tuber moth, *Phthorimaea operculella* (Zeller.) (Lepidoptera : Gelechiidae) . J.Agric. Forestry 18 (2) : 111-116.

Bradford, M. M. (1976) : A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein- dye binding . Anal .Biochem. (72) : 248- 254.

Briese, D. T. and H. A. Mende (1983) : Selection for increased resistance to a granulosis virus in the potato tuber moth , *Phthorimaea operculella* (Zeller) (Lepidoptera : Gelechiidae) . Bull. Entomol . Res. 73 (1) : 1-9 .

Broodryk, S. W.(1971) : Ecological investigations of the potato tuber moth, *Phthorimaea operculella* (Zeller.) (Lepidoptera : Gelechiidae) . Phytophactica. 3 : 73-84.

Brown, A. W. A. (1958): Insecticides resistance in arthropods Wid. Health Organ., Geneva : 178-185.

- Carlos . J. Perez , A. M. Shelton and R.C. Derksen**
(1995) : Effect of application of technology of *Bacillus thuringiensis* subspecies *Kurstaki* on management of diamondback moth.(Lepidoptera : Plutellidae) . J. Econ. Entomol. 88(5) : 1113-1119.
- Carlos . J. Perez , A. M. Shelton (1997)** : Resistance of *plutella* *xylostella* (Lepidoptera : Plutellidae) to *Bacillus thuringiensis* in control America . J. Entomol . 90(1) : 87 – 93.
- Chandramohan , N and K . Nanjan (1993)** : Protection of stored potato tubers moth . J . Agric.80(4): 239 – 240.
- Clark , A. G. ; J. N. Smith and T. W. Speir (1973)** : Cross specificity in some vertebrate and insect glutathione-transferases with methyl parathion dimethyl P. nitrophenyl n- acetylcyste amine as substrates . Biochem . J. , 135: 385.
- Clark , A. G. ; N.A. Shamaan ;M. D. Sinclair and W. C. Dauterman (1986)** : Insecticide metabolism by multiple fly,Musca domestica glutathione S-transferases in two strains of the house (L.) . Pestic. Biochem. Physiol.,25:169

- Clavijo , Sand P. A. Natz (1995) :** Ovicide effect of the carbamat insecticide thiodicarba for the control of Lepidopteran species in the families Gelehiidae and Notuidae Agr . Top . 45 (1) 27-34.
- Crow , J. F. (1957) :** Genetics of insect resistance to chemicals .Ann . Rev . Entomol . , 2: 227-246 .
- Dray ,O. ;G. P. Georghiou ; E. Parsons and N. Pasteur (1990) :** Microplate adaptation of Gomori's assay for quantitative determination of general esterase activity single insects .J. Econ . Entomol . 83 : 218-219 .
- Das , G . P . ; E . D . Magallona ; K . V . Raman and C .B .Adalla (1992) :** Effects of different components of IPM in the management of the potato tuber moth in storge . Agric Ecos&Envir . 41 : 321 – 325 .
- Dauterman , W . C . (1983) :** Role of hydrolyses and glutathione s-transferases in insecticide resistance . (C . F . Georghiou , G . P . and T . Saito (ed) , Pest Resistance to Pesticides pleneum , Publ . Crop . , New York , 229 –247 .

- Duncan , D . B . (1955)** : Multiple range and multiple F test
 .Biometrics
 11 : 1 .
- Ebora , R . V . ; M . M . Ebora and M . B . Sticklen .**
(1994) transgenic potato expressing the *Bacillus*
thuringiensis CryIA
 (c) gene effects on the survival and food
 consumption of *Phthorimaea operculella*
 (Lepidoptera : Gelechiidae) and *Ostrinia*
nubilalis (Lepidoptera : Noctuidae) . J . Econ .
 Entomol . 87 : 1122 – 1127 .
- EL –Guindy , M . A . ; G . N. El- Sayed and I . I. Guirgis**
(1975a) Comparative toxicities of certain organic
 insecticides in phosvel – resistant strain of the
 Egyptian cotton leaf worm . Bull . Ent . Soc .
 Egypt , Econ . Ser . , 9 : 137 – 146 .
- EL –Guindy , M . A . ; G . N. El- Sayed and I . I. Guirgis**
(1975b) Cross – resistance characteristic of
 Duursban – selected strain in the Egyptian cotton
 leaf worm , *Spodoptera littoralis* (Boisd.) . Bull .
 Ent . Soc . Egypt , Econ . Ser . , 9 : 157 – 182 .
- EL –Guindy , M . A . ; G . N. El- Sayed and I . I. Guirgis**
(1975c) :The effect of selection with cyolane on
 the Egyptian cotton leafworm , *Spodoptera*

littoralis (Biosd.) . Bull . Ent . Soc . Egypt,
Econ . Ser . , 9 : 183 – 189 .

**EL –Guindy , M . A . ; A . M . Elefai and M . M . Abddel
– Sattar (1983) :** The pattern of cross– resistance
to insecticides and juvenil hormone analogues in
diflubenzuron resistant strain of the cotton
leafworm , *Spodoptera littoralis* (Biosd.) .Pestic.
Sci., 14 : 235- 245 .

**EL –Guindy , M . A . ; A . M . Elefai ; A. M . El-
Samadesy and Y. F. Ghoneim .(1989a) :**
The impact of several selection procedures on
resistance to monocrotophos , diflubenzuron and
methopren in the cotton leafworm ,
Spodoptera littoralis (Biosd.) . Int . Pset . Cont .
32: 72 – 76

EL- Sheikh , A . E . S . ; G . S . A . EL- Ghar (1993) :
The influence of selected insecticide – fertilizer
combinations against the potato tuber moth ,
Phthorimaea operculella (Zeller.) on potato
plant. Bull . Entomol . Soc . Econ .Ser . Egypt .
20: 15-23 .

**EL- Tantawy , M . A . ; M. I. El- Nabarawy and A. A.
Sallam . (1993):** Selective toxicity of certain
insecticides against the different stages of the

- potato tuber worm , *Phthorimaea operculella* (Zeller.) (Lepidoptera : Gelechiidae) .Ann . Agric . Sc ., Moshtohor , 31(3): 1709-1714 .
- Escriche , B. ; A. C. Martinez ; M. D. Ramirez ; M. D. Reel ; F. J. Silva and J. Ferre . (1994) : Occurrence of three different binding sites for *Bacillus thuringiensis* delta – endotoxins in the midgut brush border membrane of the potato tuber moth , *Phthorimaea operculella* (Zeller.). Arch. Insect Biochem . Physiology , 26(4): 315 – 327 .
- FAO , (1995) : Potato in the 1990S. Situation and prospects of the World potato . Economy , Fao , Rome . P . 29 .
- Fenemore , P .G . (1977) : Oviposition of potato tuber moth , *Phthorimaea operculella* (Zeller.) (Lepidoptera : Gelechiidae) ; Fecundity in relation to mated state , age and pupal and weight . N .Z . J. Zool . 4:187 –191.
- Finney , D. J. (1971) :Probit analysis 3rd ed . Cambridge. Univ . Press, Cambridge , England . 333pp.
- Fuente- Martinez – J. M. de ; R. Farias-Rodriguez ; C. Aguilar-Zacrias ; R. Ramirez – Morales ; R. Rocha- rodriguez ;and

- J. J. Pena** – cabriales (1992): Evaluation of bioinsecticides for the control of the potato tuber moth , *Phthorimaea operculella* (Zeller.) (Lepidoptera : Gelechiidae) in potatoes in Leon , Guanajuats, Mexicana 86: 176-183 .
- Glass , E. H. (1986)** : Pesticide rsistance , strategies and tactics for management . Comitte on Strategies for the Management of pesticide resistance . Pest populations , National papers from aSymposium held in washington , Nov .27-29, 1984.NationalAcademy press , washington . D.C., 471pp.
- Georghiou , G. P. AND J. R . Calman (1969)** : Results of fentrothion selection of *Culex fatigans* Anophiles albimanus . Bull . Wld Hlth . Org . 40 : 97-101 .
- Georghiou , G. P. and R. B. Mellon (1983)** : Pesticide resistance in time and space .In : Pest Resistance to pesticides , edited by Georghiou , G. P. and T. Saito , Plenum Press , New York and London,1983 ,809 pp.
- Gomaa , E. A. ; M. A. E- tantaway and A. A. Kdider (1982)** : Result of fentrothion selection of *Culex fatigans* Anophiles albimanus Esterase activity

in different stage of the american bollworm,
Heliothis zea . proc . 2nd. Egyptian-Hungarian-
conf .plant Protection : 53-62 .

Grant, D. F. and F. Matsumura (1989) : Glutathione s-
transferase 1 and 2 susceptible and resistant
Aedes aegypti . Pestic . Biochem. Physiol. 33:
132- 143.

Grant, D. F.; D. M. Bender and B. D. Hammock (1989):
Quaantitative kinetic assay for glutathione s-
transferase and general esterase in individual
mosquitoes using EIA reader. Insect Biochem.,
19: 741- 751.

**Green , M. B. ; H. M. Lebaron and W. K. Moberg
(1990)** : Managing resistance to agrochemicals
from fundamental research to practical strategies .
Developed from a symposium sponsored by the
Division of Agrochemicals at the American
chemical. Society , Los Angeles , California, Sep.
25 – 30, 1988. American chemical. Society,
Washington, D. C. 1990. 496 pp.

**Gurigus, M. W. ; M. A. El-Tantawi; F. A. Ayad; M. S.
Abd El-Fattah and Y. Ghoneim (1985):** Studies
on the rates of development of resistance to five
insecticides in eggs of *Spodoptera littoralis*

(Boisd.) . Proc.6 th Arab Pesticide Conf. Tanta Univ. Vol.(1) : 11-20 .

- Haines ,C. P. (1997)** : The potato tuber moth , *Phthorimaea operculella* (Zeller.) abiblioraphy of recent literature and a review of its biology and control on potatoes in the field and store . Rep. Trop. Inst. G 112. 15 pp.
- Hama, H. (1984)** : Mechanism of fenitrothion and diazinon – resistance in the green rice leafhopper, *Nephotettix cinciticepe* (under): The role of aliesterase. Jep. J. Appl. Entomol. Zool.28: 143-149.
- Hayaoka, T. and W. C. Dauterman (1983)** : The effect of Phenobarbital induction on glutathione conjugation of diazinon in susceptible and resistant house flies . Pestic. Biochem. Physiol.; 19: 344.
- Hayder, M. F. and L. S. El-Sherif. (1987):** Microbial control of the potato tuber worm, *Phthorimaea operculella* in the field . Bull. Ent. Soc., Egypt. Econ. 16 : 127- 132.
- Hoskins, W. M. and H. T. Gordan (1956)** : Arthropod resistance to chemicals . Ann. Rev. Entomol. 1: 89-122.

- Ibrhim, S. A. and J. A. Ohhea. (1995)** :Biochemical and toxicological studies with laboratory and field populations of *H. virescens* (F.). Pestic. Biochem. Physio. 53 (2) : 116-126.
- Joseph. O. Lalah; C. I. Chien; N. Motoyama and W. C. Dauterman. (1995)** : glutathione s-transferases :
– Naaphthyl acetate activity and possible role in insecticide resistance. J. Econ. Entomol. 88(4) : 788- 790.
- Judy , M. C.; J. Chaufaux ; C. Buisson : N. Gilois : V. Sanghis and D. Lereclus (1996)** : *Spodoptera littoralis* (lepidoptera : Noctuidae) resistance to CryIc and cross-resistance to other *Bacillus thuringiensis* Crystal toxins . J. Econ. Entomol. 89 (4): 791-797.
- Kashyap, N. P.; D. C. Sharma; R. M. Bhagat and S. M. Suri. (1993)** : Bioefficacy of insecticides against potato tuber moth . J. Indian potato Association., 20 (2) : 179-181.
- Keiding, J. (1967)** : Persistance of resistance populations after the relaxation of the selection pressure. World Rev, Pest . Contr., 6: 115- 130.
- Khalid , M. S. (1997)** : Bichemical detection of resistance in the cotton leafworm *Spodoptera littoralis*

(Boisd.) . Ph. D. , Thesis , Fac. Agric. Al-Azhar Univ.

Kroschel, J. (1992): Integrated plant protection in potato culture in the Republic of jemen under special view of the Integrated Biological control of *Phthorimaea operculella* (Zeller) . Weikersheim (Germany). Argraf; 241 pp.

Lianderal- Cazares, C.; A. T. Lagunes: S. J. L. Carrillo; C. M. Sosa; J. G. Vera and H. M. Bravo. (1996): Susceptibility of *Phthorimaea operculella* (Zeller.) to insecticides. J. Entomol. Sci., 31 (4) : 420-426.

Loannou , Y. M. and W.C. Dauterman (1979) : In vitro metabolism of etrimfos by house flies. Pestic. Biochem. Physiol., 10: 31-59.

Madi, S. M. (1976): The insecticidal action and toxicology of certain synthetic organic insecticides on the cotton leafworm . Ph.D. Thesis, Fac.Agric., Ain Shams Univ., Egypt.

Mariy, F. M.; M. A. Daoud; G. B. El- Saadany and M. Y. Ibrahim (1999) : Biological studies on potato tuber moth, *Phthorimaea operculella* (Zeller) . Ann. Sci. Ain Shams Univ. , 44(1) : 363-378.

- Matsumure, F. (1975) :** Toxicology of insecticides.,
P.503, Plenum Press, New York, 1975.
- Melander, A. L. (1914) :** Toxicology of insecticides.,
P.503, Plenum Press, New York, 1975.
- Metcalf, R. L. (1955) :** Physiological basis for insect,
resistance to insecticides *physiol. Rev.*, 35: 179-
232
- Moawad , G. M.; R. A. El-Bedewy; S. Abd- El-Halim;
H. K. M. Bekheit; M. Mabrouk; A. Farghaly
and A. Legnaoui (1997) :** Large-scale
implementation of intergrated pest mangement in
Egypt . Interanational Potato Center (CIP) cirular,
23(1) : 8- 13.
- Moad, G. M.; H. K. Bekhiet ; A. M. Mabrouk; R. El-
Bedewy and A. Legnowi (1998 a) :** Potential uses
of neem and insect pathogens for control tuber
worm , *Phthorimaea operculella* (Zeller.) in
storge . *Ann. Agric. Sci. Cairo*. 1: 297-308.
- Moad, G. M.; S. M. Abd El-Halim : H. K. M. Bekhiet ;
A. M. Mabrouk; R. El- Bedewy and A. Lagnaui.
(1998 b) :** Effect of different IPM components in
the management of the potato tuber moth,
Phthorimaea operculella (Zeller.) in potato

cultivations. Ann. Agric. Sci., Sp. Issue., 3: 1055-1067.

Mohanna, A. H.; Allam and Y. F. Ghoneim (1996) :

Sensitivity of esterases in the development of resistance to methomyl on *Spodoptera littoralis* (Bosid.) Larvae and cross- resistance to several insecticides . J. Product. Dev. (in press) .

Motoyama, N. and W. C. Dauterman (1972) :

In vitro metabolism of azinphosmethyl in susceptible and resistant house flies. Pestic. Biochem. Physiol., 2: 113.

Motoyama, N. and W. C. Dauterman (1980) :

Glutathione s-transferases : Their role in the metabolism of organophosphorus insecticides. Res. Biochem. Toxicol., 2: 49- 69.

Nabi, M. N. (1985) :

Evaluation of cypermethrin and fenvalerate against potato tuber moth, *Phthorimaea operculella*. Ann. Appl. Biol. 6: 22-23.

Nasseh , O. M. and M. A. Al- Furassy (1992) :

Studies on control of potato tuber moth , *Phthorimaea operculella* (Zeller.) (Lepidoptera : Gelechiidae) by natural and chemical insecticides in the

- Repulic of Yemen. Anzeiger fur Schadlingskund ,
Pflanzenschutz, Umweltschutz., 65 (8): 157-159.
- Nieto, H. R.; C. L. Cazares and G. Z. Lara (1989):**
Selected attacking of *Phthorimaea operculella*
with sex pheromones in Peru. Agric. Ecosy. Enir.
21: 85-99 .
- Nomier, A. A.: A.Maharn; F. I. Moustafa and M. I.
Hashem (1982) :** Mixed function oxidase and
glutathione s-transferase activities and their
relationship insecticide resistance in Egyptian
cotton leafworm Accepted inG. Agric. Res. Alex.
- Osmelak, J. A. (1989) :** Comparison of carbaryl , maldison
and permethrin wettable powders as insecticidal
dusts to protect potato tubers for export against
potato moth, *Phthorimaea operculella* (Zeller.) .
Plant. Prot. Quarterly . 4 (2) : 84-85.
- Oppenoorth, F. J.; V. Rupes; S. Eibashir, N. W. H.
Houx and S. Voerman (1972) :** Glutathione –
dependent degradation of parathion and its
significance for resistance in the house fly. Pestic.
Biochem. Physiol., 2: 262.
- Oppenoorth, F. J.; H. R. Smissaert; W. Welling; L.J. T.
Van der pas and K. T. Hitman (1977) :**
Insensitive acetylcholinesterase, high glutathione

s-transferase, and hydrolytic activity as resistance factors in a tetrachlor- viphos-resistant strain of house fly . Pestic. Biochem. Physiol., 7: 34-47.

Oppenoorth, F. J., L. J. T. Van der pas and N. W. H.Houx (1979) : Glutathione s-transferease and hydrolytic activity in a tetra-chlorvinphos – resistant strain of house fly and their influence on resistance. Pestic. Biochem. Physiol., 11: 176 .

Paparatti, B. (1993) : Experiments in the integrated control of *Phthorimaea operculella* (Zeller) (Lepidoptera : Gelechiidae) in northern Latium. Frustula. Entomol. 16 : 9-22.

Patricia, C. Bolin ; W. D. Hutchison and D. A. Andow (1999) : Long-Term selection for resistance to *Bacillus thuringiensis* CryIAc end toxin in a Minnesota population of Europeam. Corn. Borer (Lepidoptera : Crambidae) . J.Econ. Entomol. 92 (5) : 1021- 1030.

Pimprikar, C. D. and G. P. Georghiou (1979) : Mechanism of resistance to diflubenzuron in house fly , *Musca domestical* (L.) . Pestic. Biochem. Physiol., 12: 10.

Raj, B. T. and T. P. Trived (1993) : Efficacy of insecticides against potato tuber moth,

- Phthorimaea operculella* infesting potatoes in Maharashtra. J. Indian Potato. Assoc., 20: 216- 218.
- Raman, K. V. (1988)** : Control of Potato tuber moth, *Phthorimaea operculella* with sex pheromones in Peru. Agric Ecosys. Enviro. 21: 85- 99.
- Raman, K. V. and M. Palacios (1982)** : Screening potato for resistance to potato tuberworm . J.Econ. Entomolo. 75: 47-49.
- Reed, E. M. (1969)** : Agranulosis virus of potato moth. Asut. J. Sci., 31: 300- 301.
- Reed, E. M. and B. P. Springett. (1971)** : Large- Scale field testing of agranulosis virus for the control of the potato tuber moth , (*Phthorimaea operculella*) (zell.) (Lepi : Gelechiidae) . Bull. Entomol. Res. 61: 223- 233.
- Rizk, G. A.; A. A. Abd El-Naby ; E. A. Moftah and M. F. Gergis (1990):** Potential uses of sex pheromones for suppression of potato tuber worm population and control . Purposes. Mania. J. Agric. Development ., 12- 823- 833.
- Rizkalla, M. R.; W. M. El-Deeb and M. A. El-Guindy (1979)** : Esterase activity in relation to insecticides resistance in the Egyptian cotton leafworm (*S. littoralis*). Z. Ang. Ent., 88 :70-76.

- Rofael, M. F. (1990)** : Biochemical genetics of resistance to certain insecticides in the cotton leaf worm, *Spodoptera littoralis* (Boisd.) . Ph. D. Thesis , Fac. Sci. Ain Shams Univ., Egypt.
- Rofael, M. F.; Y. F. Ghoneim ; A. M. allam; F. A. Ayed ; A. M. rashad and M. E. Keddis (1995)** : Innsensitive acetylcholin activities as factors in resistance of pin bollwrom to the organosphorothioate cyanophose. Egypt. J. Appl. Sci., 10 (9) : 477- 496.
- Rose, R. L.; S. Barbhaia; R. M. Roe and E. R. Hodyson (1995)** : Cytochrome P450- associated insecticide resistance and development of Biochemical diagnostic assays in *H. virescens* . Pestic. Biochemem, Physiol. 51 : 178- 191.
- Salah, W. S.; S. A. O. Abdeen ; N. m. Ahmed and A. M. Abd El- Kawy (1986)** : Esterase activity in relation to fenitrothion resistance in cotton leafworm *Spodoptera littoralis* (Boisd.) . Bull. Ent. Soc. Egypt, Econ. Ser. 15 : 73-77.
- Salama, H. S.; M. Regaei and M. Sadbour (1995)** : Larvae of *Phthorimaea operculella* (Zeller) as affected by various of *Bacillus thuringiensis* . J. Appl. Entomol. 119 (3) : 241-243.

- Saleh , W. S. (1981) :** Comparative biochemical studies on resistance and susceptible strains of Egyptian cotton leafworm *Spodoptera littoralis* (Boisd.) . Ph D. Thesis , fac. Agric. Al-Azhar Univ.
- Salem, I. M. (1970) :** Toxicological studies in control leafworm , *Prodenia litura* F. M. Sc. Thesis, Fac. Agric., Al-Azhar Univ., Egypt.
- Salem , S. A. (1991) :** Evaluation of neem seed oil as protectant against *Phthorimaea operculella* (Zell.) (Lepidoptera : Gelechiidae) . Ann. Agric. Science, Moshtohor., 29 (1) : 589- 595.
- Shaaban , A. M. ; M. R. Abo- El-Ghar; M. R. Abd El-Mohymen and M. A. El-Mohymen and M. A. El- Malla (1985) :** Resistance of the Egyptian cotton leafworm *Spodoptera littoralis* (Biosd.) to certain insecticides. Z. Pflanschuty, 92: 69- 75.
- Sheke, S. S.; L. D. Jadhav.; G. N. Salunkhe. (1990) :** Toxicity of some vegetable of oils extracts to *Chelonus blackburni* cam , an egg larval parasite of *Phthorimaea operculella* (Zell.) . Indian. J. Entomol. 52 (4) : 709 – 711.
- Siddig, S.A.; H. Schmutter and K. R. S. Ascher (1987) :** A proposed pest mangement program including neem treatments for combating potato pests in the

Sudan. Natural pesticides from the neem tree and other tropical plants. Proceeding of the 3rd. International Neem conference, Nairobi, Kenya, 10-15 July 1986, 1987, 419-459.

Siddig , S. A. (1988) : Cultural means of controlling potato tuber moth *Phthorimaea operculella* (Zell.) and improvement of potato yield and quality in Sudan. Acta. Horticulture 218: 281-287.

Sigfried, B. D. and j. G. Scott (1992) : Insecticide resistance mechanisms in the German Cockroach, *Blattella germanica* (L.). Acs-Symposium-Series- American- Chemical- Society (USA) , 505: 218-230.

Tabashnik, B. E. (1994) : Evolution of resistance to *Bacillus thuringiensis* . Annu. Rev. Entomol. 39: 47- 79.

Valencia, E.; C. Plata ; C. Corredor and C. Cardona (1993) : Enzymatic resistance to insecticides in larval of *Heliothis verescens* (Fabricius) (Lepidoptera : Noctuidae) . Rev. Colomiana- de- Entomol. 19 (4) : 131- 138.

Van Asperen, K.(1962) : Study of house fly esterases by means of sensitive colorimetric method . J. Insect. Physio., 8 :410-416.

- Wells, d. S.; g. C. Rok and W. c. Dauterman (1983) :**
Studies on the mechanisms responsible for variable toxicity of azinphosmethyl to various larval instars of the tufted apple bud moth , *Platynota idaevsalis*. *pestic. Biochem. Physiol.* 20 : 236.
- Wood, R. j.; N. Pasteur and G. Singre (1984) :**
Charbamate and organophosphate resistance in *Culex pipiens* (L.) (Diptera : Culicidae) in southern France and significance of Est-3 super (A). *Bull. Entomol. Res.*, 74 : 677- 687.
- Yong, B. L. and B. E. Tabashnik (1997) :** Synergism of *Bacillus thuriniensis* by ethylenediamine tetraacetate in susceptible and resistance larvae of diamondback moth, (Lepidoptera : Plutellidae) . *J. Econ. Entomol.* 90 (2) : 287- 292.
- Yu, S. J. (1992):** Detection and biochemical characterization of insecticide resistance to fall armyworm, (Lepidoptera : Noctuidae). *J. Econ. Entomol.* 85 (3) : 675-682.
- Zouh, X. S.; M. G. Zhao (ed.); J. L. Shi (ed). H. Y. Pan and Y. Y. Wang. (1992) :** Biological effect of M-one (containing 5% *Bacillus thuriniensis* var. *San diego*) on the lady beetle *Olla abdominalis* casey
collecting of achievements on the techniques

cooperation –project of R. R. –China and –F.-R.
Germany., 175-178.