

- Sahani, M. S.; Dwivedi, V. K.; Dubey, S. C. and Arora, C. L. (1980). Note on body weight from birth to first breeding and fleece weight in native and cross-bred (quarter-bred) sheep produced under field conditions in semi-arid area of Rajasthan. *Ind. J. Anim. Sci.*, 50(7): 580-581.
- Sannikov, M. E. (1952). The crosses between fine and coarse wool sheep and their improvement. *Sel-Khoz*. Moscow.
- Sanzarias, R. R.; Ovejero, F. J. and Zorita, E. (1974). Development of body weight from birth to 20 weeks of age on Charro lambs. *Anales de la Facultad de Veter de Loon*. 20(2): 101-109. (A. B. A., 44: 4807).
- Sastry, N. S. R and Thomas, C. K. (1976). *Farm Animal Management*. VIKAS Publishing PVT LTD., New Delhi, India.
- Searle, T. W. ; Graham, N. McC.; Donnelly, J. B. and Margan, D. E. (1989). Breed and sex differences in skeletal dimensions of sheep in the first year of life. *J. Agric. Sci. Camb.*, 113: 349-354.
- Seideman, S. C.; Cross, H. R.; Oltjen, R. R. and Schanbacher, B. D. (1982). Utilization of the intact male for red meat production. A review. *J. Anim. Sci.*, 55: 826.
- Selen, S. B. and Banky, E. C. (1961). Wool and body growth in lambs during the first 14 months of life. *Cand. J. Anim. Sci.*, 41: 78-88.
- Seoudy, A. M. (1966). Effect of crossing Merino and Barki sheep on some wool characteristics. M. Sc. Thesis, Ain Shams Univ., Cairo, Egypt
- Seoudy, A. M.; Ghoneim, K. E. and Ghanem, Y. S. (1969).
 I. Effect of crossing Merino with Barki sheep on some wool characteristics.
 II. Grease fleece weight, fiber diameter, crimp, density and fiber type ratio. *J. Anim. Prod., UAR.*, 9(2): 299-301.
- Seoudy, A. M.; Younis, A. A.; Galal, E. S. E.; Ghanam, S. Y. and Khishin, S. S. (1970). Effect of nutrition carcass traits for lambs raised under desert conditions. (Under publication).
- Sharma, S. K.; Bhat, P. P.; Bhat, P. N. and Garg, R. C. (1978). Genetic and nongenetic factors affecting pre-weaning body weight in Muzaffarnagri and Corriedale x Muzaffarnagri half-breds. *Ind. J. Anim. Sci.*, 48: 177-180.
- Shamma, M.; Vasileskov, T. and Sahebi, A. (1977). Growth of early weaned Merino, Zel and crossbred (Merino x Zel) lambs. *Greece.*, 2: 901-902. (A. B. A., 45: 5347).

- Sharaby, M. A. and Suleiman, I. O. (1988). Slaughter weights, carcass weight and dressing percentages of five breeds of sheep slaughtered in Saudi Arabia. *Aust. J. Exp. Agric.*, 28(5): 567-570.
- Sharafeldin, M. A. (1965). Wool characteristics of Iraqi Awassi sheep. *J. Agric. Sci. Camb.*, 65: 223-225.
- Sharafeldin, M. A. and Ghoneim, K. E. (1963). Some non-genetic factors affecting wool characters of Fleish Merino sheep. *Proc. 2nd. Anim. Prod. conf. Egypt*, 419-429.
- Sharafeldin, M. A. and Mostageer, A. (1961). Suckling in Ossimi and Rahmani lambs. *J. Anim. Prod., UAR.*, 1: 53-59.
- Sharafeldin, M. A.; Ragab, M. T. and Ramadan, I. A. (1968). Effect of breeding on lamb production. *J. Agric. Sci. Camb.*, 71: 351-354.
- Shrestha, J. N. B.; Heaney, D. P.; Fiser, P. S. and Langford, G. A. (1984). Influence of breed, birth date, age and body weight on linear body measurements of growth rams maintained in a controlled environment. *Cand. J. Anim. Sci.*, 64: 279-291.
- Sidky, A. R. (1948). The Egyptian sheep, An attempt improve quality and quantity. *Emp. J. Exp. Agric.* 16: 79-89.
- Sidwell, G. M.; Jessup, G. L. and Mcfadden, W. D. (1956). Estimation of clean fleece weight from small side samples and from wool density, body weight, staple length and grease fleece weight. *J. Anim. Sci.*, 15: 218-223.
- Sidwell, G. M.; Price, D. A. and Grandstaff, J. O. (1951). Genetic and environmental factors affecting type and condition in Havajo and Navajo crossbred weaning lambs. *J. Anim. Sci.*, 10: 455-461.
- Sidwell, G. M. and Miller, L. R. (1971). Production in some pure breeds of sheep and their crosses. II. Birth weight and weaning weight of lambs. U.S. department of Agriculture, Beltsville, Maryland 20705.
- Sierra Alfranca and Espejo Diaz, M. (1975). Preliminary results of the generation from crossing argon ewes with Finnish Landrace rams: Growth rates and sexual activity. *Anales Inst. Nac. Invest. Agric. Prod.*
- Singh, B. B. and Singh, M. R. (1974). Relative importance of body measurements for body weight and wool yield in Bikaneri lambs. *Ind. J. Anim. Health*, 12(1): 37-40.

- Singh, B.; Sangwan, M. L. and Poonia, J. S. (1987). Studies on some performance traits and associations among them in Rambouillet sheep: A note. *Ind. J. Anim. Prod. and Management*, 3(3): 139-141.
- Singh, B.; Singh, B and Sangwan, M. L. (1988). Body conformation of sheep at weaning and six months of age. *Ind. J. Anim. Prod. & Management*, 4(1): 27-30.
- Singh, V. K. and Mathur, P. B. (1971). Studies on body measurements indicating mutton production. 48: 829-824. (*Cent. Sheep Wool Res., Substn, Mannavanut*). (*A. B. A.*, 40: 1890).
- Singh, V. K.; Mathur, P. B. and Bohra, D. J. (1973). A note on carcass traits of Coimbatore male lambs weaned at two ages. *Ind. J. Anim. Sci.*, 43(8): 783-785.
- Singh, V. P.; Bapna, D. L. and Kaushik, R. C. D. (1985). Relationship between physical characteristics of Sonadi wool. *Ind. Textile. J.*, 95(9): 85-88.
- Sliwa, Z. and Kaczmarck, F. (1962). The growth of Polish Merino lambs in the first year of life. *Roczn Haukrol.*, B. 80: 435-463. (*A. B. A.*, 32: 181).
- Silwa, Z.; Kozal, E. and Stachowski, J. (1962). Value of the wool of lambs from Leszcynska ewes crossed with Polish Merino rams. (*A. B. A.*, 31: 2941).
- Sojetado, R. M. (1952). The growth and habits of lambs. *Philip Agric.*, 35: 572-578. (*A. B. A.*, 22: 1497).
- Swidan, F. A.; Aboul-Naga, A. M.; El-Shobokshy, S. and Abbass, A. (1978). Performance of Rahmani males lambs weaned at six or eight weeks of age. *Egypt. J. Anim. Prod.*, 19(2): 159-168.
- Tafta, V.; Isacov, Z. and Cirstea, E. (1970). Some morphological and production characters of Fine-wooled sheep from Muscel district. *Laur, Stiint. Inst. Organ, N: Balceson ser. C.* 12 (1969), 79-88. (*A. B. A.*, 40: 1982).
- Taneja, G. C. (1958b). Genetic correlation in Australian Merino sheep. Correlation among body length, width, depth, fore cannon length, mutton score and body weights at 5 and 11 months. *Ind. J. Vet. Sci.*, 28: 216. (*A. B. A.*, 28: 1391).
- Tekeev, E. (1973). Wool production of crossbred fat rumped sheep. *Zhivotonvod Stva, Veterinariii No.*, 5: 30.
- Terrill, C. E.; Kyle, W. H. and Hazel, L. N. (1950). Correlation between traits of range Rambouillet rams. *J. Anim. Sci.*, 9: 640.

- Terrill, C. E.; Pohle, E. M. and Emik, L. O. (1945). Estimation of clean-fleece weight from grease fleece weight and staple length. *J. Agric. Res.*, 70: 1.
- Terrill, C. E.; Sidwell, G. M and Hazel, L. N. (1948b). Effects of some environmental factors on traits of yearling and mature Rambouillet rams. *J. Anim. Sci.*, 7: 311-319.
- Thakur, A. K. B.; Mahajan, J. M. and Singh, O. M. (1967). Study on body weight and conformation of Gaddi and Romney Marsh sheep. *Ind. Vet. J.*, 44: 589-596. (A. B. A., 36: 1968).
- Thrift, F. A. and Whiteman, J. V. (1969). Wool production of Western and Dorset x Western ewes as influenced by certain environmental factors. *J. Anim. Sci.*, 29: 869-873.
- Tkachev, A. YA. (1983). Early weaning of lambs is advantageous. *Ovtsevodstvo No.*, 6: 23-24.. (A. B. A., 51(10): 6017).
- Todorovski, N.; Ristevski, K. and Popovski, K. (1973). Body measurements and weight in Awassi sheep reared in Yugoslavia. *Veterinariski Glasnik* (1973), 27(6): 427-434. (A. B. A., 41: 5285).
- Trivedi, K. R.; Bhat, P. P., Bhat, P. N. and Garg, R. C. (1978). Factors affecting preweaning growth in Muzaffarnagri and its crosses with Dorset and Suffolk breeds of sheep. *Ind. J. Anim. Sci.*, 48: 380-384.
- Vachal, J. (1959). The crossing of Mutton Merinos and Stavrapol Merinos with regard to heterosis. *Sborn. Csl. Aked. Zemed. Ved. Zivac. Vyr.* 4(32): 321-334. (A. B. A., 27: 315).
- Venter, J. J. (1967). A study of the inter relationship between fleece and fiber characteristics in the Merino. IV. Wool production in the Merino. *A. B. A.*, 36: 246.
- Vessely, J. A.; Peters, H. F. and Slen, S. B. (1965). The effect of breed and certain environmental factors on wool traits of range sheep. *Cand. J. Anim. Sci.*, 45: 91-97.
- Vinayak, A. K.; Kanaujia, A. S. and Balaine, D. S. (1982). Genetic and nongenetic factors affecting birth and weaning weights in Nali and its crosses with Merino and Corriedale. *Haryana Agric. Univ. J. Res.*, 12: 558-560.
- Wesile, c. (1963). Recombination of some body measurements in a Southdown-rambouillet F2. *J. Anim. Sci.*, 22: 817. (Abst.).
- Wildman, A. B. (1963). *Wool Ind. Res. Assoc., Torricon. Leads.* (Cited from Yeates, N. T. M., Edey, T. N. and Hill, M. K. (1975), *Animal Science program press, Australia*).

- Wood, I. D.; Macfie, H. J. H. and Brown, A. J. (1983). Effect of body weight, breed and sex on killing-out percentage and non-carass component weights in lambs. *Meat science*, 9(2): 89-99.
- Wynn, P. C. and Thwaites, C. J. (1981). The relative growth and development of the carcass tissues of Merino and crossbred rams and wethers. *Aust. J. Agric. Res.*, 32: 947-956.
- Yalcin, B. C.; Agabakan, S.; Kaseoglu, H. and Singer, N. (1977). Possibilities of exploiting the Rambouillet breed for improvement of production characteristics of Daglic sheep. 1. Fertility, lamb survival and growth rate. *Istanbul Univ. Veteriner Fakultesi Dergesi*, 3: 1-21. (A. B. A., 45: 4387).
- Yalcin, B. C.; Maftuoglu, S. and Yurtcu, B. (1979). Improvement of important production characteristics in Konyo Merino sheep through selection. 2. Some environmental factors affecting production characters. *Istanbul Univ. Veteriner Fakultesi Dergesi*, 5(1): 1-18.
- Yemen Agriculture Research Authority (1986). The animal production in Yemen Arab Republic.
- Younis, A. A.; Kotby, S. and Aamar, G. A. R. (1972). Effect of castration on live body weight and certain carcass traits in Ossimi and Rahmani lambs. *Egypt. J. Anim. Prod.* 1: 19.
- Younis, A. A.; El-shobokshy, A. S.; Hafez, M. and Danasoury, M. S. (1973). Effect of Different levels of energy intake through alteration of roughage to concentrate ratio in rations for fattening lambs. *Alex. J. Agric. Res.*, 21: 229.
- Younis, A. A.; Seoudy, A. M.; Galal, E. S. E.; Ghanam, Y. S. and Khishin, S. S. (1975). Effect of plane of nutrition on feed lot performance and carcass traits of desert sheep. *Trop. Agric.*, 52: 233.
- Youssef, A. A. (1952). Growth and development of Ossimi and Rahmani sheep. M. Sc. Thesis, Fouad 1st. Univ. Cairo, Egypt.
- Youssef, A. A. and Seleem, M. Z. (1962). Some factors affecting selection in the Ossimi and Rahmani flocks. *J. Anim. Prod.*, U. A. R., 2(2): 77-91.
- Zamarozov, A. V. (1963). The meat production of Romanov and Crossbred lambs. *A. B. A.*, 33: 374.
- Zeko, M.; Rushit, F.; Pero, J. and Xuro, K. (1985). The effect of castration on growth of calves and lambs. *Ndikimi Kastrimit per shtimine peshes ne vica e qengja*, Butetini i Shkencane Zooteknike Veterinar, 3(3): 31-36. (A. B. A., 55(8): 5423).