

CONTENTS

	Page
I. INTRODUCTION	1
II. REVIEW OF LITERATURE	3
1- Classification	3
2- The ecology	6
3- The biology	18
III. MATERIAL AND METHODS	22
V. RESULTS AND DISCUSSION	24
A. The ecology	
1- Distribution of the genus <u>Eucera</u> in Egypt	24
2- General localities of existence	32
3- Selection of nesting site	35
4- Nesting activity and characteristics of nesting site	36
5- Nest structure	40
6- The cells and its construction	40
7- Seasonal distribution of different stages	48

(b)

	Page.
B. The Biology	
1- Emergence	55
2- Activity of the newly emerged bees at the nesting site	55
3- Mating	56
4- Provisioning of cells	59
5- Life history.	
a) The egg stage	60
b) The larval stage	60
c) The post defecating larva	62
d) The pupal stage	64
VI. SUMMARY AND CONCLUSION	66
REFERENCES	69
ARABIC SUMMARY	

(a)

LIST OF TABLES

	page.
Table (1): Distribution of genus <u>Eucera</u> in Egypt from the collection of Minis- try of Agriculture	25
Table (2): Distribution of genus <u>Eucera</u> in Egypt From the collection of Entomological society of Egypt.....	26
Table (3): Distribution of genus <u>Eucera</u> in Egypt from the collection of Ain Shams University	27
Table (4): Distribution of genus <u>Eucera</u> in Egypt from the collection of Al-Azhar University	28
Table (5): Distribution of genus <u>Eucera</u> in Egypt from final technical report (study of the insect funa of Egypt, F ₄ - Ent. 2)	28
Table (6): Species of genus <u>Eucera</u> as found to be recorded in the different collec- tions in Egypt	29
Table (7): analysis of soil in the nesting site of <u>Eucera nigrilabris</u> Lep.	34
Table (8): Number of <u>E. nigrilabris</u> Lep. nests in 10 groups occupying 150 m ² of the nesting site	39

	Page
Table (9): Measurements of 10 cells of <u>E.</u> <u>nigrilabris</u> Lep. in mm	47
Table (10): Occurrence of different stages of <u>E.</u> <u>nigrilabris</u> Lep. all year round, 1979	50
Table (11): Occurrence of different stages of <u>E.</u> <u>nigrilabris</u> Lep. all year round, 1980	51
Table (12): Daily activity of newly emerged bees of <u>E. nigrilabris</u> Lep. at the nest- ing site during 3 successive weeks in December 1981	57