

References

REFERENCES

- Abbas, M.S.T. and Y.A.A. El-Deeb (1993).** The natural enemies of major pests infesting cotton in Egypt. Egypt. Nat. Agric. Res., 71 (1) : 1231-138.
- Abdel-Gawaad, A.A.; A.M. El-Sayed; and F.F. Shalaby (1990).** Natural enemies of *B. tabaci* (Genn.) and their role in suppressing the population density of the pest. Agric. Res. Rev., Cairo, 68 (1) : 185-195.
- Abdel-Salam, A.L.; A.M. Metwally; A.A. Yousef; N.A. El- Boghdady and M.F.A. Hegab (1980).** Mites associated with vegetable plants in Egypt. Proc. 1st Conf., Plant Prot. Res. Inst., Cairo, Egypt.
- Annan, I.B.; G.A. Schaefers and W.M. Tingey (1996).** Impact of density of *Aphis craccivora* Koch on growth and yield of susceptible and resistant cowpea cultivars. Ann. Appl. Biol., 128 (2) : 185-193.
- Annan, I.B.; G.A. Schaefers and W.M. Tingey (1997).** Population dynamics and clonal comparisons of cowpea aphid on resistant and susceptible cowpea cultivars. Environ. Entomol., 26 (2) : 250-255.
- Annan, I.B.; G.A. Schaefers; W.M. Tingey; A.M Pong and K. Nyarko (1997).** Interactions of fertilizer, cultivar selection and infestation by cowpea aphid (Aphididae) on growth and yield of cowpea. International of Pest Management, 43 (4) : 307-312.
- A'Holanda, Ac.de and J.V.de Oliveira (1992).** Population fluctuations of *Polyphagotarsonemus latus* Banks, 1904 and *Tetranychus desertorum* Banks on *Phaseolus vulgaris* L. and *Vigna unguiculata* L. Walp. Anais da Sociedade Entomologica do Brasil, 21 (1) : 5-14.

- Badaya, A.K.; S.B. Das; N.S. Thakur; S.S. Shaw and P.L. Bhalla (1990).** Effect of intercultural operations, fertilizer application and plant protection measures on the incidence of borer complex on pigeon pea. *Indian J. Plant Protection*, 18 (1) : 27-30.
- Balajle, K. and R. Veeravel (1995).** Effect of nitrogen on incidence of whitefly, *B. tabaci* (Genn.) in four varieties of brinjal. *Indian J. Entomol.*, 57 (4) : 356- 361.
- Becham, C.M. (1969).** Effect of nitrogen fertilization on abundance of cotton insects. *J. Econ. Entomol.*, 63 : 1219-1220.
- Bentz, J.; J. Reeves; P. Barbosa and B. Francis (1995).** Effect of nitrogen fertilizer source and level on ovipositional choice of poinsettia by *Bemisia argentifolii* (Homoptera : Aleyrodidae). *J. Econ. Entomol.*, 88 (5) : 1388-1392.
- Berg, G.N. (1984).** Effect of temperature and host species on the population growth potential of the cowpea aphid, *Aphis craccivora* Koch. *Aust. J. Zool.*, 32 : 345-352. **Bhangal, G.T. and M.B. Joi (1985).** Screening of onion cultivars for resistance to purple blotch and thrips. *J. Maharashtra Agric. Univ.*, 10 (3) : 355-356.
- Bhagabati, K.N. and B.K. Goswami (1992).** Incidence of yellow yellow vein mosaic virus disease of okra, *Abelmoschus esculentus* L. Moench in relation to whitefly, *Bemisia tabaci* (Genn.) population under different sowing date. *Indian J. Virology*, 8 (1) : 37- 39.
- Bocj, K.R.; E.J. Guthrie and R.D. Woods (1974).** Purification of maize streak virus and its relationship to viruses associated with streak diseases of sugarcane and *Panicum maximum*. *Ann. Appl. Biol.*, 77 : 289-296.

- Brar, K.S.; A.S. Sidhu and M.L. Chadha (1993).** Screening onion varieties for resistance to *Thrips tabaci* Lind. and *Helicoverpa armigera* Hubner. J. Insect Sci., 6 (1) : 123-124.
- Byers, R.A. and W.C. Templeton, Jr. (1988).** Effect of planting date placement of seed, vegetation suppression slugs and insects upon establishment of no-till alfalfa. In Report of North American Alfalfa Improvement Conference, June 10-23.
- Carman-Sanchez, M.; M. Cermeli and D. Diaz (1993).** Aphid fauna (Homoptera : Aphididae) on *Phaseolus vulgaris* in Aragua State. Boletin de Entomology Venezolana, 8 (1) : 75-81.
- Chapman, A.J.; C.A. Richmond and L.C. Fife (1947).** Comparative toxicity of benzene hexachloride and DDT to Thrips on cotton and onion. J. Econ. Entomol., 60 : 575-576.
- Coll, M. and D.G. Bottrel (1995).** Predator-prey association in mono- and dicultures effect of maize and bean vegetation. Agric. Ecosys. & Environ., 54 (1-2) : 115-125.
- Costa, H.S. and D.N. Byrne (1988).** Neutron activation of rare element to mark the sweet potato whitefly, *Bemisia tabaci* J. Ecol. Entomol., 13 (4) : 465-467.
- Darshan, Singh, A.S.; J.C. Sidha-Thakur and D. Singh (1986).** Relative resistance of onion and garlic cultivars to thrips, *Thrips tabaci* Lind. J. Res. Punjab Agric. Univ., 23 (3) : 424-427.
- Darwish, Y.A. and A.I. Farghal (1990).** Evaluation of certain pesticides activity against the cotton whitefly, *Bemisia tabaci* and associated natural enemies on cotton plants under field conditions in Assiut. Assiut J. Agric. Sci., 21 (5) : 331-339.

- Denoyes, B.; F. Bordat; H.DE. Bon and P. Daly (1988).** A new pest of vegetable crops in Martinique, *Thrips palmi* Karny (Thysanoptera). Bull. Agron. des Antilles et de la Guyane, 7 : 16-19.
- Dissemond, A. and H. Hindorf (1990).** Influence of sorghum/maize/cowpea intercropping in the insect situation at Mbita/Kinya. J. Appl. Entomol., 109 (2) : 144-150.
- Djuwarso, T. and F. Suryawan (1989).** Major insects pests of legumes. Seminar Gasil Penelitian Tanaman Pangan Balittan Bogor, Indonesia, pp. 13-14.
- Donchen, K.D. and P. Tomov (1996).** Species composition and seasonal dynamic of Thrips (Thysanoptera) on *Festuca arundinaca* Schreb. Folia Entomologica, 62 "Suppl." 33-36.
- Eichelkraut, K. and C. Cardona (1989).** Biology mass rearing and ecological aspects of the whitefly, *B. tabaci* (Genn.) as a pest of beans. Turria Lba., 39 (1) : 55-62.
- Ekukole, G. (1992).** Effect of some agronomic and chemical control practices on *A. gossypii* populations and stickiness in cotton. Cotton et Fibres Tropicales, 47 (2) : 139-143.
- El-Adl, F.E.; A.M. Abdel-Hameid and A.Z. Osman (1983).** Effect of nitrogen levels and insecticides on population density of *Eriophyes tulipae* K., *Thrips tabaci* Lind. and productivity of Chinese garlic seeds. J. Appl. Sci., Egypt, 8 (11) : 319-347.
- El-Hawary, F.M.A. (1980).** Studies on some extrinsic factors affecting the bionomics of *A. craccivora* Koch infesting beans in Egypt. M.Sc. Thesis, Fac. of Science, Cairo Univ.

- El-Heneidy, A.H.; M.S.T. Abbas and M.F. El-Dakroury (1978).** Seasonal abundance of certain predators in untreated Egyptian clover and cotton fields in fayoum Governorate, Egypt. Bull. Soc. ent. Egypté, 62 : 82-95.
- El-Heneidy, A.H.; A.H. Abbas and A.A. Khidr (1987).** Comparative population densities of certain predators in cotton fields treated with sex pheromones and insecticides in Monofia Governorate, Egypt. Bull. ent. Soc. Egypt, Econ. Ser., 16 : 181-190.
- El-Kady, W.A.; I.S. El-Hawary and S.E.D.A. Fawzy (1988).** Effect of nitrogen and potassium fertilizers on seed cotton yield and insect pests infestation. J. Agric. Res., Tanta Univ., 14 (2) : 1426-1437.
- El-Khateeb, H.M. (1977).** Ecological factors affecting *Tetranychus atabicus* Attiah infesting truck crops. M.Sc. Thesis, Fac. Agric., Ain Shams Univ., Egypt.
- El-Khawalka, M.H.M.; M.A. El-Bessomy; S.H. Mansour and H.M. Bastawisy (1996).** Susceptibility of some faba bean genotypes to field infestation with *A. craccivora* koch and *Bruchus rufimanus* Boh. with special references to yield. J. Agric. Sci., Mansoura Univ., 21 (11) : 4129- 4132.
- El-Khayat, E.F. (1994).** Effect of nitrogen fertilization on wheat aphids infesting bioenemies and grain production. Ann. Agric. Sc., Moshtohor, 32 (1) : 557-566.
- El-Khouly, A.S.; M.A. Ali; I. Ibrahim and S.A. Naga (1994).** Effect of intercropping maize and cowpea on their susceptibility to infestation with aphids. Bull. ent. Soc. Egypté, 72 : 227-235.
- El-Sayed, A.M.; F.F. Shalaby and A.A. Abdel-Gawaad (1991).** Ecological studies on *Bemisia tabaci* (Genn.) (Homoptera : Aleyrodidae) infesting different host plants. 1- Fluctuation and population density of *B. tabaci* on different host plant. J. Agric. Res., Egypt, 69 (1) : 193-207.

- El-Serwiyy, S.; H. El-Haidari and Saad (1984). Population density of whitefly, *B. tabaci* on fall cucumber in Iraq. J. Agric. & Water Resources Res., 3 (2) : 78-87.
- El-Sheikh, M.A.K.; S. El-Nagar; F.M. Maklad and S.S.A. Kabeil (1994). Aphid fauna and seasonal incidence on potato and associated with plants at Giza, Egypt. Bull. ent. Soc. Egypt, 72 : 203-215.
- Farag, A. (1966). Studies on whiteflies (Homoptear : Aleyrodidae) in Egypt. Ph.D. Thesis, Fac. of Agric., Ain Shams Univ.
- Farghali, M.A.; A.G. Ali and H.A. Hussein (1996). Susceptibility of some mungbean cultivars to Aphid infestation with special reference to pod setting and seed yield. Assiut J. Agric. Sci., 27 (2) : 125-134.
- Farrag, A.M.I.; A.S. Abdel-Salam; M.L. Wahba and M.S. Abul-Ela (1980). The spider mite *Tetranychus arabicus* Attiah on soybean plants (Acari : Tetranychidae). Proc. 1st Conf., Plant Prot. Res. Inst., Cairo, Egypt, Vol. 3 : 31-39.
- Flexner, J.L.; P.H. Westigard; P. Gonzalves and R. Hilton (1991). The effect of ground cover and herbicide treatment on two spotted spider mite density and dispersal Southern Oregon pear orchards. Ent. exp. et appl., 60 (2) : 111-123.
- Fowler, H.D. (1956). Some physiological effects of attack by whitefly, *B. tabaci* (Genn.) and spraying Parathion on cotton in the Sudan Gezira. Empire Cotton Growing Review, 33 : 288-299.
- Gerling, M. Motro and R. Horowitz (1980). Dynamic of *B. tabaci* attacking cotton in the coastal plain of Israel. Bull. Ent. Res., 70 : 213-219.
- Gillespie, D.R. (1989). Biological control of Thrips on greenhouse cucumber by *Amblyseius cucumeris*. Entomophaga, 34 (2) : 185-192.

- Habib, A.; G.N. Rezk; H.T.A. Farghaly and Z.A. Ragab (1976).** Seasonal abundance of some predators in cotton and its relationship to certain pests. Bull. Soc. ent. Egypté, 60 : 191-186.
- Habibi, J. (1975).** The cotton whitefly, *B. tabaci* : Biology and methods of control. Entomol. Phytopathol., 38 : 13-36.
- Hafez, M.; A.M. Ali; F. Mikhail and Abd El-Kawi (1975).** Fluctuation of population of certain predators in cotton and clover fields in Assiut Governorate, Egypt. Ibid, 53 (1) : 47-51.
- Hamdy, M.K. (1992).** Some ecological aspects on the leaf hopper, *Empoasca decipiens* attacking summer vegetable plants in Sharkia Governorate during 1989. Ann. Agric. Sci., Cairo, 37 (2) : 613-620.
- Hamdy, M.K. and M. Salem (1994).** The effect of planting dates of onion temperature and relative humidity on the population density of the onion thrips, *Thrips tabaci* Lind. in Egypt. Ann. Agric. Sci., Ain Shams Univ., Cairo, 39 (1) : 417-424.
- Hamid, S.; M.A. Shah and A.M. Anwar (1977).** Some ecological and behavioural studies on *A. craccivora* Koch. tech. Bull. Common Inst. Biol. Cont., 18 : 99- 111.
- Hamstead, E.D. and E. Gould (1957).** Relation of mite populations to seasonal leaf nitrogen levels in apple orchards. J. Econ. Entomol., 50 : 109-110.
- Hance, T. and A. Pasleau (1987).** Studies of tripartite relationship between haricot bean, *Tetranychus urticae* and *Phytoseiulus persimilis* and of the positive impact of the predator on the food plant. Mededlingen van de Faculteit Landbouewtenschappen Rijksuniversiteit Gent., 52 (2a) : 433-448.

- Harewijn, P. (1970).** Reproduction of aphid, *Myzus persicae* related to the mineral nitrogen of potato plants. Ent. Exp. & Appl., 13 : 307-319.
- Hassanein, M.H.; F.M. Khalil and M.A. Eisa (1970).** Contribution to the study of *Thrips tabaci* Lind. in Upper Egypt. Bull. Soc. ent. Egypté, 54 : 133-140.
- Hayder, M.F. and L.S. Sherif (1987).** Ecological aspects and developing method of onion pests control. Bull. Soc. ent. Egypt, Econ. Ser., 16 : 119-126.
- Helaly, M.M.; A.M. Ibrahi, and M.R.A. Saleh (1982).** Fluctuation of population densities of *Empoasca* sp., *A. craccivora* and *Tetranychus arabicus* Attiah attacking cowpea plants at Zagazig, Egypt. Bull. Soc. ent. Egypté, 64 : 35-43.
- Henneberry, T.J. and D. Schriver (1964).** Two-spotted spider mite feeding on bean leaf tissue of plants supplied various level of nitrogen. J. Econ. Entomol., 57 : 377-379.
- Heyer-Wilok, M.L.C. and B. Cruz (1989).** The population dynamics of the cotton whitefly, *Bemisia tabaci* (Genn.) in bean fields in the Republic of Cuba. Archiv fur Phytopathologie und Pflanzenschutz, 25 (5) : 473-479.
- Heyer, W.; R. Aviles; M.L. Chiang-Lok and B. Gruz (1994).** Towards more effective chemical control measures against pests of bean *Phaseolus vulgaris* L. in Cuba. Archives of Phytopathology & Plant Protection, 29 (1) : 89-100.
- Hijam, R.S. and T.K. Singh (1989).** Seasonal incidence of *A. craccivora* Koch on cowpea, *Vigna unguiculata* L. Walp in manipur. J. Aphidology, 3 (1-2) : 68-72.
- Hilje, L.; D. Cubillo and L. Segura (1993).** Ecological observations on the whitefly, *B. tabaci* Genn. in Costa Rica. Manejo Integrado de Plagas, 30 : 24-33.

- Hoda, F.M. and I.I. Mohamed (1982).** Susceptibility of sixteen different varieties of soybean to spider mite infestation. Egypt Natural Conf. of Entomology Organized by the Ent. Soc. of Egypt (Abst. Vol.).
- Hoda, M.F. and S.A. Doss (1984).** Studies on the susceptibility of soybean varieties of major pests infestation. Proc. 2nd General Conf. of ARC, Giza, 9-11 April.
- Isely, D. (1946).** The cotton aphid. Bull. Ark. Agric. Exp. Sta., 9 : 26-29.
- Johnson, M.W.; R.F.L. Mau; A.P. Martinez and S. Fukuda (1989).** Foliar pests of watermelon in hawaii. trop Pest Manag., 35 (1) : 90-96.
- Jasson, R.K.; G.L. Leibe; C.A. Sanchez and S. Hlecrone (1991).** Effect of nitrogen and foliar biomass on population parameters of cabbage insects. Ent. exp. et appl., 61 (1) : 7-10.
- Kaazeh, W. and J.D. Dutcher (1993).** Population parameters and probing behaviour of cowpea aphid (Homoptera : Aphididae) on preferred and non-preferred host crops.
- Kornegay, J.L.; C. Cardona; J. Van Esch and M. Acrardao (1989).** Identification of common bean lines with ovipositional resistance to *Empoasca kraemeri* (Homoptera : Cicadellidae). J. Econ. Entomol., 82 (2) : 649-654.
- Kuang, R.P. and N. Xiao (1997).** Susceptibility of broad bean cultivars to cowpea aphid, *Aphis craccivora* Koch and pea aphid, *Acyrtosiphon pisum* Harris. J. Ent. Res., 21 (3) : 219-222.
- Kumar, A.; M.K. Tripathy and K.M. Srivastava (1997).** Population dynamics of black aphid, *A. craccivora* koch in different genotypes of Indian bean, *Dolichos lablab* in relation to weather parameters at Kampur, UP. Environ. & Ecol., 15 (2) : 318-321.

- Kuroli, G.; I. Nemeth and L. Nemeth (1988).** Aphid damage to field bean in relation to population dynamics and ecological conditions. Mededelingen van de Faculteit Landbouwwetenschappen Rijksuniversiteit Gent., 53 (3a) : 1195-1201.
- Lal, O.P. (1992).** Evaluation of certain insecticides against the pest complex of french bean seed crop under field conditions. J. Ent. Res., 16 (1) : 57-61.
- Legge, J.B.B. and J. Palmer (1968).** Behaviour of *Myzus persicae* Sulzer on tobacco oil choice sites for feeding and larviposition on levels and artificial substrates of differing nutrient states. Bull. Ent. Res., 57 : 393-479.
- Li, D.X.; J. Tiang and Z.Y. Zhang (1994).** A study on the bionomics and control techniques of *A. craccivora* Koch on mung beans. Plant Protection, 20 (6) : 4-5.
- Liegh, T.F.; D.W. Greemes; H. Yamado; D. Bassett and J.R. Stochton (1974).** Insects in cotton as affected by irrigation and fertilization practices. Calif. Agric., 24 (3) : 12-14.
- Lorini, T.; L. Torres and D.R. Guimaraes (1986).** Population fluctuations of *Thrips tabaci* in an onion crop. Pesquisa em Andament Empresa Contarinese de Pesquis agropecuaria, Embrapa, Brazil, 62, 4 pp.
- Lorini, T. and V.M. Junior (1990).** Population fluctuation of *Thrips tabaci* Lindman 1988 (Thysanoptera : Thripidae) on garlic crop. Anaid da Sociad ate entomologica de Brazil, 19 (2) : 367-371.
- Mansour, M.H.; N.Z. Dimetry and I.S. Rofael (1982).** The role of coumarin as secondary substance in the food specificity of the cowpea aphid, *A. craccivora* Koch. Z. Angew. Ent., 93 : 151-157.

- Mostafa, N.R.; G.A. Karaman; M.A. Ali and A.A. Abdel- Alim (1983).** Effect of different levels of nitrogen fertilization on the infestation of soybean plants. *Minia J. Agric. Res. Res.*, 5 (1) : 143-150.
- Mrig, K.K. and Ram Singh (1985).** Incidence of insect pests on garden bean *Dolichos lablab* Linn. *Bull. Entomol.*, 26 (1) : 5-7.
- Musuna, A.C.Z. (1983).** Potentially serious pest whitefly, *B. tabaci* (Genn.). *Zimbabwe Agric. J.*, 8 (4) : 143-146.
- Nasr, E.A.; S.M. Hafez and A.M. Hossain (1982a).** Effect of cultural practices on mites inhabiting cotton plants. 1- Sowing date. *Fac. Agric., Ain Shams univ., Res. Bull.*, 963, 1-9.
- Nasr, E.A.; S.M. Hafez and A.M. Hossain (1982b).** Effect of cultural practices on mites inhabiting cotton plants. 2- Sowing date. *Fac. Agric., Ain Shams univ., Res. Bull.*, 963, 1-9.
- Naguib, M.A. (1980).** Effect of the predators population in clover fields and their population in adjacent cotton field. *Proc. 1st Conf., Plant Prot. Res. inst.*, pp. 309-321.
- Naresh, J.S. and Y.L. Nene (1980).** Host range, host preference for oviposition and development on the dispersal of *Bemisia tabaci* (Genn.), a vector of several plant virus. *Indian J. Agric. Sci.*, 50 : 620- 623.
- Nassef, M.A.; A.A. El-Feshawi; M.K.A. Abo-Sholooa and W.M. Watson (1996).** Seasonal abundance of certain predators as influenced by cotton planting dates and weather factors. *Egypt. J. Agric. Res.*, 74 (3) : 585.
- Ngugi, E.C.K.; A. Shakoori and P.G.A. Omanga (1985).** Breeding for resistance against some of the major insects of cowpea and pigeon pea. *Insect Sci. Appl.*, 6 : 365-367.

- Nour El-Din, M.M. (1984).** Ecological studies on the insects attacking Cucurbitaceae. M.Sc. Thesis, Fac. of Agric., Al-Azhar Univ., Egypt.
- Ofuya, T.I. (1986).** Predation by *Cheilomenes vicina* (Coleoptera : Coccinellidae) on the cowpea aphid, *A. craccivora* Koch : Effect of prey stage and density. Entomophaga, 31 (4) : 331-335.
- Ofuya, T.I. and A.E. Akingbohunge (1988).** Functional and numerical responses of *Cheilomenes lunata* (Fabricius) (Coleoptera : Coccinellidae) feeding on the cowpea aphid, *A. craccivora* Koch. Insect Sci. & Appl., 9 (4) : 543-546.
- Ofuya, T.I. (1991).** Aspects of predation in *Cheilomenes lunata* on cowpea aphid, *A. craccivora* Koch. Z. Angew. Zool., 78 (4) : 413-419.
- Ofuya, T.I. (1995).** Studies on the capability of *Cheilomenes lunata* to prey on the cowpea aphid, *A. craccivora* Koch. Agric. Ecosys. & Environ., 52 (1) : 35-38.
- Omar, H.I.H.; M.F. Hegab and A.E.M. El-Sorady (1993).** The impact of intercropping cotton and cowpea on pest infestation. Egypt. J. Agric. Res., 71 (3) : 709-716.
- Pai, K.M. and A.V. Dhri (1991).** Incidence of insect pest in early variety of cowpea, *Vigna unguiculata* (L.) Walp. Indian J. Entomol., 53 (2) : 329-331.
- Patial, A.P.; R.N. Nawal; D.S. Ojri and P.R. Moh;kar 91988).** Field screening of onion cultivars for their reaction to thrips. Indian Cocoa Arecanut & Species Journal, 12 (1) : 10-11.
- Patel, M.B. and K.P. Srivastava (1990).** Field screening of some high yielding genotypes of mung bean, *Vigna radiata* (Linnaeus) Wilczek to whitefly, *B. tabaci* (Genn.) and yellow mosaic virus (YMV). Indian J. Entomol., 52 (4) : 547-551.

- Patel, H.M. and R.C. Jhala (1992).** Studies on host range, host preference and population dynamics of whitefly, *B. tabaci* in South Gujarat, India. Gujarat Agric. univ. Res. J., 17 (2) : 76-81.
- Paiva, F.D.A. and A.C.P. Goulart (1995).** Variation of the whitefly population and incidence of bean golden mosaic at Dourado. MS. Fitopatologia, Brasileira, 20 (2) : 199-202.
- Pearson, E.O. (Assisted by R.C. Maxwell Darling) (1968).** The insect pests of cotton in tropical Africa, pp. 215-226.
- Perkins, H.H. (1983).** Identification and processing of honey dew contaminated cotton. Textile Research J., pp. 508-512,
- Perkins, H.H. (1987).** Sticky cotton. Proceeding of the Western Cotton Production Conference, pp. 53-55.
- Reddy, K.D.; V.P. Gargav and D.S. Misra (1983).** Studies on host preference of *A. craccivora* Koch. Entomon., 8 : 75-78.
- Rizk, G.A.; G.A. karaman; A.A. Abdel-Naby; E.A. Moftah and A.A. Abdel-Alim (1990).** Interrelation between the fluctuation in the populations of predators and parasites and the fluctuations in the population density of sucking pests associated with soybean plant in Minia region. Assiut J. Agric. Sci., 21 (3) : 105- 117.
- Rizk, G.A.; E.A. Moftah; G.A. Karaman and A.A. Abdel- Naby (1990).** Effectiveness of different planting dates on the population density of some sucking pests attacking soybean plants in Minia region. Assiut J. Agric. Sci., 21 (3) : 141-151.
- Rote, N.B. and S.N. Puri (1992).** Effect of fertilizer application on incidence of whitefly on different cotton cultivars. J. Maharashtra Agric. Univ., 17 (1) : 45-48.
- Sato, M.E. and Nakano (1990).** Influence of planting period on the main pests attacking onion in Egypt. Ann. Agric. Sc., Moshtohor, 13 : 65-75.

- Sawires, Z.R. (1978).** Ecological studies on mites associated with soybean plants. M.Sc. Thesis, Fac. of Agric., Cairo Univ.
- Sawires, Z.R.; H.A. Taha and S.T. Abdalla (1990).** Biological and ecological studies on *Tetranychus arabicus* and relative susceptibility of seventeen soybean genotypes to infestation. Agric. Res. Rev., 68 (1) : 17-24.
- Schwartz, A. (1988).** Population dynamics of *Thrips tabaci* Lindeman (Thysanoptera : Thripidae) on table grapes. S. Afr. J. Ecol. & Viticulture, (1) : 19-21.
- Sewify, G.H.; S.A. El-Arnaouty and M.H. Belal (1996).** The effect of cotton late planting on population densities of sucking insects and their associated predators in Giza region, Egypt. Bull. Fac. Agric., Univ. Cairo, 47 (4) : 664-675.
- Shalaby, F.F.; A.A. Abdel-Gawaad; A.M. El-Sayed and M.R. Abo-El-Ghar (1990).** Natural role of *Eretmocerus mundus* Mercet and *Prospaltella lutea* Masi on populations of *B. tabaci* Genn. Agric. Res. Rev., Cairo, 68 (1) : 197-208.
- Shimoda, T.; J. Takabayashi; W. Ashihara and A. Takafuji (1997).** Response of predatory insect, *Scolothrips takahashii* toward herbivore induced plant volatiles under laboratory and field conditions. J. Chem. Ecol., 23 (8) : 2033-2048.
- Singh, J. and R.N. Singh (1987).** Red spider mite *Tetranychus ludeni* Zacher, a new threat to summer bean. Manila (Philippines), 1 p.
- Singh, O.P.; K.J. Singh and P.P. Singh (1988).** Effect of dates of sowing and varieties on the incidence of major insect pest of soybean in Madhya Pradesh. Bhartiya Krishi Anusandana Patrika, 3 (1) : 47-52.
- Singh, O.P.; K.J. Singh and K.N. Kapoor (1990).** Sesaonal incidence and chemical control of red spider mites *Tetranychus telarius* Linn. on soybean in madhya Pradesh, India. Indian J. Ent., 52 (1) : 57-62.

- Singh, R.N. (1995). Studies on seasonal abundance of *T. ludeni* zacher on cowpea, *Vigna unguiculata* Sav. J. Recent Adv. Appl. Sci., 10 (1-2) : 59-63.
- Smith, L.M.; W.O. Lamp and E.J. Armrust (1994). Behaviour and reproductive response of caged *Empoasca fabae* to vegetation density of a host legume and a non host grass. J. Ent. Sci., 29 (1) : 66-81.
- Storms, J.J.H. (1969). Observation on the relationship between nutrition of apple rootstocks in gravel cultural and reproduction of *T. urticae* (Acarina : Tetranychidae). Ent. Exp. & Appl., 12 : 297-311.
- Thomas, G.D.; C.M. Ignoffo and D.B. Smith (1976). Influence of defoliation and deposing on quality of soybean. J. Econ. Entomol., 69 : 737-740.
- Tonhasca, A., Jr. (1994). Response of soybean herbivores to two agronomic practices increasing agroecosystem diversity. Agric. Ecosys. & Environ., 48 (1) : 57-65.
- Verma, S.N.; V.P. Gargav and M.V.S. Menon (1983). On the life history and host preference of the bean aphid, *A. craccivora* Koch. Pranikee, 4 : 389-395.
- Vetten, H.J. and D.J. Allen (1983). Effect of environment and host on vector biology and insecticides of two whitefly spread diseases of legumes un Nigeria. Ann. App. Biol., 102 (2) : 219-277.
- Vicente, M.; R.D. Kanthack; A.B. Noronha and M.F.S. Stradiota (1988). Incidence of golden mosaic on bean grown in two planting seasons in the president Pridente Region. Fitopatologia Brasileira, 13 (4) : 373-376.

- Vierbegen, B. (1990).** The tobacco thrips, *Thrips tabaci* Lind. and tomato spotted wilt virus in the Netherlands. *Gevasbes Cherming*, 21 (6) : 159-164.
- Vogt, H. (1994).** Effects of pesticides on *Chrysoperla carnea* (Steph.) (Neuroptera : Chrysopidae), in the field and comparison with laboratory and semi-field results. *Bull. Oil B.S.Rop*, 17 (10) : 71-82.
- Wahba, M.L.; S.A. Doss; F.S. Faris and M.K. Nakhla (1986).** *Phaseolus vulgaris* cultivars to infestation with aphids and mites. *Agric. Res. Rev., Egypt*, 69 (1) : 163-170.
-