

REFERENCES

REFERENCES

- Abbas, M. S. T. and Y. A. A. El-Deeb (1993):** On the natural enemies of the major pests infesting cotton in Egypt. Egypt. J. Agric. Res. 71 (1): 131-138.
- Abbott, W. S. (1925):** A method of computing the effectiveness of an insecticide. J. Econ. Entomol. (18):265-277.
- Abd El-Rahman - Soheir, I. (1996):** Studies on the Tetranychoid mites. Ph.D. Thesis, Fac. Agric. Cairo Univ. 350pp.
- Abd Allah, Y. E. Y. and S. M. O. Kelany (2003):** Ecological factors affecting the population activity of sap sucking Agro-Ecosystems. Arab Univ. J. Agric., Ain Shams Univ. Cairo. 11 (1): 359-371.
- Abd El- Wahed, N. M. (1997):** Toxicological and biological studies on *Tetranychus urticae* (Koch) associated with fruit orchards. M. SC. Thesis. Fac. Agric. Al- Azhar Univ 188pp.
- Abou- Awad, B. A.; El-Sawaf, B. M.; and A. A. Abd-ElKader (1999):** Life history and life table of *Pronematus ubiquitus* (McGregor) as a predator of eriophid mites in Egypt (Acari: Tydeidae). Acarologia. 40(1): 29-32.
- Abou- El-Ghar, M. R.; El- Badry, E. A.; Hassan, S. M. and S. M. El-Kilany(1971):** Effect of some pesticides on the predatory mite *Agistemus exsertus*. J. Econ. Entomol. (54): 26-27.
- Ahmed, N. F. R. (2003):** Studies on some Arthropods inhabiting cucurbits and beans M.sc. Thesis Fac. Agric. Cairo Univ. 105 pp.
- Amer, A. I. (2003):** Ecological and biological studies on certain mites infesting cotton, other field crops and associated predators. M. Sc. Thesis Fac. Agric. Al- Azhar. 104pp.
- Andreeva , I. V. and M. V. Shternshis (1995):** Microbiologica formulations against web mites in greenhouses. Zashchita-Rastenii-Moskva. (11):41-42.

REFERNCES-----

- Attiah, H. H. (1969).** The tetranychid mites of the U. A. R. I. The genus *Tetranychus* Dufour (Acarina: Tetranychidae). *Acarologia*. 11 (4): 733-741.
- Azouz, H. A. A. (2005):** Ecological and biological studies on some mites associated with cotton and some field crops in Beni-suef Governorate. Ph. D. thesis. Fac. Agric. Al- Azahr univ. 185pp.
- Baker, E. W. and G. W. Wharton (1952):** An introduction to Acarology. The MacMillan Co. New York. 465pp.
- Bayram, A; Varol, M. I. and I. H. Tozan (2000):** The spider (Araneae) fauna of the cotton fields located in the western part of Turkey *Serket*. 6 (4): 105-114.
- Botha, J. H.; Du-plassis, D. and F. J. Calitz (1994):** Effect of acaricides on phytophagous mites and selected predators in apple orchards. *J. Southern Afric. Soci. Horti. sci.* (1):47-49.
- Brickhill, C. D. (1958):** Biological studies of two species of Tydeid mites from Calif. *Hilgar*. (27): 601- 620.
- Cai, S. H. (1987):** Studies on the occurrence of *T. cinnabarinus* and its control. *Iarget. China- Cotton*. (5): 39-40.
- Canerday, T.D. and F.S. Arant (1964):** The effect of spider mite population on yield and quality of cotton . *J. Econ. Entomol*. 57(4): 553-556.
- Chant, D. A. (1959):** Phytoseiid mites (Acarina: Phytoseiidae) I. Bionomics of seven species in Southeastern England. II.A taxonomic review of the family Phytoseiidae, with descriptions of 38 new species. *Can. Entomol*. 91 (12): 164pp.
- Chant, D. A. (1965).** Generic concepts in the family Phytoseiidae (Acarina : Mesostigmata). *Can. Entomol*. (97): 351-374.
- Chandler, D.; Davidson, G.; Pell, J. K.; Ball, B. V.; Shaw, K. and K. D. Sunderland (2000):** Fungal biocontrol of Acari. *Biocont. Sci. Tech*. 10(4): 357-384.

REFERNCES

- Cividanes, F.; Tromozine, M. J. and L. H. Silvestre (1987):**
Distribution of *Tetranychus urticae* (Kock, 1936) (Acari :
Tetranychidae) on cotton plants. Cientifica-jaboticabla, 15 (2): 147-
155.
- Duncan, D. B. (1955):** Multiple range and multiple F test. Biometrics.
(11):1-42.
- EL-Adawy, A. M.; Yousri, A. Y. M. and T. EL-Sharkawy (1995):**
Effect of some acaricides and the Biocide natural – L(*Beauveria*
bassiana) on the two-spotted spider mite, *Tetranychus urticae* Koch
infesting cucumber under plastic house conditions. 6th Nat. Conf. of
pests & Dis. of Veg. & Fruit in Egypt and Arab count. Ismailia. Egypt:
136-141.
- EL- Badry, E. A. (1967):** Five new phytoseiid mites from U. A. R. with
collection notes on three other species (Acarina: Phytoseiidae). India, J.
Entomol. (29):177-184.
- EL-Hady, M. M. (1996):** Studies on tetranychoid mites and their
predators in Egypt. Ph. D. Thesis, Fac. Agric. Cairo Univ. 164
pp.
- EL-Heneidy, A. H.; Amira, A. Ibrahim; Fayad, Y. H . and G. M. Moawad**
(1996): Survey and population dynamics of common true
spiders in Egyptian cotton fields. Ann. Agric. Sci., Moshtohor. 34(3):
1177-1187.
- EL-Erksousy, M. H. M. (2000):** Studies on some true spiders in
Egypt. Ph. D. Thesis, Fac. Agric., Al-Azhar Univ. 132pp.
- EL-Maghraby, M. M. A.; El-Tantawy, M. A.; Gomaa, E. A. A. and**
M. Nada (1994): Effect of pesticide application during the
early and late cotton season on the abundance of the certain
predators associated with cotton plants. Anz. fur- schadl – pflanz -
umwelt., 67(3): 60-62.
- EL-Massry, M. K. (1978):** Studies on the predators attacking spider
mites on cotton. M .Sc. Thesis, Fac. Agric. Zagazig. Univ. 153
pp.

REFERNCES-----

- EL-Naggar, M. E.; Abd-El-Halim, M. A.; Shoeib, A. A.; Dugger, P. and D. Richter (1999):** True spiders as a biocontrol agent for controlling spider mites in Egypt. Proceed. Beltw. cotton Confer, Orlando, Florida, (4): 1125-1126
- EL-Naggar, M. E.; Sallam, G. M. and M. A. Abd-Halim (2000):** Survey of true spiders associated with some field crops in Egypt. Proceed. Beltw. Cotton Confer. (2): 1021-1022.
- EL-Safty, A. F. A. (2003):** Studies on phytophagous mites associated with fruit trees in Egypt. Ph.D. Thesis Fac. Agric. Cairo Univ. 166 pp.
- EL- Saadany, G.; El-Badry, E. A .; Hanna, M. and R. A. Rizk (1977):** Simultaneous effect of three main weather factors on two spotted Spider mite populations. Bull. Agric. Res. Rev. (55): 143-150.
- Finney, D. G. (1952):** Propit Analysis - Statistical Treatment of the Sigmoid Response Curve. Cambride. Univ. Press. 318 pp.
- Furr, R.E. and T. R. Primer (1968):** Effect of early – mid- and late- season infestation of low spotted spider mites on the yield of cotton. J. Econ. Entomol. (61): 1446-1447.
- Gerson, E. (1968):** Five tydeid mites from Israel (Acarina: Prostigmata). Israel. J. Zool. (17): 191-198.
- Grafton- Cardwell, E. E.; Granett, J. and S. M. Normington (1991):** Influence of dispersal from almonds on population dynamics and acaricides resistance frequencies of spider mite infestation neighboring cotton. Exp. and Appl. Acarology. 10(4): 187- 212.
- Gupta, S. K.; Dhooria, M. S. and A. S. Sidhu (1971):** A note on predators of citrus mites in punjab. Science and Culture. (37):484pp.
- Gutierrez, J. P. (1967):** Contribution on letude morphologigue et biology de *Tetranychus andre* Gutierrez (Acarina: Tetranychidae) Araginee rong due cotonier a Madegascer . Cotton Fibers Trop. 22(2): 183-195.

REFERENCES

- Halawa, A. M. (1998):** Studies on some mites associated with fruit trees.
M.sc. Thesis Fac. Agric. Moshtohor. Zagazig Univ. 113pp.
- Harrison, S. and R. Karban (1986):** Behavioral response of spider mite (*Tetranychus urticae*) to indicated resistance of cotton plants. Ecol. Entomol. 11(2): 181-188.
- Hassan, R. H. M. (2003):** Studies on fungal pathogens for controlling the two spotted spider mite, *Tetranychus urticae* Koch. M.sc. Thesis Fac. Agric. Cairo Univ. 70pp.
- Hassan, A. S.; El-Nahal, A. K. and E. A. El-Badoy (1959):** Infestation of cotton with spider mites found on cotton plants in Egypt and their predators. Bull. Soc. Entomol. Egypt, (43): 357-362.
- Hell, W. and M. W. Sabelis (1985):** Spider mites, their biology, Natural enemies and control. Elsevier. Amestrdam. I (B): 458pp.
- Henderson, C. F. and E. W. Tilton (1955):** Test with acaricides against the brown wheat mite. J. Econ. Entomol. (48): 157-161.
- Hessein, N. A. and T. M. Perring (1986):** Feeding habits of the Tydeidae with evidence of *Homeopronematus anconai* (Acari: Tydeidae) Perdition on *Acylops lycopersici* (Acari: Eriophidae). Int. J. Acarol. (12): 215-221.
- Hightower, B. G. and D. F. Martin (1956):** Ecological studies of spider mites attacking cotton in central Taxes. J. Econ. Entomol. 49(3): 424-425.
- Hinz, S. E. and J. E. Wright (1997)** Naturalist- L: a biological product (*Beauveria bassiana* J W-1) for the control of cotton pests.

Proceed. Beltw. Cotton. Confer. new orleans, LA, USA.
(2): 1300-1302.
- Hossain, A. M. (1981):** Effect of cultural practices, on mites inhabiting cotton plants. M. Sc. Thesis, Fac. Agric. Ain Shams. Univ. Egypt. 100pp.
- Karban, R. (1987):** Induced resistance against spider mite in cottons field verification. Rev. Appl. Entomol. 75(7): 578.

REFERENCES-----

- Krantz, G. W. (1978):** A manual of Acarology, Oregon state Univ. Book Stores. Inc. Corvallis. 509pp.
- Kuhnelt, W. (1950):** Soil biology with special reference to the animal Kingdom. Faber & Faber., London. 397pp.
- Lindquist, E. E. and G. O. Evans (1965).** Taxonomic concepts in the Ascidae, with a modification in setal nomenclature for the Idiosoma of the Gamasina (Acarina: mesostigmata). Mem. Entomol. Soc. Can. No. 47.
- Lockley, T. C.; Smith, J. W.; Scott, W. P. and C. R. Parencia (1979):** Population fluctuation of two groups of spiders from selected cotton fields in panola and pontotoc counties, Mississippi. The Southeastern. Entomol. 4(1): 20-24.
- Mansour, F. (1986):** Spiders in sprayed and unsprayed cotton fields in Israel, Their interactions with cotton pests and their importance as Predators of the Egyptian cotton leafworm, *Spodoptera littoralis*. Phytoparasitica. 15 (1): 31-41.
- Mansour , F.; Bernstein, E. and F. Moch (1995):** The potential of spiders of different taxa and a predacious mite to feed on the carmine spider mite. Phytoparsitica. 23(3):217-221.
- Michel, B. and P.Prudent (1985):** Mite and insect pests of cotton (*Gossipum hersutum* L.) in Paraguay. Cotton Fibre- Trobicals. 40 (4): 219-230.
- Mohafez, M. A. M. (2000):** Studies on true spiders in Sohag Governorate. M. Sc. Thesis, Fac. Agric. Al-Azher Univ. 115pp.
- Mostafa, S. A. (1988):** Effect of some insect growth regulators on some cotton pests. M. Sc. Thesis, Fac. Agric., Al- Azhar Univ. Egypt. 100pp.
- Muller, k. E. (1965):** Pilzkrankheiten bei Insekten. A nwendung zur biologischen Schad ling sbekamp fung und Grundlagen der Insektenmykologie. Paul parey, Berlin and Hamburg.
- Nassef, M. A.; El- Feshawi, A. A.; Abo- Sholooa, M. K. A. and W. M. Waston (1996):** Seasonal abundance of certain predator as influenced by cotton planting dates and weather factors. Egypt, J. Agric. Res., 74(3): 611- 625.

REFERNCES-----

- Nyffeler, M.; Dean, D. A. and W.L. Sterling (1987):** Evaluation of the importance of the striped Lynx spider, *Oxyopes salticus* (Araneae: Oxyopidae), as a predator in Texas cotton. Environ. Entomol. 16 (5): 1114-1123.
- Omar, H. I. H.; Hegab, M. F. and A. E. M. El-Sorady (1993):** The impact of intercropping cotton and cowpea on pest infestation. Egypt. J. Agric. Res. 71 (3): 709-716.
- Omar, H. I. H.; El-Bessony, M. A. E.; El-Magraby, H. M. and M. H. M. El-Khawalka (2000):** Effect of abamection compared with different compound on the two spotted spider mite and two insect leaf miners. J. Agric. Sci. Mansoura Univ. 25(12): 8089-8096.
- Omar, B. A. and H. M. El-Khateeb (2002):** Efficacy of some biopesticides against *Tetranychus urticae* infesting cowpea plants and their side-effects on certain predators. Egypt. J. Agric. Res. 80(3): 1157-1171.
- Pellagatti, O.; Frate, G. D.; Caretta, G. and G. Delfrate (1989):** Fungi isolated from insect and mites. Redia. (71): 255-266.
- Plagens, M. J. (1983):** Population of *Misumenops* (Araneida: thomicidae) in two Arizona cotton fields. Environ. Entomol. (12): 572-575.
- Rahil, A. A. H. (1988):** Ecological and biological studies on the spider at Fayoum. M. Sc. Thesis, Fac. Agric. Cairo Univ. 133pp.
- Sallam, Gihan, M. (2002):** Studies on true spiders in Egypt. Ph.D. Thesis, Fac. of Agric. Cairo Univ. 131pp.
- Sallam - Gihan, M. and H. K. El-Hennawy (2003):** Biological aspects of *Nurscia albomaculata* (Lucas, 1848) (Arachnida: Araneida: Titanoecidae) in Egypt. Serket. 8(4): 147-150.
- Sedrak, R. A. (1978):** Ecological studies on citrus brown mite. M. Sc. Thesis, Fac. Agric. Ain Shams Univ. Egypt. 88pp.
- Schoenig, S. E. and L. T. Wilson (1992):** Patterns of spatial association between spider mites (Acari: Tetranychidae) and their natural enemies on cotton. Environ.- entomol. 21(3): 471-477.

REFERENCES-----

- Sewify, G. H; Aboul-Ela, S. and M. S. El-Oin (2000):** Effects of the entomopathogenic fungus *Metarhizium anisopliae* (Metsch.) and granulosus virus (GV) combinations on the potato tuber moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). Bull. Fac. Agric. Cairo Univ. (51): 95-106.
- Shi, X. and Z. A. Wu (1987):** A preliminary evaluation of a space distribution model for *Tetranychus. urticae* and sampling quality in cotton field. Rev. Appl. Entomol. 75(6): 530.
- Simittly, D. R. and G. G. Kennedy (1988):** Aerial dispersal of the two-spotted spider mite (*Tetranychus urticae*) from field corn. Exp. and Appli. Acarolo. 5(1-2): 33 - 46.
- Steel, R. G. and J. H. Torrie (1968):** Principles and procedures of statistics. McGraw Hill Book Co. Inc. New York.
- Taha, H. A.; Sawires, Z. R. and A. M. Mostafa (1990):** Population dynamics of the two- spotted spider mite, *T. arabicus* Attiah on two cotton varieties. J. Agric. Res. Egypt. (68): 5-10.
- Taha, H. A. and S.A. El-Raies (1996):** Influence of water salinity irrigation on spider mite infestation and cotton leaves contents. Egypt. J. Agric. Res. 74 (1): 83 - 90.
- Taha, H. A.; Soliman, S. M.; Abd-El-Haleem, A. and S. A. El-Raies (2001):** Field studies on the main pests infesting cotton with refer to their natural enemies'. Proceedings of Beltwide cotton conference U. S. A. (2): 885-888.
- Taha, H. A.; Ahmed, M. A. and A. A. Younes (2002):** Induced resistance to spider mite *T. urticae* (Koch) infestation of cotton and soybean plants. J. Agric. Sci. Mansoura Univ. 27 (1): 625-630.
- Tamai, M. A.; Alves, S. B.; Lopes, R. B. And P. S. Neves (1998):** Avaliacao de fungos entomopatogenicos para control de *Tetranychus urticae* Koch. Abstracts 17th Brazil Congr. Entomol. Riode Janeiro. 1066pp.
- Tamai, M. A.; Avies, S. B. and P. j. Neves (1999):** Pathogenicity of *Beauveria bassiana* (Bals.) Vuill. against *Tetranicus* Koch. Sci.-Agric. 56 (2): 285-288.

REFERNCES-----

- Trichile, P. J. and T. F. Leight (1985):** The use of life tables to asses varietals resistance of cotton to spider mite. Entomol. Exp. Appl. 39 (1): 27-33.
- Thulasi, R. K.; Patil, B. V. and M. Bheemanna (1994):** Survey of the spider mites species and their natural enemies on cotton in Tungabhadra project area. Karnat.-J. Agric. Sci. 7(2): 240-242.
- Wahba, M. L. (1976).** Studies on tydeid mites (Family: Tydeidae), especially as related to mango and citrus trees. Ph. D. Thesis, Fac. Sci. Ain Shams Univ. Egypt. 119pp.
- Wehner, F. (1990):** The effect of time and duration of infestation by *Tetranychus turkestanii* on plant growth, yield and quality of cotton in Central Asia. Beitrage Zur Tropisches Landwirt. Veter. Remed. 27 (2): 203- 209.
- Wilson, L. T.; Trichilo, P. J. and D. Gonzalez (1991):** Spider mite (Acari: Tetranychidae) infestation rate and initiation effect on cotton yield. J. Econ. Entomol, 84(92): 593-600.
- Yassin, E. M. A. (1997):** Biological and ecological studies on some phytophagous mites associated with some field crops. M. Sc. Thesis, Fac. Sci. Cairo Univ. 155pp.
- Yassin, E. M. (2004)** studies on some Tydeid mites in Egypt. Ph. D. Thesis, Fac. Sci Cairo Univ., 319pp.
- Yousri, H. (1994):** Laboratory evaluation of *Beauveria bassiana* (Balsamo) Vuill. As pathogen of adult female stage of the two-spotted spider mite *Tetranychus urticae* Koch. Egypt. J. Appl. Sci. 9(2): 380-387
- Yousef, A. A.; El-Badry, E. A. and I. H. Heykel (1976):** Mites inhabiting cotton and associated weeds in Egypt. Bull. Soc. Entomol. Egypt. (60): 223-227.
- Zaher, M. A. and E. A. El-Badry (1962):** Abundance of mites and insects predaceous on tetranychid and tenupalpid mites in Giza. Bull. Soc. Entomol., Egypt. (46): 429-441.
- Zaher, M. A. (1986):** Survey and ecological studies on phytophagous, predaceous and soil mites in Egypt. (Nile Valley and Delta). II. Predaceous and non phytophagous mites PL. 480 program U.S. A. Project No. EG-ARS- 30 Grant No. FG- 139:567pp.
- REFERENCES-----**