

REFERENCES

- Abdel-Moneim, A. M. (1976):
Eosinophilia in Schistosomiasis, M. D. Thesis, Cairo University.
- Abdel-Salam, M. and Abdel-Fattah, S (1977):
Prevalence and morbidity of *S. haematobium* in Egyptian children, Am. J. Trop. Med. Hyg. 26(3): 463-469.
- Abdel-Wahab, M.F., Strickland, C. T., El. Sohly, A., El-Kady, N. Zakaria, S. and Ahmad, L. (1979):
Changing pattern of schistosomiasis in Egypt: 1935-1979. lancet 2:242.
- Abdel-Wahab, M.F., Strickland, C. T., El-Sahly, A., Ahmed, L., Zakaria, S., El-Kady, N. and Mahmoud, S. (1980):
Schistosoma mansoni in an Egyptian village in the Nile delta. Am. J. Trop. Med. Hyg 29:868.
- Aidaros, S. M., Soliman, A. and Lily, A. M. (1961):
Partial vascular changes in human bilharzial cirrroses. J. Path. and Bact. 82:91.
- Al-Alamy, M.A. and Clin B.L. (1977):
Prevalence and intensity of *Schistosoma haematobium* and *Schistosoma mansoni* infection in Qolyub, Egypt. Am. J. Trop. Med. Hyg. 26-470.
- Archer, R. K. (1959):
Eosinophil leucocyte and their reaction to histamine. J. of Pathology and Bacteriology 78: 95-100.

Armand-Delile (1930):

Quoted from Britton p.90.

Arnoldsson, H. and Helander, E. (1958):

Acta.Allerg. 12:96 Quoted from Britton p:90.

Atta, A. A. (1960):

Tropical Medicine and Infectious Disease Text Book, first Edition. Modern Cairo Book shop. p. 367.

Ayad. N. (1974):

A short review of epidemiology of schistosomiasis in Africa. Egypt. J. Bilharz. 1,9.

Baehner, R.L and Johnston, R. B. (1971):

Metabolic and bactericidal activity of human eosinophil. British Journal of Haematology 20,277.

Baston, A.and Besson, A. (1970):

Mechanism of eosinophil. Role of the lymphocyte. J. Exp. Med. 131: 1288-1305.

Bell, A. (1963):

A new method in counting *S. mansoni* eggs in faeces. Bull. World Health Org. 29, 525.

Bell, A. (1969):

Ann. of New York Academy of Science 160: 593-601.

Berkow, R. (1982):

In, the Merck Manual of Disease and Therapy, (Robert Berkow, (chief Editor) Pub. Merck Sharp and Dohme Research Laboratory p. 323, 1213.

Bogitsh, B. J. (1971):

Tissue eosinophilia in infested rats. *Exp. Parasitol*
292; 493.

Britton, G. J. C. (1963):

Eosinophilia. In: "Whitby and Britton", *Disorder of the blood.*
Ninth Edition. Pub. J. and A. Churchill L. T. D, New York,
4: 89-92.

Butterworth, A.E (1979):

Damage to schistosomules of *S. mansoni* induced directly by
eosinophilic major basic protein. *J. Immunol.* 122(1):
221-229.

Byram, J.E. (1977):

Altered schistosoma granuloma formation in nude mice. *Am. J.*
Trop. Med. Hyg. 26, 944-956.

Byram, J. E. (1979):

S. mansoni infection in T-cell deprived mice. *Am. J. Trop.*
Med. Hyg. 28(2): 224.

Cheever, A. W. (1968):

A quantitative post-mortem study of *Schistosomiasis mansoni*
in man. *Am. J. Trop. Med. Hyg.* 17, 38-64.

Cheever, A. W., Kamel, I.A, Elwi, A.M, Mosimann, J. E., and
Danner, R. (1977):

Schistosoma mansoni and *Schistosoma haematobium* in Egypt.
II. Quantitative parasitological findings at necropsy.
Am. J. Trop. Med. Hyg. 26(4), 702-716.

Cheever, A. W., Kamel, I.A., Elwi, A.M., Mosiman, J.E., Danner, R. and Sippel J. E. (1978):

S. mansoni and S. heamatobium in Egypt.III extrahepatic pathology. Am. J. Trop. Med. Hyg. 27: 55-75.

Cohen, S. G and Sapp, T. H (1965):

Experimental eosinophilia. J. allergy 36: 45-422.

Cohen, S. G; Vassalli, B., Benacerrof, B. and McClusky R.T. (1966):

The distribution of antigenic and non antigenic compound in dranng lymph node. Laboratory Investigation, 15: 1143-1155.

Cohen, S. G and Word, P. A. (1971):

In vitro and in vivo activity of a lymphocyte and immune complex-dependent chemotactic for eosinophil. J. Exp. Med. 33: 133-140.

Colley, D. G. (1972):

S. mansoni: eosinophilia and the develepement of lymphocyte blastogenesis in response to soluble antigen in mice. Exp. Parositol. 32, 520-526.

Colley, D. G. (1977):

Immune response during human schistosomiasis mansoni. Am. J. Trop. Med. Hyg. 26:917.

Crofton, J. W., Lininestone, J. L., Oswald, N. C. and Reberts A., T. (1952):

Pulmonary eosinophilia. Thorax 7:1.

- Dacie, J. V. and Lewis, S. M. (1984):
Basic haematological technique. In "Practical haematology"
6th edition. Pub. Churchill Livingstone, London 3:22.
- Discombe G. (1946):
Lancet 1:195 Quoted from. Penington P. 400
- Elaine, C. J., Emil, y. c. and Seymour; J. (1984):
Human neutrophil-mediated Killing of schistosomula of
S. mansoni. Am. J. Trop. Med. Hyg. 33(1): 104.
- Elwi, A. M. (1967):
In Bilharziasis, 1st Ed. (Eds) F.K. Mostafi, Springer-Ver-
alog, New York p. 39.
- Engleman, E. P and Shearn, M. A. (1983):
Artheritis and allied rhumatic disorders—progressive sys-
temic sclerosis. In. Current Medical Diagnosis and Treat-
ment, Marcus. A. Krupp, Chief. Editor). Pub. Lange
Medical Publication, California 13: 493.
- Farid, Z. (1984):
Schistosomiasis. Medicine International. J. Middle Esteran
Edition 193-199
- Farooq, M. and Samman J. A. (1967):
The relative potential of different age group in the trans-
mission of Schistosomiosis in Egypt., 49 project area. Ann.
Trop. Med. Parasit. 61(3): 315-320.
- Feldmeier, H., Doehring, E. and Dafalla, A., A. (1982):
Simultaneous use of a sensitive filtration technique and
reagent strips of urinary schistosomiasis. Trans. Roy.
Soc. Trop. Med. Hyg. 76: 416-421.

Fine, D. P. (1973):

Schistosoma mansoni infection in mice depleted of thymus-
-dependant lymphocyte. Am. J. Path. 71, 193.

Gonnert, R., (1955):

Studies on schistosomiasis. Trop. Med. Parasit. 6:279.

Hamilton, P. K., Hutchinson, H. S., Jansion, P. W. and Jones,
H. L. (1959):

The pathology and pathogenesis of hepatosplenic. Disease
associated with bilharziasis, Egyptian bilharziasis. Am.
J. Clin. Path. 32-18.

Hamman, H. M., Fapha, P. H. and Allan, F. M. (1976):

Epidemiology of schistosomiasis. Am.J. Trop. Med. Hyg 3:10.

Hartnett, B.S. (1955):

Eosinophilia after administration of some drug. Brit. Med.
J. 1: 1458.

Hosinger, J.(1972):

The eosinophil and allergy: Why ? J. allergy and Clin.
Immunol. 94(3): 142

Hsu , S.Y.L, Hsu, H. F., Mitors, F. A, Helms, G. M. and Solo-
mon, R.I (1980):

Eosinophils as effector cell in the destruction of S. mansoni
eggs in granulomas. Ann. Trop. Med. Parasitol. 74, 179-183.

James, S. L. and Colley, D. G. (1975):

Eosinophil and immune mechanism, production of the lymphokine
eosinophil stimulation promoter in vitro by isolated intact
granuloma. J. Reticuloendothelial Soc. 18, 283-293.

- James, S.L. and Colley, D. G. (1978):
Eosinophil mediated destruction of *S. mansoni* eggs in Vitro
II. The rol of eytophilic antibody. *Cell Immunol* 38, 35-47.
- Jardan, S., Weingatan, M. and Staven, F. (1985):
Eosinophilic meningitis and hypereosinophilic syndrom. *Am. J. Med.* 78: 674.
- Kabil, S. M, Hifney, A.A and Youssef, S. M. (1986):
The value of rectal snip in the diagnosis of schistosomiasis
mansoni in Upper Egypt (Sohag Province). *Benha Medical Journal*, Accepted 1986.
- Kamil, I. A., Smith, J. H., Elwi, A. and Vonlichtenberg, F (1974):
A quantitative post-mortam analysis of urinary schistosomiasis
in Egypt. I pathology and pathogenesis. *Am. J. Trop. Med. Hyg* 23:1054.
- Kamel, I. A., Elwi, A. M., Cheever, A. W., Mosiman, J. E. and Danner. R. (1978):
S.mansoni and *S. haematobium* in Egypt IV hepatic lesion
Am. J. Trop. Med. Hyg. 27:931.
- Kato, K. and Miura, M. (1954):
Comparative examinations. *Jap. J. Parasitol*, 3: 35. Quoted
from (Martine and Beaver, 1968).
- Kloetzel, K. (1963):
Some quantitative aspects of diagnosis and epidemiology in
S. Mansoni. *Am. J. Trop. Med. Hyg.* 12: 337-339.

- King, M. A. (1976):
"A Medical Laboratory for Developing Countries". 1st edition. Pub, The English Language Book Society and Oxford. University Press 8:15, 10:2.
- Knopf, P. M. (1979):
S. mansoni: Peripheral and tissue eosinophilia in infected rats. Exp. Parasitol 47(2), 232-244.
- Kuberski, T. and Wallace, G. D. (1979):
Clinical manifestation of eosinophilic meningitis due to Angiostrongylus cantonensis Neurology 29: 1566.
- Lani, S., S., Micheal, C. L., Kathleen, M. K. and Dennis, M. (1985):
Urinary iron loss and physical fitness of Kenyan children with urinary schistosomiasis. Am. J. Trop. Med. Hyg. 34(2), 322-330.
- Lehman, J. S., Farid, Z., Smith, J. H and El-Masry, N.A. (1973):
Urinary schistosomiasis in Egypt. Clinical, radiological and parasitological correlation. Tran. Roy. Soc. Trop. Med. Hyg. 67(3): 384-399.
- Lehman, J. S., Mott, K. E., Muniz, T. M. and Boyer, M. H. (1976):
The intensity and effects of infection with S. mansoni in rural community in northeast Brazil. Am. J. Trop. Med. Hyg. 25: 285-294.
- Lewis, F. A., Capter, G. E. and Colley, D. G. (1977):
Eosinophil and immune mechanism. Cell. Immunol 32, 86-96.

- Litt, M. (1964):
Eosinophilia and antigen-antibody reaction. *Annals of the New York academy of science* 116, 964-989.
- Mahmoud, A. A. F. (1975):
A role for eosinophils in acquired resistance to *S. mansoni* infection. *J. Exp. Med.* 142: 805-813.
- Manson-Bahr, B. E. C. and Apted, F. I. C. (1983):
Schistosomiasis. *Manson's Tropical Diseases*. Pub. The English Language Book Society and Bialliere Tindall. London p. 206.
- Martin, L. K. (1965):
Randomness of particle distribution in human faeces and the resulting influence on helminth eggs counting. *Am. J. Trop. Med. Hyg.* 14: 747, 759.
- Martine L. K., and Beaver (1968):
Evaluation of Kato-thick smear technique for quantitative diagnosis of helminth infection. *Am. J. Trop. Med. Hyg.* 17, 382-391.
- Masaba, S. C., Awiti, I. E. and Muruka, J. F. (1983):
Morbidity of urinary schistosomiasis in relation to the intensity of infection via Kisumu, Kenya. *J. Trop. Med. Hyg.* 86(2), 65-66.
- McMahon, J. E (1976):
Egg excretion of *Schistosoma haematobium*. *East. Afr. Med. J.* 44:250.

Mousa, A. H. and El-Garem, A. S. (1964):

The haemodynamic study of Egypt hepatosplenic bitharziasis.
J. Egy. Med. Ass. 42:444.

Mousa, A. H. (1976):

Pathological lesion associated with *S. mansoni* and *S. haematobium* infection in man. Trans. R. Soc. Trop. Med. Hyg. 61: 620.

Milburn, C. L. and Ernst, K. P (1953):

Eosinophilia-hepatomegaly syndrome of infant and young children-report of a case due to invasion of liver by nematode larva. Pediatrics 11: 358-367, Quoted from Thompson p. 461.

Muehrcke, R. C., Eckert, E. L and Kark, R.C (1952):

J. Lab. Clin. Med, 40: 161 Quoted from, Penington P, 401.

Olds, G. R. and Mahmoud, A., A., F. (1980):

Role of host granulomatous response in murine schistosomiasis *mansoni*. Eosinophil-mediated destruction of eggs. J. Clin. Invest. 66, 1191-1199.

Olds, G. R. and Mahmoud, A. A. F. (1981):

Kinetics and mechanisms of pulmonary granuloma formation around *S. Japonicum* eggs injected into mice. Cell Immunol 60, 251-260.

Onori, (1962):

The diurnal cycle of egg excretion of *Schistosoma haematobium*
Ann. Trop. Med. Parasit. 56: 292.

Ottson, E. A., Stonley, A. M., Gifand, A. G. and Godek, J. E.
(1977):

Immunoglobulin and complement receptors on human eosinophils
and their role in cellular adherence to schistosomiasis.
Am. J. Trop. Med. Hyg. 26; 134-141.

Owhashi, M., Horii, Y. and Ishii, A. (1983):

Eosinophil chemotactic factor in schistosome eggs. Am. J.
Trop. Med. Hyg. 32(2): 359-366.

Pathak, G. L and Kahali, B. S (1975):

Cyclic Variation in the eosinophil count during the phases
of the menstrual cycle. J. of clinical endocrinology
17, 862-869.

Pelley, R. P. (1977):

Splitt suppressor in chronic schistosomiasis mansoni.
Suppression of cell mediated immunity and enhancement of
humoral immunity. Clin. Res. 25: 491 A.

Penington, D., Rush, B., and Castoldi, P. (1983):

Eosinophilia In "G. C. deGrachy", Clinical hematology in
medical practice 4 th edition. Pub. Blackwell Scientific
Publications, Oxford P: 400-402, 603.

Pepper, H. and Lindsay, S. (1960):

Level of eosinophil, Platelets, leucocyte and 17 hydroxy
Corticosteroid during menstrual cycle. Pro. Soc. Exp.
Biol. Med. 104, 145-147.

Peters, P. A., Mahmoud, A. F. Warren, K. S. Ouma, J.H. and Arapsiongok, T. K. (1976).
Field studies of a rapid accurate means of quantifying schistosoma Haematobium eggs in urine samples. Bull. Wld. Health. Org. 54-59.

Phillips, M. (1977):

Schistosomiasis in the congenitally athymic (nude) mice.
I. Thymic dependency of eosinophilia, granuloma formation and host morbidity. J. Immunol. 118(2): 594-599.

Pope, Rt., Clin, B. L. and El-Alamy, M. A (1980):

Evaluation of schistosomal morbidity in subject with high intensity infection in Qalyub, Egypt. Am. J. Trop. Med. Hyg. 29, 416.

Pugh, R. N. H.; Bell, D. R. and Gilles, H. M. (1980):

The potential medical importance of bilharzia in Northern Nigeria. Ann. Trop. Med. Parasit. 74: 597-613.

Punyagupta, S. Jutti, P. and Bunnagt. (1975):

Eosinophilic meningitis in Thailand, clinical studies of 484 typical cases probably caused by *Angiostrongylus cantonensis*. Am. J. Trop. Med. Hyg. 24:921.

Rescetti, M. (1978):

Specific release of neutrophilic and eosinophilic stimulatory factors from sensitised lymphocytes. Blood 47: 757-765.

Riddle, J. M. and Barhort, M. I. (1965):

The eosinophil as a source of profibrinolysin in acute inflammation Blood 25, 776.

Riisager, P. M. (1959):

Study on blood cell in G. I. T. disease Lancet 2: 1008.

Ritchie, L. S., Knight, W. B., Hiatt, R. A. and Cline, B. L.
(1976):

A modification of the formol-ether concentration technique for increased sensitivity in detecting schistosoma mansoni eggs. Am. J. Trop. Med. Hyg. 25, 818-823.

Saif, M. (1961):

A simple method for counting bilharzialova in urine. J. Egypt. Med. Assoc. 44:1016.

Salih, S. Y. (1977):

Eosinophilia in S. mansoni infection. East. Afr. Med. J. 54(8): 421-424.

Scott, J. A. (1968):

The regularity of egg out put of helminth infections with special reference to schistosoma mansoni. Am. J. Trop. Med. Hyg. 27: 155-175.

Sher, A. (1977):

Immunity against S. mansoni in the mouse. Am. J. Trop. Med. Hyg. 26: 20-28.

Shrewsbury, M. M. and Reinhardt, W. O. (1955):

Relationships of adrenals, gonads and thyroid to thymus, lymphnode and blood leucocytes. Blood 10: 633-645.

Smith, J. H., Kamal, I. A., Elwi, A. and Von-Lichtenberg, F,
(1974):

A quantitative post mortem analysis of urinary schistosomiasis in Egypt. I- Pathology and pathogenesis. Am. J. Trop. Med. Hyg. 23(4): 1054-1071.

Stewart, S. G (1933):

Am. J. Med Sci, Quoted from Britton P. 91.

Stimmel, G. M. and Scott, J. A. (1969):

The regularity of egg out-put of *Schistosoma haematobium*. Texas. Rep. Biol, Med. 14: 440-458.

Stoll, N. R. (1926):

Concerning two options in dilution egg counting, small drop and displacement. Am. J. Trop. Med. Hyg 6: 134-145
Quoted from Martin (1965).

Sturrock, R. F. (1977):

Eosinophilia in kenyan baboons induced by *T. spiralis* and *S. mansoni* Exp. Parasitol 75: 89-100.

Tai, P. C and Spry C. J. F. (1967):

Studies on blood eosinophil. Clin. Exp. Immunol. 24: 415-423.

Tessdale, C. H. and Amin, M. A. (1976):

Comparison of the Bell technique, a modified Kato-thick Smear technique and a digestion method for the field diagnosis of schistosomiasis mansoni. J. of Helminth. 50: 17-20.

Thompson, R. B. (1977):

Eosinophilia, IN "Disorders of the Blood", A text book of clinical haematology Pub. Churchill Livingstone, New York 29. p.460-464.

Tsuda, S., Fukuyama, K. and Epstein, W. L. (1979):

Low molecular weight eosinophil chemotactic factor in granulomatous liver of murine schistosomiasis. J. Immunol., 122, 2554-2557.

Wadee, A. A. and Sher, R. (1980):

The effects of a soluble factor released by sensitized mononuclear cells incubated with *S. haematobium* ova on eosinophil migration. J. Immunol. 14. 989-995.

Walker, A. R. P. (1968):

A simple method for large scale collection and preservation of urins for counting bilharzia eggs. Trans. Roy. Soc. Trop. Med. Hyg. 62: 570.

Wallerstein, R. O. (1983):

Histiocytosis in Current Medical Diagnosis and Treatment. (Marcus A. Krupp, (Chief Editor). Pub. Lange Medical Publication, California 9:320.

Ward, P. A. (1969):

Chemotaxis of human eosinophil. Am. J. Pathology 54: 121-128.

Warren, K. S. (1961):

The aetiology of hepatosplenic schistosomiasis mansoni in mice. Am. J. Trop. Med. and Hyg. 10, 870.

Warren K. S. Mohmoud A. F., Muruka J. F., Whittakesl, R. and Singok T.K. (1979):

Relationship between egg out put and morbidity. Am. J. Trop. Med. Hyg. 28(5): 864-840.

Wasserman, S. L., Goetzl, E. J. and Austen, K. F. (1975):

Inactivation of slow reacting substance of anaphylaxis by human arylsulfatase. J. Immunol 114.

WHO (1980):

World Health Organisation. Technical Report Series 643: 1-63.

Wilkins, H. A., Gool, P., and Moore, P. J. (1979):

The significance of proteinuria and haematuria in Schistosoma haematobium infection. Trans. Roy. Soc. Trop. Med. Hyg. 73: 74-80.

Wilkins, H. A., Goll, P. H. and Moore, P. J. (1985):

Schistosoma haematobium infection and haemoglobin concentration in a Gambian community. Ann. Trop. Med. Parasitol. 79(2): 159-161.

William, W. J., Beutlar, E., Erslev, A. J. and Rundles, W., R. (1977):

In, Heamatology, Text book. Puf., Magrow-Hill Book Company. A Blakiston Publication. New York p. 659, 668, 670.

Wintrobe, M. M., Richard, G. L, BoGGs, D. R. and Foerster, J. (1975):

Causes of eosinophilia. In "Clinical Heomotology" Seventh Edition. Pub. Henry Kimpton, London 3:262, 41:1282.

Woodruff, A. W. (1970):

Toxocariasis. British. Med. J. iii: 663-669.

Yam, L. T. and Crosby, W. H. (1971):

Cyto-chemical identification of monocyte and granulocyte
Am. J. Clin. Pathology 55, 283.

Zuelser, W. W. and Apt. L (1949):

Disseminated visceral lesion associated with extreme eos-
inophilia. Am. J. disease of children. Quoted from,
Thompson. p. 481.