

REFERENCES

A

Abman S H, Accurso FG, Wikenning, et al ;(1987):

Persistent fetal pulmonary hypertension after acute hypoxia
Am J. Physiol .,253: H 941-H 948.

Acquatella, H; Schiller, NB; Sharpe, DN. et al, (1979):

Lack of correlation between echocardiographic patterns of
pulmonary valve morphology and simultaneous pulmonary
arterial pressure.
Am. J. Cardiol 43 . 946.

Allen, HD; Sahn, Dj; Goldberg Sj, et al., (1989) :

Doppler echocardiography. An overview,
Indian pediatr. 19 : 79.

Ansert, H. S., (1983) :

Textbook of diagnostic ultrasonography, 2nd ed st. Louis :
CV Mosby.

B

Baker, D.W., Rubenstein S.A., and Lorch D.S., (1977) :

Pulsed Doppler echocardiography : Principles and
applications. Am J. Med. : 80, 63.

Barst R.J., (1986):

Pharmacologically induced pulmonary vasodilation in children and young adults with pulmonary hypertension .
Chest. 89 : 497 - 503.

Beitzke A, Suppan C, Justich E., (1982) :

Complications in 1000 cardiac catheter examinations in childhood. Rontgen - blatter 35 : 430 - 437 .

Berger AM, Popp RL, (1985):

Estimation of right ventricular systolic pressure by Doppler echocardiography in patient with tricuspid regurgitation.
Circulation 70:657-662,.

Bommer W.J, (1985) :

Basic principles of flow imaging echocardiography
2 : 507 - 509 .

Braunwald N.S., Ross, J.J., Morrow, A.G. (1980) :

Conservative management of tricuspid regurgitation in patients undergoing mitral valve replacement.
Circulation, 35 (Suppl I) : 1-63.

Brecker S.J., Gibbs - J. S., Fox - K.M. et al., (1994):

Comparison of Doppler derived haemodynamic variables and simultaneous high fidelity pressure measurements in severe pulmonary hypertension .
Br - heart J. 72 (4) : 384 - 9.

C

Caceres J, Santos D.J., Gavilan J.L, et al; (1995) :

Doppler echocardiographic evaluation of pulmonary hypertension in children .

Rev.Esp. Cariol , 48 (2) :122-127 .

Carins. K.B., Kloster, F.F., Bristow, J.D., et al; (1968) :

Problems in the haemodynamic diagnosis of tricuspid insufficiency.

Am. Heart J. 75 : 173 - 9.

Celemajer , D.S., Dollerg, C., Burch, et al., (1994):

Role of endothelium in development of pulmonary hypertension .

Circulation, 89 (5) :2041 - 4.

Craig, M., (1991) :

Echocardiography instruments And principles.
Echocardiography textbook volume II. Edited by Marveen Carig, international ultrasound institute, Dallas, Texas, J. B, Lppincott company, Philadelphia, London page 43 .

Currie PJ, Seward JB, Chan KL, Fyfe DA, et al; (1985) :

Continuous Wave Doppler determination of right ventricular pressure : A simultaneous Doppler catheterization study in 127 Patients.

Cullinane C., Cox P.N. Silver M.M., (1992):

Persistent pulmonary hypertension of the newborn due to alveolar capillary dysplasia. *Pediatr Pathol* ; 12 : 499 - 514.

D

Das, M. Sternmank KR, Dempsey E.C., (1995) :

Enhanced growth of fetal and neonatal pulmonary artery adventitial fibroblasts is dependant on protein Kinase C., *Am J physiol* 269 :L 660 - L 667 .

Diebold, B., Touat, R., Blanchard, D., Colonna, G., J.C., et al; (1983) :

Quantitative assesment of tricuspid regurgitation using pulsed Doppler echocardiography .
Br. Heart J. 1983, 50 443 - 9.

Diseassa ; T.G., Hagen, A.D, and Isablejones, J.B., (1981) :

Two - Dimensional echocardiographic evaluation of the discrete subaortic stenosis from the apical long axis view.
Am . Heart, J., 101 : 774 - 782.

Durmowicz A G, Parks WC, Hyde DM. Et al (1994):

persistant reexpression ,and induction of pulmonary arterial fibronectin, tropoelastin, and type one procollagen mRNA in neonates with pulmonary hypertension.

Am J Pathol 145: 1411-1420,.

E

Eqqleton, R.C., (1978):

Intesin AIUM standard nomenclature .

Reflections, 4 : 275 .

F

Fantini. F., Magherini, A., (1978) :

L'echo cardiographia Dopplernella diagnosi dei vizi Tricuspidalici.

G. Ital. Cardiol. 8 : 879 .

Fiegenbaum, H., (1994) :

Aquired valvular heart Disease : In echocardiography. 5th ed philadelphia : lead and febigur 286 - 8.

Fiegenbaum, H. (1994):

Echocardiographic examination, Echocardiography 5th ed.

Pheladelphia : lea and febiger. P. 68-125.

Frid, M.G., Moiseeva, E.P., Stemmark, K. R. (1994) :

Multiple phenotypically distinct smooth muscle cell populations exist in adult and developing bovin pulmonary arterial media in vivo;. *circ Res* 75 : 689 - 681.

Fridman, I., Baufer, A., Ron, N. And Davies, A.M. (1997):

Experimental myocarditis : in vitro and in vivo studies of lymphocytes sensitized to heart extracts and group A streptococci.

Immunology 20 225- 232.

Fuster V, steel PM, Edward WD ,et al ;(1984):

primary pulmonary hypertension :Natural history and importance of thrombosis .

circulation 70: 580-587.

Fyler , D.C. (1992) :

Pulmonary hypertention, Nadas Pediatric Cardiology textbook . Mosby Year book INC, Boston Philadiphia,

G

Garcia, Dorado, D., Falzgraf, S., et al., (1982) :

Diagnosis of functional tricuspid rurgurgitation by pulsed wave Doppler echocardiography .

Circulation 66 : 1315.

Geggel RL, Dosor AJ, Fyler Dc, et al, (1984) :

The structural basis of pulmonary hypertension of newborns,
Clinical perinatology 2 : 525-549.

Gleason, W.L., Braunwald, E., (1982) :

Studies on the first derivative of the ventricular pressure
pulse in man.

J. Clin . invest., 41 : 40 - 91 .

Goldberg Walds , Cruz, Feldman, L., and Satin, et al; (1981) :

Range gated ultrasound detection of contrast
echocardiographic microbubbles for cardiac and great
vessels blood flow patterns .

Am. Heart J. 101 : 793 , 1981.

Gramiak, R., Nada, N.C. and Shah, P.M (1972) :

Echocardiographic detection of pulmonary valve.

Radiology 102, 153 - 157.

Greistarfer AAT, Peach Mj, Owens GK, (1988):

Angiotensin II induces hypertrophy ,not hyperplasia ,of
cultured rate aortic smooth muscle cells.

Circulation Resp, 62 :749-756.

Grossman W., (1983) :

Echocardiographic features of pulmonary hypertension .

J. Am. Coll. Cardiol 4 : 343 - 550.

Groves B.M., Turkevich D., Donnellan K., et al., (1988):

Current approach to therapy of primary pulmonary hypertension. Chest 93 (Suppl III) : 1755-1785.

H

Hansing C.E., Rowe G.G., (1972) :

Tricuspid insufficiency. A study of hemodynamics and pathogenesis .

Circulation 45 : 793.

Hammerman, C., Lass N., Strates, E., et al., (1987) :

Prostanoids in neonate with persistent pulmonary hypertension.

J. Periatr. 110 : 470 - 472.

Harrigan P., (1991) :

Two dimensional and Doppler echocardiography. Diagnostic Medical sonography vol II edited by Marveen craig. Lippincott company philadelphia page 91 .

Hatle L., Angelesen B. A., Tromsdal A., (1981) :

Non-invasive estimation of pulmonary artery systolic pressure with Doppler ultrasound.

Br. Heart J., 45 : 157 - 65 .

Hatle, L., Angelesen B. A. (1982) :

Doppler ultrasound in cardiology
philadelphia : Lea Febiger 151 - 5.

Hatle L., and Angelsen B., (1985) :

Doppler ultrasound in cardiology. 2nd ed. Philadelphia, Lea,
Febiger page 216 - 220 .

Hawarth G., (1984) :

Pulmonary vascular disease in different types of congenital
heart disease. Implication for interpretation of lung biopsy
finding in early childhood.

Br. Heart J. 52 : 537-571.

Hawarth S.G., (1987) :

Pulmonary vascular disease in ventricular septal defect :
structural and functional correlations in lung biopsies from
85 patients, with outcome of intracardiac repair .

J. Pathol 152 - 168 .

Hawarth S.G., Hislop A., (1983) :

Pulmonary vascular development : normal value of
peripheral vascular structure .

Am J. Cardiol . 52 : 578 - 583.

Health D., Smith P., Gosney J. Et al., (1987):

The pathology of early and late stages of primary pulmonary hypertension.

Br. Heart J. 58 204 - 213.

Holen J. A., aslid R., Landmaek K., (1977) :

Determination of effective orifice area in mitral stenosis from non invasive ultrasound Doppler data and mitral flow rate.

Acta Med. Scand 210 - 83 .

J

James E., Lock, M.D., John, F. et al., (1992) :

Cardiac catheterization , Nadas Pediatric cardiology,

Mosby - year book inc. St. Louis , Boston Page 187 - 220 .

Johnes, C.J., Song. G.J. and Gibson D.G., (1991) :

An echocardiographic assessment of atrial mechanical behavior .

Br. Heart J., 65 : 31.

Jones, P.C., Rabinovich, M., (1995):

Induction of tensascin and endothelial cell apoptosis are features of monocrotaline - induced progressive pulmonary vascular disease .

Circulation , 92 : 1 - 74 .

Jones, C. J., Song. G. J. And Gibson, D. G., (1991) :

An echocardiographic assessment of atrial mechanical behavior.

Br. Heart. J., 65 : 31 .

K

Kaku R., Neumann, A., Bommer W., Weinert L., Mason D.T. and Demaria, A.N., (1978) :

Sensitivity and specificity of the pulmonic valve echogram in the detection of pulmonary hypertension (Abst.)

Am J. Cardio., 41 436 .

Kalmanson D., Vegrate C., Abitbol G., (1979) :

Two dimensional echo Doppler velocimetry in mitral and tricuspid valve disease. In proceedings of the international symposium on Recent advance in ultrasound Diagnosis, edited by Kujak A., Dubrounik, Excerpta Medica, pp 335 - 348.

Keane J.F., Fyler D.C., Nadas A.S., (1978) :

Hazards of cardiac catheterization in children with primary pulmonary vascular obstruction.

Am : heart J., 96 : 556 - 558 .

Kisslo J., Van Ramm O.T. and Thursturn F., (1976) :

Cardiac imaging using a phased array ultrasound system II.
Clinical technique and applications circulation 53:262 - 267.

Knap and Meximoy M. (1977) :

Determinats of pulmonic valve opening velocity -
experimental echo cardiographic studies (abstract).
Circ. 53 and 54 . Suppl. II. 61.

Kuwako K., Tohda E., Ino T., et al ; (1980):

Echocardiographic evaluation of the tricuspid valve and
ring .
J. Cardiogr., 10 : 947 - 966.

L

Laurenceau I.L., Duesnil J.G., (1976):

Left ventricular dimensions as determinants of ventricular
septal motion.
Chest 69 : 388 .

Lew W., Karliner S.J., (1979) :

Assessment of pulmonary valve echogram in normal
subjects and in patients with pulmonary arterial
hypertension .
Br . Heart J. : 42 : 147 - 161 .

Lima C.O., Sahn D.J., Valdes -J.D, Allen H.D (1983) :

Non - invasive prediction of trans-valvular pressure gradient in patients with pulmonary stenosis by quantitative two dimensional echocardiographic Doppler studies.
Circ. 67 : 866 .

M

Marvin Kawabori I., Guntheroth W.G., (1985):

Validation of Doppler diagnosis of tricuspid regurgitation (abstract).
Circ. 64 (Suppl IV) : IV - 255 .

Milnor W.R., Conti C.R., Lewis K.B., et al ; (1969):

Pulmonary arterial pulse wave velocity and impedance in man.
Circ. Res. 25 : 637 - 49 .

Miyatake K., Okamoto M., Kinoshita W., et al ; (1982):

Evaluation of tricuspid regurgitation by pulsed Doppler and two - dimensional echocardiography .
Circ. 60 ; 777-784 .

Miyatake K., Oka moto M., Kinoshita W., et al ; (1982) :

Pulmonary regurgitation studied with ultrasonic pulsed Doppler technique .
Circ. 65 : 969 .

*Murphy, J (1980) : Persistent pulmonary hypertension of newborn
pediatr. Resp. 14:606*

*Murray G.B., (1971) :
Angiographic evaluation of valvular insufficiency .
Circ. 43 : 599 .*

N

*Nada N., Gramiak R., Robinson T.I., Shah P. M (1974):
Echocardiographic evaluation of pulmonary hypertension .
Circ., 50 : 575 .*

*Nishimura R.A., Jajik A.J., (1994):
Quantitative haemodynamics by Doppler echocardiography .
Prog. Cardiovasc. Dis. 35 (4) : 309 - 342.*

O

*Okamoto M., Miyatake K., Kinoshita, Nimura Y., (1984):
Analysis of blood flow in pulmonary hypertension with the
pulsed Doppler flowmeter combined with cross sectional
echocardiography .
Br. Heart J., 51 : 407 - 15.*

Omoto R., Kasiae C., (1986):

Basic principles of Doppler colour flow imaging .

Echocardiography 3 : 463 - 473.

P

Pocoski D.J., Shah P.M. and Sylvester L., (1978):

Physiologic Correlates of pulmonary valve motion in diastole (abst.)

Am J. Cardio , 41 : 436 .

Popp R.L., Fowles R. And Collart J., (1979) :

Cardiac anatomy viewed systematically with two dimensional echocardiography .

Chest, 75 , 579 - 585.

Pornin M., Morin D., Guize L., (1994):

Diagnosis, Coarse and Prognosis of pulmonary arterial hypertension .

Ann - Cardiol - Angeiol - Paris . 43 (8) 479 - 85.

R

Rabinovitch M., Reid L., (1981) :

Structural analysis of the pulmonary vascular bed in congenital heart Defects.

In Engle, M.E. (ed) : Pediatric cardiovascular disease.

Philadelphia, F.A. Davis pp 149 - 169 .

Requarth , Goldberg S.J., Vasko S.D., Allen H.D., (1984) :

In vitro verification of Doppler prediction of trans-valvular pressure gradient and orifice area in stenosis .

Am . J. Cardiol. 53 : 1369 .

Rice M.J., Seward J.B., Hagler D.J. et al., (1983) :

Impact of 2-dimensional echocardiography on the management of distressed newborns in whom cardiac disease is suspected.

Am. J. Cardiol 51 : 288 - 292.

Rich S., Dantzker D.R., Ayres S.M., (1987):

Primary pulmonary hypertension

Ann Intern Med. 107 : 216 - 223.

Riemnschnieder, J.A.,Nielsen H.C., Ruttenberg H.D.et al., (1976):

Disturbances of transitional circulation : Spectrum of pulmonary hypertesion and myocardial dysfunction

J. Pediatr. 89 : 622 - 625.

Rieves J.T., Groves B.M., Turkevich D., (1986):

The case for treatment of selective patients with primary pulmonary hypertension .

Am. Rev. respir Dis. 134 : 342 - 346.

S

Saal A.K., Otto C.M., Janko C.L., (1984):

Measurement of pulmonary systolic pressure in adults with tricuspid regurgitation using high pulsed repetition frequency. Doppler (abst.)

Circ.; 70 (Supp 11) 11 - 117.

Sharma S; Goswami- kc Shrivastavas (1994) :

Non- invasive estimation of pulmonary artery wedge pressure by dual M-Mode echocardiography.

J.Assoc Physicians - india 42 (4) : 298 - 9.

Siassi B., Goldberg S., Emmanouilides G.C. et al., (1986):

Persistent pulmonary hypertension in newborn infants

J. Pediatr 78 - 610-615.

Skinner J.R., Boys R.J., Hunter S., et al., (1991) :

Non- invasive assessment of pulmonary arterial pressure in healthy neonate .

Arch - Dis - child 66 (4) : 386 - 90 .

Skjaerpc T., Hatle L., (1983):

Non - invasive estimation of pulmonary artery pressure by Doppler ultrasound in tricuspid regurgitation. In : Sponcer MP, ed. Cardiac Doppler Diagnosis. Bostan : Martinus , Nijhoff, : 247 - 54.

Stephen P. And Sanders, M.D. (1992):

Echocardiography ; Nadas pediatric cardiology Edited by Fyler DC., Mosby year book inc., st-louis., Boston, Philadelphia , page 159 .

Stoermer J. Hentrich F., Galal D. et al., (1984) :

Risks der herztatheterisi emng and angiokardiographic in sauglings und kindersalter. Klin Pediatr . 196 : 191 - 194.

Sundler H., Dodge H.T., May R.C., (1963) :

Quantitation of valvular insufficiency in man by angiocardiography .

Am. Heart J., 65 : 501 .

Switzer, D.F. and Nada N.C., (1986):

Colour Doppler evaluation of the valvular regurgitation, echocardiography 2 : 543 - 553.

T

Tahara M., Tanaka H., Nakao (1981):

Haemodynamic Determinants of pulmonary valve motion during systole in experimental pulmonary hypertension .
Circ. 64 1249 - 1255.

Tavel, M.E., (1972):

Clinical phonocardiography and external pulse recording .
Chicago, year Book Medical . pp 274 - 278.

Tei C., Kisanuki A., Arina S., (1984) :

Non- invasive assessment of right ventricular pressure in patients with tricuspid regurgitation by continuous wave Doppler echocardiography .
Circ. 70 (Supp 11) 11- 116.

U

Ubago J.L., Figueroa A., Ochoteco, (1983):

Analysis of the amount of tricuspid valve annular dilatation required to produce functional tricuspid regurgitation.
Am . J. Card. 53 , 155-158 .

W

Waggoner A.D., Quinones M.A., Young T.B., (1981):

Pulsed Doppler echocardiographic detection of right sided valve regurgitation .

Am. J. Cardiol 47 279 - 66.

Wagenvoort C.A., (1984):

Lung biopsies and pulmonary vascular disease. In weir E.K., Raees J.T. (eds) : Pulmonary hypertension Mt. Kisco , NY, Future publishing

PP. 73 - 114.

Wells P.N.T., (1975):

Absorption and dispersion of ultrasound in biological tissue.

Ultrasound Med. Biol., 1 : 369 .

Wessel D.L., Kean J.F., Fellows K.E., et al., (1986) :

Fibrinolytic therapy for femoral arterial thrombosis following cardiac catheterization in infants and children.

Am. J. Cardiol 58 : 347 - 351.

Weymn A.E., Dillon J.C., Feigenhalm H., (1974b):

Echocardiographic patterns of pulmonic valve motion with pulmonary hypertension.

Circ 50 - 905 - 910 .

Weyman A.E., Dillon J.C., Feigenbaum H., (1974 a) :

Echocardiographic patterns of pulmonary valve motion in
valvular pulmonary stenosis.

Am. J. Cardiol 34 , 644 - 651.

Y

Yook P.G., Popp R.L., (1984) :

Non - invasive estimation of right ventricular pressures by
Doppler ultrasound in patients with tricuspid and aortic
regurgitation.

Circ. 68 (Supp 111) 111 - 230.

Z

Zellers, T. (1989):

persistant pulmonary hypertension of the newborn .

J. Pediatr. 114:735 .